



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 06:55 PM EST

PDB ID : 4WSR
Title : The crystal structure of hemagglutinin form A/chicken/New York/14677-13/1998
Authors : Yang, H.; Carney, P.J.; Chang, J.C.; Villanueva, J.M.; Stevens, J.
Deposited on : 2014-10-28
Resolution : 2.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	 6% 81% 11% • 6%
1	B	514	 8% 78% 13% • 6%
1	C	514	 6% 83% 9% • 6%
1	D	514	 9% 80% 16% ••
1	E	514	 8% 79% 14% •• 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	514	 17% 79% 11% 6%
2	G	2	 100%
2	H	2	 100%
2	I	2	 100%
2	J	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	I	2	-	-	-	X
3	NAG	A	601	X	-	-	-
3	NAG	B	601	X	-	-	-
3	NAG	C	605	X	-	-	-
3	NAG	D	601	X	-	-	-
3	NAG	D	602	X	-	-	-
3	NAG	E	601	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 24430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	483	3841	2410	669	743	19	0	0	0
1	B	482	3832	2405	667	741	19	0	0	0
1	C	482	3832	2405	667	741	19	0	0	0
1	E	489	3886	2441	675	751	19	0	0	0
1	D	498	3952	2484	688	761	19	0	0	0
1	F	481	3825	2400	666	740	19	0	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ALA	-	expression tag	UNP Q0A3A5
A	-2	ASP	-	expression tag	UNP Q0A3A5
A	-1	PRO	-	expression tag	UNP Q0A3A5
A	0	GLY	-	expression tag	UNP Q0A3A5
A	317	THR	UNK	conflict	UNP Q0A3A5
A	504	GLY	VAL	conflict	UNP Q0A3A5
A	505	ARG	MET	conflict	UNP Q0A3A5
A	507	VAL	-	expression tag	UNP Q0A3A5
A	508	PRO	-	expression tag	UNP Q0A3A5
A	509	ARG	-	expression tag	UNP Q0A3A5
A	510	GLY	-	expression tag	UNP Q0A3A5
B	-3	ALA	-	expression tag	UNP Q0A3A5
B	-2	ASP	-	expression tag	UNP Q0A3A5
B	-1	PRO	-	expression tag	UNP Q0A3A5
B	0	GLY	-	expression tag	UNP Q0A3A5
B	317	THR	UNK	conflict	UNP Q0A3A5
B	504	GLY	VAL	conflict	UNP Q0A3A5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	505	ARG	MET	conflict	UNP Q0A3A5
B	507	VAL	-	expression tag	UNP Q0A3A5
B	508	PRO	-	expression tag	UNP Q0A3A5
B	509	ARG	-	expression tag	UNP Q0A3A5
B	510	GLY	-	expression tag	UNP Q0A3A5
C	-3	ALA	-	expression tag	UNP Q0A3A5
C	-2	ASP	-	expression tag	UNP Q0A3A5
C	-1	PRO	-	expression tag	UNP Q0A3A5
C	0	GLY	-	expression tag	UNP Q0A3A5
C	317	THR	UNK	conflict	UNP Q0A3A5
C	504	GLY	VAL	conflict	UNP Q0A3A5
C	505	ARG	MET	conflict	UNP Q0A3A5
C	507	VAL	-	expression tag	UNP Q0A3A5
C	508	PRO	-	expression tag	UNP Q0A3A5
C	509	ARG	-	expression tag	UNP Q0A3A5
C	510	GLY	-	expression tag	UNP Q0A3A5
E	-3	ALA	-	expression tag	UNP Q0A3A5
E	-2	ASP	-	expression tag	UNP Q0A3A5
E	-1	PRO	-	expression tag	UNP Q0A3A5
E	0	GLY	-	expression tag	UNP Q0A3A5
E	317	THR	UNK	conflict	UNP Q0A3A5
E	504	GLY	VAL	conflict	UNP Q0A3A5
E	505	ARG	MET	conflict	UNP Q0A3A5
E	507	VAL	-	expression tag	UNP Q0A3A5
E	508	PRO	-	expression tag	UNP Q0A3A5
E	509	ARG	-	expression tag	UNP Q0A3A5
E	510	GLY	-	expression tag	UNP Q0A3A5
D	-3	ALA	-	expression tag	UNP Q0A3A5
D	-2	ASP	-	expression tag	UNP Q0A3A5
D	-1	PRO	-	expression tag	UNP Q0A3A5
D	0	GLY	-	expression tag	UNP Q0A3A5
D	317	THR	UNK	conflict	UNP Q0A3A5
D	504	GLY	VAL	conflict	UNP Q0A3A5
D	505	ARG	MET	conflict	UNP Q0A3A5
D	507	VAL	-	expression tag	UNP Q0A3A5
D	508	PRO	-	expression tag	UNP Q0A3A5
D	509	ARG	-	expression tag	UNP Q0A3A5
D	510	GLY	-	expression tag	UNP Q0A3A5
F	-3	ALA	-	expression tag	UNP Q0A3A5
F	-2	ASP	-	expression tag	UNP Q0A3A5
F	-1	PRO	-	expression tag	UNP Q0A3A5
F	0	GLY	-	expression tag	UNP Q0A3A5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	317	THR	UNK	conflict	UNP Q0A3A5
F	504	GLY	VAL	conflict	UNP Q0A3A5
F	505	ARG	MET	conflict	UNP Q0A3A5
F	507	VAL	-	expression tag	UNP Q0A3A5
F	508	PRO	-	expression tag	UNP Q0A3A5
F	509	ARG	-	expression tag	UNP Q0A3A5
F	510	GLY	-	expression tag	UNP Q0A3A5

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	2	28	16	2	10	0	0	0
2	H	2	28	16	2	10	0	0	0
2	I	2	28	16	2	10	0	0	0
2	J	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	B	1	Total 14	C 8	N 1	O 5	0	0
3	C	1	Total 14	C 8	N 1	O 5	0	0
3	C	1	Total 14	C 8	N 1	O 5	0	0
3	C	1	Total 14	C 8	N 1	O 5	0	0
3	E	1	Total 14	C 8	N 1	O 5	0	0
3	E	1	Total 14	C 8	N 1	O 5	0	0
3	D	1	Total 14	C 8	N 1	O 5	0	0
3	D	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	226	Total 226	O 226	0	0
4	B	219	Total 219	O 219	0	0
4	C	175	Total 175	O 175	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	85	Total O 85 85	0	0
4	D	212	Total O 212 212	0	0
4	F	107	Total O 107 107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin




- Molecule 1: Hemagglutinin





- Molecule 1: Hemagglutinin








• Molecule 1: Hemagglutinin




• Molecule 1: Hemagglutinin



• Molecule 1: Hemagglutinin





- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	252.54Å 134.82Å 123.09Å 90.00° 113.23° 90.00°	Depositor
Resolution (Å)	48.00 – 2.50 47.29 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.9 (48.00-2.50) 98.9 (47.29-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.65 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.210 , 0.246 0.213 , 0.247	Depositor DCC
R_{free} test set	6504 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	49.1	Xtrriage
Anisotropy	0.338	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24430	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.73	2/3928 (0.1%)	0.80	5/5323 (0.1%)
1	B	0.70	0/3919	0.83	9/5311 (0.2%)
1	C	0.70	0/3919	0.81	6/5311 (0.1%)
1	D	0.68	1/4042 (0.0%)	0.82	8/5477 (0.1%)
1	E	0.64	0/3974	0.82	11/5385 (0.2%)
1	F	0.67	1/3911 (0.0%)	0.86	12/5299 (0.2%)
All	All	0.69	4/23693 (0.0%)	0.82	51/32106 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	60	TRP	CB-CG	-5.61	1.40	1.50
1	F	485	TYR	N-CA	5.58	1.57	1.46
1	A	109	ASP	CB-CG	-5.22	1.40	1.51
1	D	109	ASP	CB-CG	-5.13	1.41	1.51

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	201	MET	CG-SD-CE	9.91	116.06	100.20
1	F	156	ASP	CB-CG-OD2	9.56	126.91	118.30
1	F	156	ASP	CB-CG-OD1	-9.17	110.05	118.30
1	F	201	MET	CG-SD-CE	8.19	113.30	100.20
1	F	461	LEU	CA-CB-CG	7.27	132.03	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3841	0	3701	51	0
1	B	3832	0	3693	51	0
1	C	3832	0	3691	32	0
1	D	3952	0	3819	62	1
1	E	3886	0	3746	72	0
1	F	3825	0	3687	63	0
2	G	28	0	25	0	0
2	H	28	0	25	0	0
2	I	28	0	25	0	0
2	J	28	0	25	0	0
3	A	14	0	13	0	0
3	B	14	0	13	1	0
3	C	42	0	39	2	0
3	D	28	0	26	0	0
3	E	28	0	26	0	0
4	A	226	0	0	4	0
4	B	219	0	0	8	0
4	C	175	0	0	4	0
4	D	212	0	0	14	0
4	E	85	0	0	15	0
4	F	107	0	0	1	0
All	All	24430	0	22554	299	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 299 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:145:ARG:O	1:E:252:TRP:NE1	1.57	1.35
1:F:480:VAL:HG23	1:F:485:TYR:OH	1.47	1.13
1:F:476:CYS:O	1:F:485:TYR:OH	1.75	1.04
1:A:447:HIS:CE1	1:A:451:LYS:NZ	2.28	1.01
1:E:241:ASN:O	4:E:701:HOH:O	1.86	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:137:TYR:OH	1:D:137:TYR:OH[2_555]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	479/514 (93%)	464 (97%)	14 (3%)	1 (0%)	47 68
1	B	478/514 (93%)	464 (97%)	13 (3%)	1 (0%)	47 68
1	C	478/514 (93%)	465 (97%)	12 (2%)	1 (0%)	47 68
1	D	496/514 (96%)	479 (97%)	15 (3%)	2 (0%)	34 54
1	E	485/514 (94%)	472 (97%)	12 (2%)	1 (0%)	47 68
1	F	477/514 (93%)	461 (97%)	14 (3%)	2 (0%)	34 54
All	All	2893/3084 (94%)	2805 (97%)	80 (3%)	8 (0%)	41 61

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	486	ASN
1	D	497	ASN
1	D	346	VAL
1	A	346	VAL
1	E	346	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/443 (95%)	401 (95%)	20 (5%)	25	48
1	B	420/443 (95%)	396 (94%)	24 (6%)	20	39
1	C	420/443 (95%)	402 (96%)	18 (4%)	29	53
1	D	431/443 (97%)	409 (95%)	22 (5%)	24	45
1	E	425/443 (96%)	410 (96%)	15 (4%)	36	62
1	F	419/443 (95%)	399 (95%)	20 (5%)	25	48
All	All	2536/2658 (95%)	2417 (95%)	119 (5%)	26	49

5 of 119 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	294	GLN
1	F	294	GLN
1	E	301	ILE
1	F	217	ARG
1	F	490	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	247	ASN
1	E	445	ASN
1	F	354	HIS
1	D	8	HIS
1	B	171	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

17 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	601	1	14,14,15	0.67	0	17,19,21	1.63	5 (29%)
2	NAG	I	1	1,2	14,14,15	0.83	1 (7%)	17,19,21	1.08	1 (5%)
3	NAG	C	602	1	14,14,15	0.73	1 (7%)	17,19,21	1.76	4 (23%)
2	NAG	I	2	2	14,14,15	0.69	0	17,19,21	1.93	3 (17%)
3	NAG	D	601	1	14,14,15	0.75	0	17,19,21	2.08	5 (29%)
3	NAG	C	601	1	14,14,15	0.99	1 (7%)	17,19,21	3.10	6 (35%)
3	NAG	E	601	1	14,14,15	0.80	0	17,19,21	2.49	6 (35%)
3	NAG	D	602	1	14,14,15	0.63	0	17,19,21	1.79	2 (11%)
3	NAG	C	605	1	14,14,15	1.00	1 (7%)	17,19,21	1.69	5 (29%)
3	NAG	B	601	1	14,14,15	0.63	0	17,19,21	2.68	7 (41%)
2	NAG	J	1	1,2	14,14,15	0.90	1 (7%)	17,19,21	1.56	4 (23%)
2	NAG	J	2	2	14,14,15	0.66	0	17,19,21	2.11	3 (17%)
2	NAG	G	2	2	14,14,15	0.63	0	17,19,21	1.20	2 (11%)
2	NAG	H	1	1,2	14,14,15	0.70	0	17,19,21	1.91	4 (23%)
3	NAG	E	602	1	14,14,15	0.50	0	17,19,21	1.56	1 (5%)
2	NAG	H	2	2	14,14,15	0.75	0	17,19,21	1.51	2 (11%)
2	NAG	G	1	1,2	14,14,15	0.76	1 (7%)	17,19,21	2.07	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	H	2	2	-	1/6/23/26	0/1/1/1
3	NAG	A	601	1	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	D	601	1	1/1/5/7	1/6/23/26	0/1/1/1
3	NAG	C	602	1	-	0/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	2/6/23/26	0/1/1/1
3	NAG	D	602	1	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	C	601	1	-	4/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
3	NAG	C	605	1	1/1/5/7	3/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	601	1	1/1/5/7	1/6/23/26	0/1/1/1
2	NAG	J	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	2/6/23/26	0/1/1/1
3	NAG	E	602	1	-	2/6/23/26	0/1/1/1
3	NAG	E	601	1	1/1/5/7	4/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	601	NAG	C1-C2	2.66	1.56	1.52
2	I	1	NAG	O5-C1	-2.23	1.40	1.43
3	C	602	NAG	C1-C2	2.22	1.55	1.52
2	G	1	NAG	C1-C2	2.16	1.55	1.52
3	C	605	NAG	C1-C2	2.08	1.55	1.52

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	NAG	C1-O5-C5	9.61	125.21	112.19
3	B	601	NAG	C1-O5-C5	8.45	123.65	112.19
3	D	602	NAG	O5-C1-C2	-6.29	101.36	111.29
2	J	2	NAG	C1-O5-C5	6.26	120.67	112.19
2	I	2	NAG	C1-O5-C5	6.00	120.32	112.19

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	601	NAG	C1
3	B	601	NAG	C1
3	C	605	NAG	C1
3	E	601	NAG	C1
3	D	601	NAG	C1

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	601	NAG	C3-C2-N2-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	601	NAG	C1-C2-N2-C7
2	J	1	NAG	O5-C5-C6-O6
3	C	601	NAG	O5-C5-C6-O6
2	I	1	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	601	NAG	1	0
3	C	605	NAG	1	0
3	B	601	NAG	1	0

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	G	1	1,2	14,14,15	0.76	1 (7%)	17,19,21	2.07	6 (35%)
2	NAG	G	2	2	14,14,15	0.63	0	17,19,21	1.20	2 (11%)
2	NAG	H	1	1,2	14,14,15	0.70	0	17,19,21	1.91	4 (23%)
2	NAG	H	2	2	14,14,15	0.75	0	17,19,21	1.51	2 (11%)
2	NAG	I	1	1,2	14,14,15	0.83	1 (7%)	17,19,21	1.08	1 (5%)
2	NAG	I	2	2	14,14,15	0.69	0	17,19,21	1.93	3 (17%)
2	NAG	J	1	1,2	14,14,15	0.90	1 (7%)	17,19,21	1.56	4 (23%)
2	NAG	J	2	2	14,14,15	0.66	0	17,19,21	2.11	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	H	2	2	-	1/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
2	NAG	J	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	1	NAG	O5-C1	-2.23	1.40	1.43
2	G	1	NAG	C1-C2	2.16	1.55	1.52
2	J	1	NAG	C1-C2	2.05	1.55	1.52

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	2	NAG	C1-O5-C5	6.26	120.67	112.19
2	I	2	NAG	C1-O5-C5	6.00	120.32	112.19
2	G	1	NAG	C1-C2-N2	4.29	117.82	110.49
2	H	2	NAG	C4-C3-C2	4.20	117.17	111.02
2	H	2	NAG	C3-C4-C5	3.70	116.85	110.24

There are no chirality outliers.


5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	J	1	NAG	O5-C5-C6-O6
2	I	1	NAG	O5-C5-C6-O6
2	J	1	NAG	C4-C5-C6-O6
2	G	2	NAG	O5-C5-C6-O6
2	I	1	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	601	1	14,14,15	0.67	0	17,19,21	1.63	5 (29%)
3	NAG	D	601	1	14,14,15	0.75	0	17,19,21	2.08	5 (29%)
3	NAG	C	602	1	14,14,15	0.73	1 (7%)	17,19,21	1.76	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	602	1	14,14,15	0.63	0	17,19,21	1.79	2 (11%)
3	NAG	C	601	1	14,14,15	0.99	1 (7%)	17,19,21	3.10	6 (35%)
3	NAG	C	605	1	14,14,15	1.00	1 (7%)	17,19,21	1.69	5 (29%)
3	NAG	B	601	1	14,14,15	0.63	0	17,19,21	2.68	7 (41%)
3	NAG	E	602	1	14,14,15	0.50	0	17,19,21	1.56	1 (5%)
3	NAG	E	601	1	14,14,15	0.80	0	17,19,21	2.49	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	D	601	1	1/1/5/7	1/6/23/26	0/1/1/1
3	NAG	C	602	1	-	0/6/23/26	0/1/1/1
3	NAG	D	602	1	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	C	601	1	-	4/6/23/26	0/1/1/1
3	NAG	C	605	1	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	B	601	1	1/1/5/7	1/6/23/26	0/1/1/1
3	NAG	E	602	1	-	2/6/23/26	0/1/1/1
3	NAG	E	601	1	1/1/5/7	4/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	601	NAG	C1-C2	2.66	1.56	1.52
3	C	602	NAG	C1-C2	2.22	1.55	1.52
3	C	605	NAG	C1-C2	2.08	1.55	1.52

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	NAG	C1-O5-C5	9.61	125.21	112.19
3	B	601	NAG	C1-O5-C5	8.45	123.65	112.19
3	D	602	NAG	O5-C1-C2	-6.29	101.36	111.29
3	D	601	NAG	O5-C1-C2	-5.81	102.12	111.29
3	C	601	NAG	O5-C5-C6	5.72	116.16	107.20

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	601	NAG	C1
3	B	601	NAG	C1
3	C	605	NAG	C1
3	E	601	NAG	C1
3	D	601	NAG	C1

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	601	NAG	C3-C2-N2-C7
3	C	601	NAG	C1-C2-N2-C7
3	C	601	NAG	O5-C5-C6-O6
3	A	601	NAG	O5-C5-C6-O6
3	C	605	NAG	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	601	NAG	1	0
3	C	605	NAG	1	0
3	B	601	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/514 (93%)	0.18	29 (6%) 21 22	29, 47, 101, 126	0
1	B	482/514 (93%)	0.45	43 (8%) 9 9	30, 51, 125, 173	0
1	C	482/514 (93%)	0.24	33 (6%) 17 17	31, 53, 104, 133	0
1	D	498/514 (96%)	0.55	48 (9%) 8 7	31, 59, 151, 182	0
1	E	489/514 (95%)	0.53	42 (8%) 10 10	39, 73, 112, 140	0
1	F	481/514 (93%)	0.79	86 (17%) 1 1	36, 72, 146, 195	0
All	All	2915/3084 (94%)	0.46	281 (9%) 8 7	29, 60, 126, 195	0

The worst 5 of 281 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	3	ILE	9.0
1	D	456	ASP	7.5
1	D	469	TRP	7.0
1	B	485	TYR	6.9
1	F	361	GLY	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	602	14/15	0.62	0.34	102,130,143,148	0
3	NAG	D	601	14/15	0.72	0.33	127,140,146,151	0
3	NAG	B	601	14/15	0.73	0.24	117,125,140,143	0
2	NAG	I	2	14/15	0.76	0.41	105,124,129,130	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	C	601	14/15	0.78	0.17	99,105,113,114	0
3	NAG	C	602	14/15	0.81	0.26	106,118,123,131	0
2	NAG	J	2	14/15	0.82	0.25	86,95,101,102	0
2	NAG	H	2	14/15	0.82	0.27	104,115,119,121	0
3	NAG	E	602	14/15	0.82	0.16	101,111,115,119	0
3	NAG	A	601	14/15	0.83	0.24	87,105,115,115	0
3	NAG	E	601	14/15	0.83	0.22	61,88,96,124	0
2	NAG	G	2	14/15	0.84	0.29	87,96,100,105	0
2	NAG	I	1	14/15	0.84	0.18	95,108,115,124	0
3	NAG	C	605	14/15	0.86	0.23	61,80,98,109	0
2	NAG	G	1	14/15	0.91	0.29	63,71,78,86	0
2	NAG	H	1	14/15	0.91	0.25	78,86,92,99	0
2	NAG	J	1	14/15	0.91	0.13	60,72,76,81	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	I	2	14/15	0.76	0.41	105,124,129,130	0
2	NAG	H	2	14/15	0.82	0.27	104,115,119,121	0
2	NAG	J	2	14/15	0.82	0.25	86,95,101,102	0
2	NAG	G	2	14/15	0.84	0.29	87,96,100,105	0
2	NAG	I	1	14/15	0.84	0.18	95,108,115,124	0
2	NAG	G	1	14/15	0.91	0.29	63,71,78,86	0
2	NAG	J	1	14/15	0.91	0.13	60,72,76,81	0
2	NAG	H	1	14/15	0.91	0.25	78,86,92,99	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	602	14/15	0.62	0.34	102,130,143,148	0
3	NAG	D	601	14/15	0.72	0.33	127,140,146,151	0
3	NAG	B	601	14/15	0.73	0.24	117,125,140,143	0
3	NAG	C	601	14/15	0.78	0.17	99,105,113,114	0
3	NAG	C	602	14/15	0.81	0.26	106,118,123,131	0
3	NAG	E	602	14/15	0.82	0.16	101,111,115,119	0
3	NAG	E	601	14/15	0.83	0.22	61,88,96,124	0
3	NAG	A	601	14/15	0.83	0.24	87,105,115,115	0
3	NAG	C	605	14/15	0.86	0.23	61,80,98,109	0

6.5 Other polymers [i](#)

There are no such residues in this entry.