



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2023 – 11:15 pm GMT

PDB ID : 2X1D
Title : The crystal structure of mature acyl coenzyme A:isopenicillin N acyltransferase from *Penicillium chrysogenum*
Authors : Bokhove, M.; Yoshida, H.; Hensgens, C.M.H.; van der Laan, J.M.; Sutherland, J.D.; Dijkstra, B.W.
Deposited on : 2009-12-23
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 4% 95% 5% 5%
1	B	357	 7% 91% 8% 5%
1	C	357	 8% 93% 5% 5%
1	D	357	 4% 94% 5% 5%

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 12327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACYL-COENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	354	Total 2848	C 1784	N 513	O 538	S 13	0	8	0
1	B	352	Total 2856	C 1790	N 513	O 538	S 15	0	10	0
1	C	351	Total 2835	C 1777	N 509	O 534	S 15	0	8	0
1	D	351	Total 2828	C 1776	N 503	O 532	S 17	0	8	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	2	Total Cl 2 2	0	0
4	C	1	Total Cl 1 1	0	0
4	D	1	Total Cl 1 1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is (4S,5S)-1,2-DITHIANE-4,5-DIOL (three-letter code: D1D) (formula: C₄H₈O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	202	Total	O	0	0
			202	202		
7	B	260	Total	O	0	0
			260	260		
7	C	226	Total	O	0	0
			226	226		
7	D	221	Total	O	0	0
			221	221		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ACYL-COENZYME




- Molecule 1: ACYL-COENZYME



- Molecule 1: ACYL-COENZYME



- Molecule 1: ACYL-COENZYME



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	198.71Å 68.22Å 146.96Å 90.00° 128.64° 90.00°	Depositor
Resolution (Å)	34.10 – 1.64 33.81 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.2 (34.10-1.64) 98.2 (33.81-1.64)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.64Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.179 , 0.207 0.190 , 0.215	Depositor DCC
R_{free} test set	9288 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12327	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, D1D, GOL, CSD, CL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2898	0.62	0/3913
1	B	0.56	0/2905	0.66	0/3923
1	C	0.53	0/2884	0.64	0/3891
1	D	0.49	0/2877	0.61	0/3884
All	All	0.52	0/11564	0.63	0/15611

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2848	0	2785	9	0
1	B	2856	0	2799	22	0
1	C	2835	0	2777	8	0
1	D	2828	0	2774	12	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	D	10	0	0	0	0
3	A	6	0	8	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	4	0	3	1	0
5	C	4	0	3	0	0
5	D	4	0	3	0	0
6	D	8	0	8	0	0
7	A	202	0	0	0	0
7	B	260	0	0	1	0
7	C	226	0	0	1	0
7	D	221	0	0	0	0
All	All	12327	0	11160	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:TRP:CZ2	1:B:179[B]:VAL:HG12	2.02	0.95
1:B:62:TRP:HZ2	1:B:179[B]:VAL:HG12	1.38	0.86
1:C:94:TYR:HA	1:C:97:LYS:HB2	1.70	0.73
1:C:96:LEU:HG	1:C:174:LEU:HD22	1.73	0.70
1:B:256:GLU:OE2	5:B:1358:ACT:H2	1.92	0.69
1:B:179[B]:VAL:HG11	1:B:204:GLN:HB3	1.75	0.66
1:B:177[B]:THR:O	1:B:177[B]:THR:HG22	1.97	0.64
1:B:179[B]:VAL:HG11	1:B:204:GLN:CB	2.28	0.64
1:A:236:LEU:HD23	1:A:242:MET:HG3	1.80	0.63
1:D:179[B]:VAL:HG11	1:D:204:GLN:CB	2.30	0.62
1:A:95:GLY:HA2	1:A:174:LEU:CD1	2.34	0.56
1:B:179[B]:VAL:CG1	1:B:204:GLN:HB3	2.37	0.55
1:B:236:LEU:HD23	1:B:242[B]:MET:HG3	1.87	0.55
1:C:42:LYS:HE2	1:C:96:LEU:HD22	1.92	0.51
1:A:203:GLU:HG3	1:B:232:ARG:HD2	1.92	0.51
1:A:350:ARG:HG3	1:A:355:ALA:HB3	1.93	0.50
1:C:350:ARG:HG3	1:C:355:ALA:HB3	1.93	0.50
1:B:207:MET:HG3	1:B:226:PHE:CD1	2.47	0.50
1:D:67:GLU:OE1	1:D:70:ARG:NH2	2.42	0.50
1:B:60[A]:GLU:HG3	7:B:2061:HOH:O	2.12	0.49
1:A:252:HIS:HB3	1:A:256:GLU:HG3	1.95	0.49
1:A:95:GLY:HA2	1:A:174:LEU:HD13	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:45:GLU:O	1:C:49:GLN:HG2	2.14	0.48
1:B:94:TYR:HA	1:B:97:LYS:HB2	1.96	0.47
1:B:112:ASN:HB3	7:C:2112:HOH:O	2.14	0.46
1:A:36:ILE:HG23	1:A:96:LEU:CD1	2.45	0.46
1:D:241:ARG:HD2	1:D:276:LEU:HD13	1.98	0.46
1:D:275:LEU:HB3	1:D:289:LEU:HD22	1.97	0.46
1:D:252:HIS:HB3	1:D:256:GLU:HG3	1.98	0.46
1:B:144:LYS:HE3	1:B:343[A]:ARG:O	2.16	0.45
1:B:177[B]:THR:O	1:B:177[B]:THR:CG2	2.64	0.45
1:B:133:LEU:HD12	1:B:145:PHE:CZ	2.52	0.45
1:D:179[B]:VAL:HG11	1:D:204:GLN:HB2	1.98	0.44
1:D:236:LEU:HA	1:D:242[B]:MET:HG3	2.00	0.44
1:B:179[B]:VAL:HG23	1:B:179[B]:VAL:O	2.19	0.43
1:C:215:VAL:HB	1:C:222:PHE:CE1	2.53	0.43
1:B:49:GLN:O	1:B:53[B]:GLN:HG3	2.18	0.43
1:D:179[B]:VAL:CG1	1:D:204:GLN:HB3	2.48	0.43
1:B:261:PRO:HG2	1:B:266:TRP:HH2	1.84	0.43
1:B:42:LYS:HE3	1:B:96:LEU:HD22	2.00	0.43
1:C:36:ILE:HG12	1:C:92:PHE:HB3	1.99	0.43
1:D:236:LEU:HD23	1:D:242[B]:MET:HG3	2.00	0.42
1:C:232:ARG:NE	1:C:251:GLN:O	2.49	0.42
1:A:211:ALA:HB3	1:A:226:PHE:CZ	2.55	0.41
1:A:160:ALA:HA	1:D:112:ASN:HA	2.01	0.41
1:B:179[B]:VAL:HG11	1:B:204:GLN:HG3	2.02	0.41
1:B:258:GLU:HG2	1:B:261:PRO:HG3	2.03	0.41
1:D:133:LEU:HD12	1:D:145:PHE:CZ	2.55	0.41
1:D:179[B]:VAL:O	1:D:179[B]:VAL:CG2	2.69	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/357 (100%)	346 (97%)	12 (3%)	0	100	100
1	B	358/357 (100%)	349 (98%)	8 (2%)	1 (0%)	41	21
1	C	355/357 (99%)	347 (98%)	7 (2%)	1 (0%)	41	21
1	D	354/357 (99%)	345 (98%)	8 (2%)	1 (0%)	41	21
All	All	1425/1428 (100%)	1387 (97%)	35 (2%)	3 (0%)	47	26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	154	LYS
1	D	154	LYS
1	C	154	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/291 (102%)	296 (100%)	1 (0%)	92	87
1	B	298/291 (102%)	297 (100%)	1 (0%)	92	87
1	C	296/291 (102%)	294 (99%)	2 (1%)	84	71
1	D	295/291 (101%)	294 (100%)	1 (0%)	92	87
All	All	1186/1164 (102%)	1181 (100%)	5 (0%)	91	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	157	PHE
1	B	157	PHE
1	C	44	ASP
1	C	157	PHE
1	D	157	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	C	53	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	A	103	1	3,7,8	1.24	0	1,8,10	0.80	0
1	CSD	B	103	1	3,7,8	1.31	0	1,8,10	2.16	1 (100%)
1	CSD	D	103	1	3,7,8	1.27	0	1,8,10	0.69	0
1	CSD	C	103	1	3,7,8	1.23	0	1,8,10	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	103	1	-	0/2/6/8	-
1	CSD	B	103	1	-	0/2/6/8	-
1	CSD	D	103	1	-	0/2/6/8	-
1	CSD	C	103	1	-	0/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	103	CSD	OD1-SG-CB	2.16	109.64	105.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	B	1358	-	3,3,3	0.69	0	3,3,3	1.12	0
2	PO4	D	1356	-	4,4,4	0.79	0	6,6,6	0.71	0
6	D1D	D	1359	-	6,8,8	0.53	0	6,10,10	0.95	0
2	PO4	D	1357	-	4,4,4	0.81	0	6,6,6	0.39	0
3	GOL	A	1357	-	5,5,5	0.36	0	5,5,5	0.13	0
5	ACT	D	1358	-	3,3,3	0.83	0	3,3,3	1.04	0
2	PO4	A	1356	-	4,4,4	0.45	0	6,6,6	0.68	0
5	ACT	C	1357	-	3,3,3	0.71	0	3,3,3	1.33	0
2	PO4	B	1357	-	4,4,4	0.85	0	6,6,6	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1357	-	-	0/4/4/4	-
6	D1D	D	1359	-	-	-	0/0/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1358	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/357 (98%)	0.12	14 (3%) 38 36	17, 26, 40, 63	0
1	B	351/357 (98%)	0.25	26 (7%) 14 13	16, 23, 41, 53	0
1	C	350/357 (98%)	0.28	29 (8%) 11 9	16, 24, 49, 66	0
1	D	350/357 (98%)	0.13	15 (4%) 35 32	18, 26, 39, 47	0
All	All	1404/1428 (98%)	0.20	84 (5%) 21 19	16, 25, 42, 66	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	307	GLY	7.7
1	C	254	LYS	6.5
1	C	356	ARG	6.3
1	B	304	TYR	6.2
1	C	304	TYR	5.9
1	C	307	GLY	5.7
1	B	306	GLU	5.7
1	B	98	ALA	5.3
1	A	100	ARG	5.0
1	B	41	LYS	4.9
1	B	310	ARG	4.8
1	C	260	ASP	4.8
1	C	43	THR	4.6
1	D	41	LYS	4.6
1	C	308	LYS	4.6
1	C	259	LEU	4.4
1	C	310	ARG	4.4
1	C	44	ASP	4.4
1	B	94	TYR	4.3
1	A	101	ASP	4.2
1	D	307	GLY	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	254	LYS	4.1
1	B	260	ASP	3.9
1	C	251	GLN	3.8
1	C	306	GLU	3.8
1	B	356	ARG	3.6
1	C	253	GLY	3.6
1	D	98	ALA	3.4
1	C	49	GLN	3.4
1	D	306	GLU	3.4
1	C	46	GLU	3.4
1	C	96	LEU	3.3
1	C	262	LEU	3.3
1	A	251	GLN	3.2
1	C	45	GLU	3.2
1	B	97	LYS	3.1
1	C	42	LYS	3.1
1	A	56	ARG	3.1
1	C	41	LYS	3.1
1	B	251	GLN	3.1
1	A	41	LYS	3.0
1	B	185	LEU	3.0
1	A	98	ALA	2.9
1	D	277	ASP	2.9
1	A	277	ASP	2.9
1	C	120	TRP	2.9
1	D	15	TYR	2.8
1	C	305	GLU	2.8
1	B	257	LYS	2.8
1	C	97	LYS	2.8
1	C	47	LEU	2.7
1	B	308	LYS	2.7
1	C	257	LYS	2.7
1	C	309	SER	2.7
1	C	355	ALA	2.7
1	B	111	PRO	2.7
1	D	308	LYS	2.7
1	D	355	ALA	2.6
1	B	277	ASP	2.6
1	A	99	ALA	2.5
1	A	254	LYS	2.5
1	C	238	ALA	2.5
1	B	259	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	120	TRP	2.3
1	C	298	PHE	2.3
1	A	164[A]	VAL	2.3
1	B	298	PHE	2.3
1	B	309	SER	2.2
1	D	354	ASN	2.2
1	B	120	TRP	2.2
1	D	350	ARG	2.1
1	B	164	VAL	2.1
1	D	97	LYS	2.1
1	B	181[A]	SER	2.1
1	B	305	GLU	2.1
1	D	251	GLN	2.1
1	A	15	TYR	2.1
1	A	155	VAL	2.1
1	D	238	ALA	2.1
1	B	112	ASN	2.1
1	A	238	ALA	2.0
1	B	166	TYR	2.0
1	D	185	LEU	2.0
1	D	274	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	103	8/9	0.97	0.09	20,22,25,26	0
1	CSD	C	103	8/9	0.97	0.09	21,23,29,30	0
1	CSD	D	103	8/9	0.98	0.08	20,21,26,26	0
1	CSD	B	103	8/9	0.99	0.09	19,21,26,30	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	A	1358	1/1	0.69	0.09	69,69,69,69	0
2	PO4	D	1357	5/5	0.74	0.34	56,56,57,58	0
5	ACT	B	1358	4/4	0.75	0.20	34,35,36,36	0
5	ACT	D	1358	4/4	0.81	0.21	37,38,38,38	0
5	ACT	C	1357	4/4	0.86	0.17	30,31,31,31	0
2	PO4	A	1356	5/5	0.91	0.14	27,28,33,34	0
4	CL	B	1359	1/1	0.91	0.08	56,56,56,56	0
4	CL	B	1360	1/1	0.91	0.09	32,32,32,32	1
4	CL	D	1360	1/1	0.92	0.12	52,52,52,52	0
3	GOL	A	1357	6/6	0.92	0.16	27,29,30,31	0
6	D1D	D	1359	8/8	0.92	0.22	25,26,28,28	8
4	CL	C	1358	1/1	0.95	0.13	48,48,48,48	0
2	PO4	B	1357	5/5	0.96	0.11	29,31,32,33	0
2	PO4	D	1356	5/5	0.98	0.14	23,25,27,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.