



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 04:05 AM EDT

PDB ID : 6X1I
Title : Two-Component D3 Assembly Constructed by Fusing Symmetric Oligomers to Coiled Coils
Authors : Laniado, J.; Yeates, T.O.; Sawaya, M.R.
Deposited on : 2020-05-18
Resolution : 4.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1018 (4.84-3.80)
Clashscore	141614	1081 (4.84-3.80)
Ramachandran outliers	138981	1033 (4.84-3.80)
Sidechain outliers	138945	1016 (4.84-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	172	 82% 17% .
2	B	157	 67% 25% 7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cob_adeno_trans domain-containing protein PH0671 fused to a coiled coil.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	171	1354	861	226	263	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	initiating methionine	UNP O58404

- Molecule 2 is a protein called SnoaL-like Protein fused to a coiled coil.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	146	1153	726	214	211	2	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	221	HIS	-	expression tag	UNP Q7D0S4
B	222	HIS	-	expression tag	UNP Q7D0S4
B	223	HIS	-	expression tag	UNP Q7D0S4
B	224	HIS	-	expression tag	UNP Q7D0S4
B	225	HIS	-	expression tag	UNP Q7D0S4
B	226	HIS	-	expression tag	UNP Q7D0S4

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 3 2	Depositor
Cell constants a, b, c, α , β , γ	146.34Å 146.34Å 146.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.49 – 4.32 84.49 – 4.32	Depositor EDS
% Data completeness (in resolution range)	99.5 (84.49-4.32) 90.8 (84.49-4.32)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 4.30Å)	Xtrriage
Refinement program	PHENIX dev 3724	Depositor
R, R_{free}	0.278 , 0.293 0.277 , 0.293	Depositor DCC
R_{free} test set	392 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	217.9	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 330.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	2507	wwPDB-VP
Average B, all atoms (Å ²)	267.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1368	0.54	0/1837
2	B	0.29	0/1175	0.53	0/1584
All	All	0.29	0/2543	0.54	0/3421

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1354	0	1393	17	0
2	B	1153	0	1152	31	0
All	All	2507	0	2545	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:GLN:HA	1:A:170:LYS:HE3	1.70	0.74
1:A:104:VAL:HG21	1:A:158:ARG:HD2	1.76	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:203:ASN:HB2	2:B:209:ASP:HB3	1.78	0.64
1:A:33:THR:HG21	1:A:124:ARG:HG3	1.80	0.63
2:B:76:LYS:O	2:B:80:LEU:HD23	2.03	0.58
1:A:27:ILE:HD13	1:A:132:VAL:HG22	1.88	0.56
2:B:200:VAL:HG22	2:B:211:VAL:HG22	1.88	0.55
2:B:123:ARG:HH11	2:B:154:ILE:HG22	1.70	0.55
2:B:136:ARG:HG3	2:B:210:ASN:OD1	2.08	0.53
2:B:158:ARG:NH1	2:B:164:TYR:O	2.42	0.52
2:B:132:ALA:HA	2:B:207:ARG:NE	2.25	0.52
1:A:165:ALA:O	1:A:169:GLU:HG3	2.09	0.51
2:B:90:LYS:O	2:B:94:LEU:HB2	2.11	0.51
2:B:72:GLN:O	2:B:76:LYS:HG2	2.11	0.51
2:B:123:ARG:HH12	2:B:158:ARG:HB2	1.74	0.50
1:A:33:THR:HG22	1:A:125:ALA:HB2	1.93	0.49
2:B:127:VAL:HA	2:B:131:TRP:CE3	2.46	0.49
1:A:92:SER:O	1:A:96:GLU:HG3	2.13	0.49
2:B:169:ALA:HB2	2:B:184:ARG:HG2	1.95	0.49
1:A:40:PHE:HB3	1:A:118:CYS:SG	2.53	0.49
2:B:117:GLU:O	2:B:123:ARG:NH2	2.46	0.49
2:B:117:GLU:O	2:B:123:ARG:NE	2.47	0.48
2:B:127:VAL:O	2:B:131:TRP:HB2	2.13	0.48
2:B:179:THR:O	2:B:199:VAL:HA	2.14	0.48
1:A:38:THR:HB	1:A:63:ILE:HG21	1.95	0.48
2:B:117:GLU:HB3	2:B:123:ARG:HE	1.79	0.47
2:B:135:THR:O	2:B:146:GLY:N	2.44	0.47
1:A:38:THR:OG1	1:A:60:GLN:HG3	2.15	0.46
2:B:123:ARG:HH22	2:B:158:ARG:HD3	1.82	0.45
2:B:85:ALA:HA	2:B:88:LYS:HE3	1.98	0.44
2:B:136:ARG:O	2:B:210:ASN:HA	2.18	0.44
1:A:84:ILE:O	1:A:88:GLU:HG3	2.18	0.43
1:A:117:VAL:O	1:A:121:ILE:HG12	2.19	0.43
2:B:132:ALA:HA	2:B:207:ARG:HE	1.83	0.43
2:B:137:TYR:HD1	2:B:211:VAL:HB	1.82	0.42
1:A:177:LYS:HE2	1:A:181:GLN:HE21	1.85	0.42
2:B:181:PHE:CZ	2:B:198:ASP:HB2	2.55	0.41
2:B:201:SER:HB2	2:B:210:ASN:HB3	2.02	0.41
2:B:112:LEU:HD22	2:B:166:PHE:HB3	2.02	0.41
2:B:72:GLN:HB2	2:B:76:LYS:HE2	2.02	0.41
1:A:36:GLU:OE1	1:A:124:ARG:NH2	2.54	0.41
1:A:177:LYS:HE2	1:A:181:GLN:NE2	2.36	0.41
2:B:96:LYS:O	2:B:100:GLN:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:104:HIS:CE1	2:B:174:GLY:HA3	2.56	0.41
2:B:123:ARG:NH1	2:B:154:ILE:HG22	2.33	0.41
2:B:181:PHE:O	2:B:197:THR:HG23	2.21	0.40
1:A:91:ILE:O	1:A:95:GLU:HB2	2.22	0.40
1:A:171:GLU:O	1:A:175:LEU:HD23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/172 (98%)	166 (98%)	3 (2%)	0	100	100
2	B	144/157 (92%)	140 (97%)	4 (3%)	0	100	100
All	All	313/329 (95%)	306 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/144 (99%)	142 (99%)	1 (1%)	84	90
2	B	116/125 (93%)	113 (97%)	3 (3%)	46	67
All	All	259/269 (96%)	255 (98%)	4 (2%)	65	80

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	LEU
2	B	84	ASN
2	B	94	LEU
2	B	103	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.