



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 01:29 PM EDT

PDB ID : 6X4D
Title : Crystal Structure of HIV-1 Reverse Transcriptase in Complex with 5-(cyclopropylmethyl)-7-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-8-methyl-2-naphthonitrile (JLJ678), a Non-nucleoside Inhibitor
Authors : Chan, A.H.; Duong, V.N.; Ippolito, J.A.; Jorgensen, W.L.; Anderson, K.S.
Deposited on : 2020-05-22
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	
2	B	428	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	547	4415	2861	728	818	8	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 2 is a protein called p51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	401	3319	2160	550	603	6	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is 5-(cyclopropylmethyl)-7-{2-[2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy]phenoxy}-8-methylnaphthalene-2-carbonitrile (three-letter code: UO4) (formula: C₂₈H₂₅N₃O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	35	28	3	4	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
			O	S				
4	A	1	Total	5	4	1	0	0
4	A	1	Total	5	4	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	38	Total 38	O 38	0	0
5	B	33	Total 33	O 33	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.74Å 74.27Å 108.73Å 90.00° 100.02° 90.00°	Depositor
Resolution (Å)	43.37 – 2.65 43.37 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.37-2.65) 94.4 (43.37-2.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.224 , 0.258 0.224 , 0.259	Depositor DCC
R_{free} test set	1865 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7850	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UO4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/4532	0.42	0/6172
2	B	0.24	0/3413	0.41	0/4635
All	All	0.24	0/7945	0.42	0/10807

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4415	0	4419	29	0
2	B	3319	0	3341	32	0
3	A	35	0	0	0	0
4	A	10	0	0	1	0
5	A	38	0	0	0	0
5	B	33	0	0	1	0
All	All	7850	0	7760	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:TYR:O	1:A:143:ARG:NH2	2.29	0.66
1:A:28:GLU:HG3	1:A:135:ILE:HD12	1.79	0.65
2:B:308:GLU:HA	2:B:311:LYS:HE2	1.80	0.64
1:A:544:GLY:HA2	2:B:286:THR:HG22	1.78	0.63
2:B:107:THR:OG1	2:B:198:HIS:NE2	2.34	0.60
2:B:107:THR:HG1	2:B:198:HIS:HE2	1.50	0.60
1:A:543:GLY:HA3	2:B:284:ARG:HA	1.86	0.58
2:B:168:LEU:HD22	2:B:180:ILE:HD13	1.86	0.56
1:A:443:ASP:HB3	1:A:550:LYS:HD3	1.86	0.56
1:A:244:ILE:HG12	1:A:310:LEU:HD13	1.87	0.56
1:A:317:VAL:HG23	1:A:349:LEU:HD23	1.87	0.55
2:B:315:HIS:O	2:B:347:LYS:NZ	2.40	0.54
1:A:180:ILE:HG12	1:A:189:VAL:HG13	1.90	0.53
1:A:88:TRP:CD1	2:B:143:ARG:HD2	2.44	0.52
2:B:203:GLU:HA	2:B:206:ARG:HG2	1.92	0.52
1:A:491:LEU:HB3	1:A:529:GLU:HG2	1.92	0.52
2:B:72:ARG:NH2	2:B:151:GLN:OE1	2.43	0.51
2:B:73:LYS:NZ	2:B:146:TYR:OH	2.42	0.51
2:B:101:LYS:HD3	2:B:382:ILE:HG23	1.92	0.51
2:B:275:LYS:HD3	2:B:276:VAL:H	1.75	0.51
2:B:21:VAL:HB	2:B:59:PRO:HD3	1.92	0.51
2:B:425:LEU:HD23	2:B:428:GLN:HB3	1.93	0.51
2:B:275:LYS:HD3	2:B:276:VAL:N	2.27	0.50
1:A:340:GLN:HG3	1:A:351:THR:HG22	1.95	0.49
1:A:183:TYR:HB3	1:A:188:TYR:HE2	1.78	0.48
1:A:90:VAL:HG21	2:B:141:GLY:H	1.79	0.48
1:A:369:THR:O	1:A:373:GLN:HG2	2.13	0.48
2:B:323:LYS:NZ	2:B:344:GLU:OE2	2.38	0.48
2:B:328:GLU:O	2:B:339:TYR:HA	2.14	0.47
1:A:108:VAL:HG22	1:A:188:TYR:HD1	1.80	0.46
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.51	0.46
2:B:172:LYS:HG2	2:B:180:ILE:HD12	1.99	0.45
2:B:136:ASN:HB3	2:B:138:GLU:HG3	1.99	0.45
1:A:363:ASN:ND2	1:A:366:LYS:HE3	2.33	0.44
1:A:512:LYS:NZ	4:A:602:SO4:O3	2.51	0.44
1:A:110:ASP:O	1:A:217:PRO:HD3	2.17	0.43
1:A:503:LEU:HD22	1:A:535:TRP:HB2	2.01	0.43
1:A:363:ASN:HA	1:A:511:ASP:OD1	2.19	0.43
2:B:194:GLU:O	2:B:198:HIS:N	2.37	0.43
1:A:475:GLN:HB3	1:A:501:TYR:CE2	2.55	0.42
2:B:339:TYR:OH	2:B:378:GLU:OE1	2.30	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:38:CYS:SG	2:B:132:ILE:HD11	2.59	0.42
1:A:363:ASN:HD21	1:A:366:LYS:HE3	1.85	0.42
1:A:38:CYS:SG	1:A:132:ILE:HD11	2.59	0.42
1:A:90:VAL:HG11	2:B:140:PRO:HB3	2.00	0.42
2:B:82:LYS:NZ	5:B:505:HOH:O	2.53	0.42
2:B:104:LYS:HB2	2:B:192:ASP:HA	2.02	0.41
1:A:342:TYR:HB3	1:A:348:ASN:HA	2.03	0.41
2:B:281:LYS:HG3	2:B:284:ARG:NH1	2.35	0.41
2:B:393:ILE:HD13	2:B:398:TRP:HB2	2.01	0.41
2:B:164:MET:HG3	2:B:168:LEU:HD12	2.03	0.41
2:B:167:ILE:HG12	2:B:212:TRP:CE3	2.56	0.41
2:B:275:LYS:HD2	2:B:277:ARG:HG2	2.03	0.41
1:A:454:LYS:HZ2	1:A:552:VAL:HB	1.85	0.41
1:A:177:ASP:OD1	1:A:177:ASP:N	2.45	0.41
1:A:134:SER:HG	1:A:139:THR:HG1	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	543/557 (98%)	528 (97%)	15 (3%)	0	100	100
2	B	395/428 (92%)	380 (96%)	15 (4%)	0	100	100
All	All	938/985 (95%)	908 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	479/495 (97%)	470 (98%)	9 (2%)	57	74
2	B	364/390 (93%)	357 (98%)	7 (2%)	57	74
All	All	843/885 (95%)	827 (98%)	16 (2%)	57	74

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	MET
1	A	74	LEU
1	A	89	GLU
1	A	106	VAL
1	A	212	TRP
1	A	286	THR
1	A	295	LEU
1	A	399	GLU
1	A	546	GLU
2	B	8	VAL
2	B	69	THR
2	B	74	LEU
2	B	123	ASP
2	B	232	TYR
2	B	275	LYS
2	B	417	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	306	ASN
1	A	336	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	602	-	4,4,4	0.13	0	6,6,6	0.06	0
3	UO4	A	601	-	39,39,39	1.12	2 (5%)	55,55,55	1.53	12 (21%)
4	SO4	A	603	-	4,4,4	0.14	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UO4	A	601	-	-	3/16/18/18	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	UO4	C0K-N0H	2.67	1.41	1.37
3	A	601	UO4	C0Z-C0Y	2.55	1.42	1.37

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	UO4	C0Z-C0C-C0V	-3.64	118.20	122.45
3	A	601	UO4	C0C-O0A-C04	3.61	126.64	118.04
3	A	601	UO4	C10-C0V-C0C	-3.45	115.52	120.68
3	A	601	UO4	C13-C0Y-C0Z	-3.35	113.83	119.76
3	A	601	UO4	C15-C11-C0W	3.34	124.40	121.05
3	A	601	UO4	C13-C0Y-C0X	3.33	126.01	120.76
3	A	601	UO4	C11-C15-C18	2.73	122.01	119.65
3	A	601	UO4	C0C-C0V-C0W	2.62	122.17	118.84
3	A	601	UO4	C16-C12-C0X	2.30	124.33	121.13
3	A	601	UO4	C12-C0X-C0W	-2.14	115.79	118.45
3	A	601	UO4	C0Y-C13-C1C	2.12	119.26	114.34
3	A	601	UO4	C0O-C0P-N0H	-2.06	120.51	122.44

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	UO4	C0Y-C13-C1C-C1G
3	A	601	UO4	C0Z-C0Y-C13-C1C
3	A	601	UO4	O0B-C0D-C0E-N0H

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	602	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	547/557 (98%)	0.48	53 (9%) 7 6	25, 62, 122, 150	0
2	B	401/428 (93%)	0.34	26 (6%) 18 16	29, 54, 109, 128	0
All	All	948/985 (96%)	0.42	79 (8%) 11 9	25, 58, 116, 150	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	VAL	8.4
1	A	292	VAL	7.7
1	A	303	LEU	7.7
2	B	358	ARG	6.7
1	A	286	THR	5.8
2	B	359	GLY	5.8
1	A	289	LEU	5.3
1	A	300	GLU	5.2
1	A	283	LEU	5.1
1	A	218	ASP	5.0
1	A	297	GLU	4.9
1	A	291	GLU	4.9
1	A	255	ASN	4.8
1	A	295	LEU	4.7
1	A	287	LYS	4.7
2	B	241	VAL	4.6
1	A	311	LYS	4.5
1	A	252	TRP	4.5
1	A	548	VAL	4.5
2	B	356	ARG	4.4
1	A	302	GLU	4.2
1	A	246	LEU	4.0
1	A	299	ALA	3.9
2	B	85	GLN	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	237	ASP	3.9
2	B	172	LYS	3.8
1	A	288	ALA	3.8
2	B	239	TRP	3.8
1	A	221	HIS	3.6
1	A	222	GLN	3.6
1	A	551	LEU	3.6
1	A	293	ILE	3.6
2	B	238	LYS	3.6
2	B	207	GLN	3.5
2	B	232	TYR	3.4
1	A	290	THR	3.3
1	A	261	VAL	3.3
1	A	306	ASN	3.3
1	A	257	ILE	3.3
1	A	294	PRO	3.2
2	B	88	TRP	3.2
1	A	259	LYS	3.1
2	B	209	LEU	3.1
1	A	552	VAL	3.1
1	A	251	SER	2.9
1	A	253	THR	2.8
1	A	38	CYS	2.8
1	A	309	ILE	2.6
1	A	248	GLU	2.6
1	A	247	PRO	2.6
1	A	115	TYR	2.6
1	A	298	GLU	2.4
2	B	14	PRO	2.4
2	B	197	GLN	2.4
1	A	214	LEU	2.4
2	B	240	THR	2.4
2	B	206	ARG	2.4
1	A	90	VAL	2.3
1	A	280	SER	2.3
1	A	279	LEU	2.3
1	A	304	ALA	2.3
1	A	130	PHE	2.2
1	A	72	ARG	2.2
2	B	360	ALA	2.2
1	A	256	ASP	2.2
1	A	73	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	285	GLY	2.1
1	A	346	PHE	2.1
2	B	200	THR	2.1
2	B	87	PHE	2.1
1	A	24	TRP	2.1
1	A	301	LEU	2.1
2	B	208	HIS	2.1
2	B	69	THR	2.1
2	B	168	LEU	2.0
2	B	178	ILE	2.0
2	B	191	SER	2.0
1	A	116	PHE	2.0
2	B	199	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	602	5/5	0.88	0.18	73,74,92,92	0
3	UO4	A	601	35/35	0.95	0.19	41,50,61,66	0
4	SO4	A	603	5/5	0.96	0.28	88,92,105,106	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.