

wwPDB X-ray Structure Validation Summary Report (i)

Nov 22, 2023 – 10:22 PM JST

PDB ID : 7X52

Title: Crystal structure of Bacteroides thetaiotaomicron glutamate decarboxylase

BTGAD-PLP complex

Authors: Liu, S.; Du, G.; Wang, Y.; Wen, B.; Xin, F.

Deposited on : 2022-03-03

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	502	83%	8%	9%
1	В	502	86%	5%	9%
1	С	502	86%	5%	9%
1	D	502	84%	7%	9%
1	Е	502	85%	6%	9%
1	F	502	85%	6%	9%



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	В	501	-	-	X	-
3	ACT	Е	501	-	-	X	-
3	ACT	F	501	-	-	X	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 23934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glutamate decarboxylase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	458	Total	С	N	О	S	4	0	0
1	Λ	450	3670	2373	600	677	20	4	0	
1	В	459	Total	С	N	О	S	4	0	0
1	Ъ	409	3669	2372	601	676	20	4	0	0
1	С	459	Total	С	N	О	S	4	0	0
1		409	3686	2383	602	680	21	4		
1	D	458	Total	С	N	О	S	4	0	0
1	D	450	3660	2366	599	675	20	4	0	
1	Е	458	Total	С	N	О	S	4	0	0
1	l L	450	3666	2372	601	673	20	4	0	
1	F	458	Total	С	N	О	S	4	0	0
1	Г	400	3670	2374	601	675	20	4	U	0

There are 126 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP Q8A4M9
A	-19	GLY	-	expression tag	UNP Q8A4M9
A	-18	SER	-	expression tag	UNP Q8A4M9
A	-17	SER	-	expression tag	UNP Q8A4M9
A	-16	HIS	-	expression tag	UNP Q8A4M9
A	-15	HIS	-	expression tag	UNP Q8A4M9
A	-14	HIS	-	expression tag	UNP Q8A4M9
A	-13	HIS	-	expression tag	UNP Q8A4M9
A	-12	HIS	-	expression tag	UNP Q8A4M9
A	-11	HIS	-	expression tag	UNP Q8A4M9
A	-10	SER	-	expression tag	UNP Q8A4M9
A	-9	SER	-	expression tag	UNP Q8A4M9
A	-8	GLY	-	expression tag	UNP Q8A4M9
A	-7	LEU	-	expression tag	UNP Q8A4M9
A	-6	VAL	-	expression tag	UNP Q8A4M9
A	-5	PRO	-	expression tag	UNP Q8A4M9
A	-4	ARG	-	expression tag	UNP Q8A4M9



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q8A4M9
A	-2	SER	-	expression tag	UNP Q8A4M9
A	-1	HIS	-	expression tag	UNP Q8A4M9
A	0	MET	-	expression tag	UNP Q8A4M9
В	-20	MET	-	initiating methionine	UNP Q8A4M9
В	-19	GLY	-	expression tag	UNP Q8A4M9
В	-18	SER	-	expression tag	UNP Q8A4M9
В	-17	SER	-	expression tag	UNP Q8A4M9
В	-16	HIS	-	expression tag	UNP Q8A4M9
В	-15	HIS	-	expression tag	UNP Q8A4M9
В	-14	HIS	-	expression tag	UNP Q8A4M9
В	-13	HIS	-	expression tag	UNP Q8A4M9
В	-12	HIS	-	expression tag	UNP Q8A4M9
В	-11	HIS	-	expression tag	UNP Q8A4M9
В	-10	SER	-	expression tag	UNP Q8A4M9
В	-9	SER	-	expression tag	UNP Q8A4M9
В	-8	GLY	-	expression tag	UNP Q8A4M9
В	-7	LEU	ı	expression tag	UNP Q8A4M9
В	-6	VAL	-	expression tag	UNP Q8A4M9
В	-5	PRO	-	expression tag	UNP Q8A4M9
В	-4	ARG	-	expression tag	UNP Q8A4M9
В	-3	GLY	-	expression tag	UNP Q8A4M9
В	-2	SER	-	expression tag	UNP Q8A4M9
В	-1	HIS	-	expression tag	UNP Q8A4M9
В	0	MET	-	expression tag	UNP Q8A4M9
С	-20	MET	-	initiating methionine	UNP Q8A4M9
С	-19	GLY	-	expression tag	UNP Q8A4M9
С	-18	SER	-	expression tag	UNP Q8A4M9
С	-17	SER	-	expression tag	UNP Q8A4M9
С	-16	HIS	-	expression tag	UNP Q8A4M9
С	-15	HIS	-	expression tag	UNP Q8A4M9
С	-14	HIS	=	expression tag	UNP Q8A4M9
С	-13	HIS	-	expression tag	UNP Q8A4M9
С	-12	HIS	-	expression tag	UNP Q8A4M9
С	-11	HIS	-	expression tag	UNP Q8A4M9
С	-10	SER	-	expression tag	UNP Q8A4M9
С	-9	SER	-	expression tag	UNP Q8A4M9
С	-8	GLY	-	expression tag	UNP Q8A4M9
С	-7	LEU	=	expression tag	UNP Q8A4M9
С	-6	VAL	-	expression tag	UNP Q8A4M9
С	-5	PRO	=	expression tag	UNP Q8A4M9
С	-4	ARG		expression tag	UNP Q8A4M9



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	-3	GLY	-	expression tag	UNP Q8A4M9
С	-2	SER	-	expression tag	UNP Q8A4M9
С	-1	HIS	-	expression tag	UNP Q8A4M9
С	0	MET	-	expression tag	UNP Q8A4M9
D	-20	MET	-	initiating methionine	UNP Q8A4M9
D	-19	GLY	ı	expression tag	UNP Q8A4M9
D	-18	SER	-	expression tag	UNP Q8A4M9
D	-17	SER	-	expression tag	UNP Q8A4M9
D	-16	HIS	-	expression tag	UNP Q8A4M9
D	-15	HIS	-	expression tag	UNP Q8A4M9
D	-14	HIS	-	expression tag	UNP Q8A4M9
D	-13	HIS	-	expression tag	UNP Q8A4M9
D	-12	HIS	-	expression tag	UNP Q8A4M9
D	-11	HIS	-	expression tag	UNP Q8A4M9
D	-10	SER	-	expression tag	UNP Q8A4M9
D	-9	SER	-	expression tag	UNP Q8A4M9
D	-8	GLY	-	expression tag	UNP Q8A4M9
D	-7	LEU	-	expression tag	UNP Q8A4M9
D	-6	VAL	-	expression tag	UNP Q8A4M9
D	-5	PRO	-	expression tag	UNP Q8A4M9
D	-4	ARG	-	expression tag	UNP Q8A4M9
D	-3	GLY	-	expression tag	UNP Q8A4M9
D	-2	SER	-	expression tag	UNP Q8A4M9
D	-1	HIS	-	expression tag	UNP Q8A4M9
D	0	MET	-	expression tag	UNP Q8A4M9
Е	-20	MET	-	initiating methionine	UNP Q8A4M9
Е	-19	GLY	-	expression tag	UNP Q8A4M9
Е	-18	SER	ı	expression tag	UNP Q8A4M9
Е	-17	SER	ı	expression tag	UNP Q8A4M9
Е	-16	HIS	-	expression tag	UNP Q8A4M9
Е	-15	HIS	ı	expression tag	UNP Q8A4M9
Е	-14	HIS	ı	expression tag	UNP Q8A4M9
Е	-13	HIS	-	expression tag	UNP Q8A4M9
Е	-12	HIS	ı	expression tag	UNP Q8A4M9
Е	-11	HIS	-	expression tag	UNP Q8A4M9
Е	-10	SER	-	expression tag	UNP Q8A4M9
Е	-9	SER	-	expression tag	UNP Q8A4M9
Е	-8	GLY	-	expression tag	UNP Q8A4M9
Е	-7	LEU		expression tag	UNP Q8A4M9
Е	-6	VAL	-	expression tag	UNP Q8A4M9
Е	-5	PRO	-	expression tag	UNP Q8A4M9
Е	-4	ARG	-	expression tag	UNP Q8A4M9




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-3	GLY	-	expression tag	UNP Q8A4M9
Е	-2	SER	-	expression tag	UNP Q8A4M9
Е	-1	HIS	-	expression tag	UNP Q8A4M9
Е	0	MET	-	expression tag	UNP Q8A4M9
F	-20	MET	-	initiating methionine	UNP Q8A4M9
F	-19	GLY	-	expression tag	UNP Q8A4M9
F	-18	SER	-	expression tag	UNP Q8A4M9
F	-17	SER	-	expression tag	UNP Q8A4M9
F	-16	HIS	-	expression tag	UNP Q8A4M9
F	-15	HIS	-	expression tag	UNP Q8A4M9
F	-14	HIS	-	expression tag	UNP Q8A4M9
F	-13	HIS	-	expression tag	UNP Q8A4M9
F	-12	HIS	-	expression tag	UNP Q8A4M9
F	-11	HIS	-	expression tag	UNP Q8A4M9
F	-10	SER	-	expression tag	UNP Q8A4M9
F	-9	SER	-	expression tag	UNP Q8A4M9
F	-8	GLY	-	expression tag	UNP Q8A4M9
F	-7	LEU	ı	expression tag	UNP Q8A4M9
F	-6	VAL	=	expression tag	UNP Q8A4M9
F	-5	PRO	-	expression tag	UNP Q8A4M9
F	-4	ARG	-	expression tag	UNP Q8A4M9
F	-3	GLY	-	expression tag	UNP Q8A4M9
F	-2	SER	-	expression tag	UNP Q8A4M9
F	-1	HIS	-	expression tag	UNP Q8A4M9
F	0	MET	=	expression tag	UNP Q8A4M9

• Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
2	Λ	1	15	8	1	5	1		0
2	В	1	Total	С	N	О	Р	0	0
	D	1	15	8	1	5	1	0	U
2	С	1	Total	С	N	Ο	Р	0	0
		1	15	8	1	5	1	0	U
2	D	1	Total	С	N	Ο	Р	0	0
	D	1	15	8	1	5	1	0	U
2	E	1	Total	С	N	Ο	Р	0	0
	Ľ	1	15	8	1	5	1	0	U
2	F	1	Total	С	N	О	Р	0	0
	I.	1	15	8	1	5	1		U

 \bullet Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2)$ (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0

• Molecule 4 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 3 4	0	0
4	В	1	Total C O 7 3 4	0	0
4	D	1	Total C O 7 3 4	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	354	Total O 354 354	0	0
5	В	295	Total O 295 295	0	0
5	С	354	Total O 354 354	0	0
5	D	282	Total O 282 282	0	0
5	E	242	Total O 242 242	0	0
5	F	247	Total O 247 247	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate decarboxylase









4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	109.77Å 131.27Å 109.85Å	Depositor	
a, b, c, α , β , γ	90.00° 100.28° 90.00°	Depositor	
Resolution (Å)	36.00 - 1.90	Depositor	
resolution (A)	36.00 - 1.90	EDS	
% Data completeness	98.7 (36.00-1.90)	Depositor	
(in resolution range)	98.7 (36.00-1.90)	EDS	
R_{merge}	0.07	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.11 (at 1.89Å)	Xtriage	
Refinement program	PHENIX 1.19.1_4122	Depositor	
R, R_{free}	0.165 , 0.194	Depositor	
it, it free	0.165 , 0.194	DCC	
R_{free} test set	12067 reflections $(5.07%)$	wwPDB-VP	
Wilson B-factor (Å ²)	27.3	Xtriage	
Anisotropy	0.452	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.32 \; , 32.6$	EDS	
L-test for twinning ²	$< L > = 0.46, < L^2> = 0.29$	Xtriage	
Estimated twinning fraction	0.260 for l,-k,h	Xtriage	
F_o, F_c correlation	0.97	EDS	
Total number of atoms	23934	wwPDB-VP	
Average B, all atoms (Å ²)	34.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, ACT, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.41	0/3769	0.59	0/5127	
1	В	0.38	0/3768	0.57	0/5126	
1	С	0.41	0/3785	0.59	0/5146	
1	D	0.37	0/3759	0.58	0/5115	
1	Е	0.34	0/3765	0.54	0/5121	
1	F	0.35	0/3769	0.55	0/5126	
All	All	0.38	0/22615	0.57	0/30761	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3670	0	3606	26	0
1	В	3669	0	3600	20	0
1	С	3686	0	3633	21	0
1	D	3660	0	3584	32	0
1	Е	3666	0	3609	26	0
1	F	3670	0	3612	19	0
2	A	15	0	7	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	15	0	7	2	0
2	С	15	0	7	2	0
2	D	15	0	7	1	0
2	Ε	15	0	7	1	0
2	F	15	0	7	1	0
3	A	4	0	3	0	0
3	В	4	0	3	3	0
3	С	8	0	6	0	0
3	Ε	8	0	6	4	0
3	F	4	0	3	3	0
4	A	7	0	2	0	0
4	В	7	0	2	0	0
4	D	7	0	2	0	0
5	A	354	0	0	3	0
5	В	295	0	0	1	0
5	С	354	0	0	1	0
5	D	282	0	0	3	0
5	Ε	242	0	0	2	0
5	F	247	0	0	2	0
All	All	23934	0	21713	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$	
3:B:501:ACT:H1	1:E:61:PHE:H	1.24	1.00	
1:D:310:GLN:HG2	1:D:311:VAL:H	1.45	0.81	
1:B:141:LYS:HZ3	1:B:142:ARG:HH12	1.29	0.79	
1:B:141:LYS:NZ	1:B:142:ARG:HH12	1.80	0.79	
1:B:60:THR:OG1	3:E:501:ACT:H1	1.82	0.78	

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	456/502 (91%)	444 (97%)	12 (3%)	0	100	100
1	В	457/502 (91%)	445 (97%)	12 (3%)	0	100	100
1	\mathbf{C}	457/502 (91%)	445 (97%)	12 (3%)	0	100	100
1	D	456/502 (91%)	442 (97%)	14 (3%)	0	100	100
1	E	456/502 (91%)	443 (97%)	13 (3%)	0	100	100
1	F	456/502 (91%)	441 (97%)	15 (3%)	0	100	100
All	All	2738/3012 (91%)	2660 (97%)	78 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	392/432 (91%)	390 (100%)	2 (0%)	88	89	
1	В	390/432 (90%)	388 (100%)	2 (0%)	88	89	
1	С	395/432 (91%)	393 (100%)	2 (0%)	88	89	
1	D	389/432 (90%)	386 (99%)	3 (1%)	81	82	
1	E	391/432 (90%)	389 (100%)	2 (0%)	88	89	
1	F	392/432 (91%)	389 (99%)	3 (1%)	81	82	
All	All	2349/2592 (91%)	2335 (99%)	14 (1%)	86	87	

5 of 14 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	D	372	LEU
1	D	413	GLU
1	F	372	LEU
1	F	151	LYS
1	F	331	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	302	ASN
1	D	310	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type	Type	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Res	Link	Bo	Bond lengths		Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2													
3	ACT	E	501	-	3,3,3	1.05	0	3,3,3	1.70	1 (33%)													
2	PLP	D	502	1	15,15,16	0.91	1 (6%)	20,22,23	1.25	2 (10%)													



Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	F	501	-	3,3,3	0.57	0	3,3,3	1.52	0
2	PLP	E	503	1	15,15,16	1.13	1 (6%)	20,22,23	1.27	3 (15%)
4	MLI	В	502	-	6,6,6	1.47	1 (16%)	7,7,7	1.47	1 (14%)
2	PLP	С	503	1	15,15,16	1.11	1 (6%)	20,22,23	1.44	3 (15%)
2	PLP	В	503	1	15,15,16	1.06	1 (6%)	20,22,23	1.50	3 (15%)
3	ACT	В	501	-	3,3,3	0.94	0	3,3,3	1.59	1 (33%)
3	ACT	A	502	-	3,3,3	1.24	0	3,3,3	1.31	0
4	MLI	A	503	-	6,6,6	1.30	0	7,7,7	1.36	0
3	ACT	С	502	-	3,3,3	1.40	0	3,3,3	1.35	0
2	PLP	A	501	1	15,15,16	1.00	1 (6%)	20,22,23	1.32	3 (15%)
4	MLI	D	501	-	6,6,6	1.41	0	7,7,7	1.70	2 (28%)
3	ACT	С	501	-	3,3,3	0.94	0	3,3,3	0.86	0
3	ACT	Е	502	-	3,3,3	1.35	0	3,3,3	1.42	0
2	PLP	F	502	1	15,15,16	1.00	1 (6%)	20,22,23	1.15	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	D	502	1	-	0/6/6/8	0/1/1/1
2	PLP	Е	503	1	-	1/6/6/8	0/1/1/1
4	MLI	В	502	_	-	2/4/4/4	-
2	PLP	С	503	1	-	0/6/6/8	0/1/1/1
2	PLP	В	503	1	-	3/6/6/8	0/1/1/1
4	MLI	A	503	-	-	3/4/4/4	_
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1
4	MLI	D	501	_	-	2/4/4/4	-
2	PLP	F	502	1	-	5/6/6/8	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
2	С	503	PLP	C2-N1	2.83	1.39	1.33
2	Ε	503	PLP	C2-N1	2.68	1.38	1.33
2	F	502	PLP	C2-N1	2.54	1.38	1.33
2	В	503	PLP	C2-N1	2.43	1.38	1.33
2	A	501	PLP	C2-N1	2.28	1.38	1.33



The worst	5	of 21	bond	angle	outliers	are	listed	below:
110 110100	$\overline{}$	OI -I	OILG	ari Sic	Catheren	COL C	IID CCC	CIC III .

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	503	PLP	C6-C5-C4	3.82	121.16	118.16
2	С	503	PLP	C6-C5-C4	3.52	120.93	118.16
2	В	503	PLP	C4A-C4-C5	-3.14	117.70	120.94
2	D	502	PLP	C6-C5-C4	3.14	120.63	118.16
2	F	502	PLP	C6-C5-C4	3.12	120.61	118.16

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	503	PLP	C5A-O4P-P-O1P
2	В	503	PLP	C5A-O4P-P-O3P
2	F	502	PLP	C5A-O4P-P-O2P
2	F	502	PLP	C5A-O4P-P-O3P
4	A	503	MLI	C2-C1-C3-O9

There are no ring outliers.


9 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Е	501	ACT	4	0
2	D	502	PLP	1	0
3	F	501	ACT	3	0
2	Е	503	PLP	1	0
2	С	503	PLP	2	0
2	В	503	PLP	2	0
3	В	501	ACT	3	0
2	A	501	PLP	1	0
2	F	502	PLP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient




equivalents in the CSD to analyse the geometry.



























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	458/502 (91%)	-0.58	0 100 100	17, 26, 45, 69	1 (0%)
1	В	459/502 (91%)	-0.52	8 (1%) 70 72	22, 31, 53, 79	1 (0%)
1	С	459/502 (91%)	-0.58	0 100 100	17, 26, 44, 70	1 (0%)
1	D	458/502 (91%)	-0.51	10 (2%) 62 64	22, 32, 54, 76	1 (0%)
1	E	458/502 (91%)	-0.45	6 (1%) 77 79	21, 36, 58, 86	1 (0%)
1	F	458/502 (91%)	-0.45	10 (2%) 62 64	22, 36, 62, 98	1 (0%)
All	All	2750/3012 (91%)	-0.51	34 (1%) 79 81	17, 31, 55, 98	6 (0%)

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	308	ILE	4.9
1	В	301	VAL	4.3
1	D	302	ASN	4.3
1	D	309	THR	4.2
1	D	310	GLN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	ACT	Е	502	4/4	0.73	0.15	50,58,59,59	0
4	MLI	D	501	7/7	0.83	0.25	32,42,44,47	0
4	MLI	A	503	7/7	0.84	0.23	37,39,53,53	0
3	ACT	С	502	4/4	0.85	0.14	34,36,36,38	0
3	ACT	С	501	4/4	0.87	0.15	15,24,24,33	0
4	MLI	В	502	7/7	0.89	0.21	33,40,41,45	0
3	ACT	Е	501	4/4	0.89	0.11	18,26,29,38	0
3	ACT	F	501	4/4	0.91	0.11	23,27,28,38	0
3	ACT	В	501	4/4	0.93	0.09	26,30,31,39	0
2	PLP	F	502	15/16	0.96	0.10	28,38,45,47	0
2	PLP	D	502	15/16	0.96	0.11	28,32,35,35	0
2	PLP	Е	503	15/16	0.97	0.12	28,33,37,38	0
2	PLP	В	503	15/16	0.98	0.10	24,29,32,32	0
2	PLP	С	503	15/16	0.98	0.08	20,22,26,28	0
3	ACT	A	502	4/4	0.98	0.07	21,22,23,25	0
2	PLP	A	501	15/16	0.98	0.10	19,22,25,26	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ACT E 502:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ACT C 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








Electron density around ACT E 501: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ACT B 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around PLP F 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around PLP D 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



















6.5 Other polymers (i)

There are no such residues in this entry.

