

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2023 – 05:55 AM EDT

PDB ID	:	6X9D
Title	:	Structure of proline utilization A with trans-4-hydroxy-L-proline bound in the
		L-glutamate-gamma-semialdehyde dehydrogenase active site
Authors	:	Tanner, J.J.; Campbell, A.C.
Deposited on	:	2020-06-02
Resolution	:	1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	$2580 \ (1.56-1.52)$
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	1235	^{2%} 95% •	
1	В	1235	4% 93% 5% •	I



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 20619 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bifunctional protein PutA.

Mol	Chain	Residues		Α	toms			ZeroOcc	AltConf	Trace
1	А	1218	Total 9088	C 5717	N 1631	O 1705	$\begin{array}{c} \mathrm{S} \\ \mathrm{35} \end{array}$	0	9	0
1	В	1217	Total 9046	C 5699	N 1621	O 1692	S 34	0	9	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-1	SER	-	expression tag	UNP F7X6I3
А	0	MET	-	expression tag	UNP F7X6I3
В	-1	SER	-	expression tag	UNP F7X6I3
В	0	MET	-	expression tag	UNP F7X6I3

• Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
0	Δ	1	Total	С	Ν	Ο	Р	0	0
	A	1	53	27	9	15	2	0	0
0	р	1	Total	С	Ν	Ο	Р	0	0
	D	1	53	27	9	15	2	0	0

• Molecule 3 is 4-HYDROXYPROLINE (three-letter code: HYP) (formula: $C_5H_9NO_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Δ	1	Total C N O	0	0
0	3 A	T	9 5 1 3	0	0
2	Р	1	Total C N O	0	0
5	D	L	9 5 1 3	0	0

• Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
4	Λ	1	Total C O	0	0 0	
4	Л	T	10 6 4	0	0	
4	В	1	Total C O	0	0	
4	D	L	10 6 4	0	0	

• Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	А	1	Total 3	С 1	O 2	0	0

 \bullet Molecule 6 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD)



(formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	Λ	1	Total	С	Ν	Ο	Р	0	0
0	Л	T	44	21	7	14	2	0	0
6	D	1	Total	С	Ν	Ο	Р	0	0
0	D	L	44	21	7	14	2	0	0



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	А	1	Total 5	0 4	S 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	1	Total Mg 1 1	0	0
8	В	1	Total Mg 1 1	0	0

• Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	В	1	Total C 7 4	O 3	0	0

• Molecule 10 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	1099	Total O 1099 1099	0	0
10	В	1112	Total O 1112 1112	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Bifunctional protein PutA



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	101.32Å 102.14Å 126.46Å	Depositor
a, b, c, α , β , γ	90.00° 106.42° 90.00°	Depositor
Bosolution(Å)	48.59 - 1.54	Depositor
Resolution (A)	48.59 - 1.54	EDS
% Data completeness	99.6 (48.59-1.54)	Depositor
(in resolution range)	99.6~(48.59-1.54)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.54 (at 1.54 Å)	Xtriage
Refinement program	PHENIX 1.14	Depositor
P. P.	0.179 , 0.210	Depositor
n, n_{free}	0.179 , 0.211	DCC
R_{free} test set	18022 reflections (4.97%)	wwPDB-VP
Wilson B-factor $(Å^2)$	15.9	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 40.3	EDS
L-test for $twinning^2$	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20619	wwPDB-VP
Average B, all atoms $(Å^2)$	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NAD, HYP, PGE, MG, SO4, FAD, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.33	0/9275	0.54	1/12615~(0.0%)	
1	В	0.34	0/9232	0.54	0/12564	
All	All	0.33	0/18507	0.54	1/25179~(0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	174	ARG	NE-CZ-NH1	5.64	123.12	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	9088	0	9141	30	0
1	В	9046	0	9090	38	0
2	А	53	0	30	4	0
2	В	53	0	31	4	0
3	А	9	0	8	0	0
3	В	9	0	8	0	0
4	А	10	0	14	0	0
4	В	10	0	14	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	А	3	0	1	0	0
6	А	44	0	26	2	0
6	В	44	0	26	2	0
7	А	20	0	0	0	0
7	В	10	0	0	0	0
8	А	1	0	0	0	0
8	В	1	0	0	0	0
9	В	7	0	10	0	0
10	А	1099	0	0	6	0
10	В	1112	0	0	5	0
All	All	20619	0	18399	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:473:TYR:HB2	2:B:2001:FAD:HM72	1.65	0.78
1:B:844:CYS:SG	6:B:2005:NAD:C4N	2.79	0.69
1:A:473:TYR:HB2	2:A:2001:FAD:HM72	1.74	0.69
1:B:793:GLN:OE1	10:B:2101:HOH:O	2.12	0.68
1:A:844:CYS:SG	6:A:2005:NAD:C4N	2.83	0.65
1:A:1183:LEU:O	10:A:2101:HOH:O	2.15	0.64
1:B:674:GLU:HG2	1:B:711:ALA:HB1	1.80	0.64
1:B:539:GLU:OE1	10:B:2102:HOH:O	2.16	0.63
1:A:873:ILE:HG13	1:A:883:VAL:HB	1.80	0.62
1:B:339:VAL:HG21	1:B:350:LEU:HD21	1.84	0.60
1:A:958:ARG:NH1	10:A:2108:HOH:O	2.34	0.59
1:A:1147:LEU:HD22	1:B:1147:LEU:HD13	1.86	0.58
1:B:297:LYS:HD2	1:B:329:LEU:HA	1.85	0.58
1:A:458:GLU:OE1	10:A:2102:HOH:O	2.17	0.57
1:B:856:VAL:HG12	1:B:859:ARG:HH22	1.70	0.56
1:A:286:LEU:HD21	1:A:322:VAL:HG11	1.88	0.56
1:B:358:ARG:HG2	1:B:415:VAL:HG11	1.90	0.54
1:B:615:ARG:NH2	10:B:2106:HOH:O	2.37	0.52
1:B:286:LEU:HD21	1:B:322:VAL:HG11	1.92	0.51
1:B:650:MET:O	1:B:654:LEU:HG	2.10	0.51
1:B:1143:GLN:O	1:B:1147:LEU:HG	2.10	0.51
1:A:650:MET:O	1:A:654:LEU:HG	2.11	0.51
1:B:492:GLU:HB3	2:B:2001:FAD:H4'	1.94	0.50



6X9D

		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:339:VAL:HG21	1:A:350:LEU:HD21	1.94	0.50
1:A:961:ASP:OD2	1:B:1055:LYS:NZ	2.38	0.49
1:B:473:TYR:HE2	1:B:489:ARG:HH22	1.60	0.48
1:A:298:ASN:ND2	10:A:2105:HOH:O	2.31	0.48
1:A:878:ARG:HD2	10:A:2413:HOH:O	2.13	0.48
1:B:782:GLY:O	1:B:811:THR:HA	2.13	0.48
1:B:844:CYS:SG	6:B:2005:NAD:C3N	3.02	0.47
1:A:937:LEU:HD21	1:A:947:HIS:CD2	2.49	0.47
1:B:323:LEU:HD13	1:B:335:MET:HE3	1.96	0.47
1:A:1056:TRP:CD1	1:A:1142:PRO:HD3	2.51	0.46
1:A:803:ARG:NH2	1:A:1197:GLU:OE1	2.41	0.45
1:A:473:TYR:CB	2:A:2001:FAD:HM72	2.44	0.45
1:A:952:ARG:HE	1:A:952:ARG:HB3	1.61	0.45
1:B:371:GLY:N	2:B:2001:FAD:O2'	2.50	0.44
1:B:1069:ALA:HA	1:B:1117:ALA:HB1	2.00	0.44
1:A:562:LEU:HD11	1:A:654:LEU:HD12	2.00	0.43
1:B:124:LYS:HD2	1:B:133:HIS:CE1	2.52	0.43
1:A:272:ARG:HB3	1:A:277:GLN:HG3	1.98	0.43
1:B:128:GLY:O	1:B:130:TRP:N	2.48	0.43
1:B:473:TYR:CB	2:B:2001:FAD:HM72	2.41	0.43
1:A:918:THR:HB	1:A:923:PHE:CD1	2.54	0.43
1:B:1056:TRP:CD1	1:B:1142:PRO:HD3	2.53	0.43
1:A:74:ALA:O	1:A:78:LYS:HG2	2.19	0.43
1:A:958:ARG:O	1:A:961:ASP:HB2	2.19	0.42
1:A:492:GLU:HB3	2:A:2001:FAD:H4'	2.00	0.42
1:A:782:GLY:O	1:A:811:THR:HA	2.20	0.42
1:B:374:TRP:HZ3	1:B:1229:LEU:HB3	1.84	0.42
1:B:700[B]:ILE:HG12	1:B:725:ASN:HB3	2.00	0.42
1:B:717:ILE:HG12	1:B:727:VAL:HG11	2.02	0.42
1:B:323:LEU:HB3	1:B:335:MET:HE3	2.01	0.42
1:B:1178:LYS:HB3	1:B:1178:LYS:HE3	1.89	0.41
1:A:1167:GLU:HA	1:A:1193:ALA:O	2.20	0.41
1:A:844:CYS:SG	6:A:2005:NAD:C3N	3.09	0.41
1:B:892:LYS:NZ	10:B:2142:HOH:O	2.54	0.41
1:A:1143:GLN:O	1:A:1147:LEU:HG	2.21	0.41
1:A:371:GLY:N	2:A:2001:FAD:O2'	2.54	0.41
1:B:297:LYS:HG3	1:B:332:TRP:HB2	2.02	0.40
1:B:873:ILE:HD11	1:B:912:ILE:HD11	2.03	0.40
1:B:1058:ASP:OD2	1:B:1070[B]:ARG:NH2	2.26	0.40
1:A:440:HIS:HE1	10:A:3040:HOH:O	2.03	0.40
1:B:643:ALA:HB1	1:B:676:ILE:HG23	2.02	0.40



Atom-1	Atom-1 Atom-2		Clash overlap (Å)
1:B:662:GLY:HA2	10:B:2247:HOH:O	2.21	0.40
1:B:78:LYS:NZ	1:B:455:PRO:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	1223/1235~(99%)	1206 (99%)	17 (1%)	0	100	100
1	В	1223/1235~(99%)	1200 (98%)	23~(2%)	0	100	100
All	All	2446/2470 (99%)	2406 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	919/951~(97%)	911 (99%)	8 (1%)	78 60		
1	В	909/951~(96%)	901 (99%)	8 (1%)	78 60		
All	All	1828/1902~(96%)	1812 (99%)	16 (1%)	78 60		

All (16) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	39	TYR
1	А	342	TYR
1	А	730	LYS
1	А	764	ARG
1	А	796	ASP
1	А	861	LEU
1	А	872	HIS
1	А	934	LEU
1	В	39	TYR
1	В	342	TYR
1	В	604	SER
1	В	730	LYS
1	В	833	ILE
1	В	872	HIS
1	В	934	LEU
1	В	937	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	440	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Dog	Link	B	ond leng	gths	Bond angles		
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HYP	А	2002	-	$9,\!9,\!9$	1.13	1 (11%)	10,12,12	1.16	1 (10%)
5	FMT	А	2004	-	2,2,2	0.64	0	1,1,1	0.12	0
7	SO4	А	2006	-	$4,\!4,\!4$	0.10	0	6,6,6	0.18	0
7	SO4	В	2007	-	$4,\!4,\!4$	0.14	0	6,6,6	0.11	0
7	SO4	А	2008	-	$4,\!4,\!4$	0.16	0	6,6,6	0.17	0
2	FAD	В	2001	-	$53,\!58,\!58$	1.97	13 (24%)	68,89,89	1.30	9 (13%)
3	HYP	В	2002	-	$9,\!9,\!9$	1.15	1 (11%)	10,12,12	0.68	0
4	PGE	А	2003	-	$9,\!9,\!9$	0.51	0	8,8,8	0.20	0
4	PGE	В	2003	-	$9,\!9,\!9$	0.53	0	8,8,8	0.31	0
9	PEG	В	2004	-	$6,\!6,\!6$	0.46	0	$5,\!5,\!5$	0.19	0
7	SO4	В	2006	-	$4,\!4,\!4$	0.26	0	6,6,6	0.27	0
6	NAD	А	2005	8	42,48,48	3.54	16 (38%)	50,73,73	1.67	6 (12%)
6	NAD	В	2005	8	42,48,48	3.43	15 (35%)	50,73,73	1.64	5 (10%)
7	SO4	А	2009	-	4,4,4	0.17	0	6,6,6	0.16	0
7	SO4	A	2007	-	4,4,4	0.16	0	6,6,6	0.20	0
2	FAD	А	2001	-	$53,\!58,\!58$	1.88	13 (24%)	68,89,89	1.43	10 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HYP	А	2002	-	-	0/4/13/13	0/1/1/1
2	FAD	В	2001	-	-	8/30/50/50	0/6/6/6
3	HYP	В	2002	-	-	1/4/13/13	0/1/1/1
4	PGE	А	2003	-	-	2/7/7/7	-
4	PGE	В	2003	-	-	3/7/7/7	-
9	PEG	В	2004	-	-	3/4/4/4	-
6	NAD	А	2005	8	-	5/26/62/62	0/5/5/5
6	NAD	В	2005	8	-	1/26/62/62	0/5/5/5
2	FAD	А	2001	-	-	7/30/50/50	0/6/6/6

All (59) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\operatorname{\AA})$	
6	А	2005	NAD	O4D-C1D	-8.51	1.29	1.41	
6	В	2005	NAD	O4D-C1D	-8.38	1.29	1.41	
6	А	2005	NAD	C3B-C4B	-8.32	1.31	1.53	
6	В	2005	NAD	C3B-C4B	-8.19	1.32	1.53	
6	А	2005	NAD	C7N-N7N	8.10	1.48	1.33	
6	В	2005	NAD	C7N-N7N	7.94	1.48	1.33	
6	А	2005	NAD	C3D-C4D	-7.65	1.33	1.53	
6	А	2005	NAD	O4D-C4D	7.19	1.61	1.45	
6	А	2005	NAD	O4B-C4B	7.19	1.61	1.45	
2	В	2001	FAD	O4-C4	7.09	1.37	1.23	
6	В	2005	NAD	C3D-C4D	-7.08	1.34	1.53	
6	В	2005	NAD	O4B-C4B	6.88	1.60	1.45	
6	В	2005	NAD	O4D-C4D	6.82	1.60	1.45	
2	А	2001	FAD	O4-C4	6.75	1.36	1.23	
2	В	2001	FAD	O2-C2	5.57	1.34	1.24	
6	А	2005	NAD	O4B-C1B	-5.52	1.33	1.41	
2	А	2001	FAD	O2-C2	5.03	1.33	1.24	
6	В	2005	NAD	O4B-C1B	-4.95	1.34	1.41	
6	В	2005	NAD	C3N-C7N	4.55	1.57	1.50	
2	А	2001	FAD	C4X-N5	4.48	1.39	1.30	
6	А	2005	NAD	C3N-C7N	4.41	1.57	1.50	
2	В	2001	FAD	C4X-N5	4.32	1.39	1.30	
6	А	2005	NAD	O3D-C3D	4.31	1.53	1.43	
6	В	2005	NAD	C6A-N6A	4.28	1.49	1.34	
6	В	2005	NAD	O3D-C3D	4.12	1.52	1.43	
6	В	2005	NAD	C2N-N1N	3.96	1.39	1.35	
6	А	2005	NAD	C6A-N6A	3.96	1.48	1.34	
6	А	2005	NAD	O3B-C3B	3.31	1.50	1.43	
2	А	2001	FAD	O2'-C2'	-3.12	1.36	1.43	
2	В	2001	FAD	C2-N1	3.11	1.44	1.36	
6	А	2005	NAD	C2N-N1N	3.06	1.38	1.35	
2	В	2001	FAD	C2B-C1B	-3.02	1.49	1.53	
2	В	2001	FAD	C2A-N3A	2.88	1.36	1.32	
2	A	2001	FAD	C2-N1	2.84	1.43	1.36	
2	A	2001	FAD	C6A-N6A	2.83	1.44	1.34	
2	В	2001	FAD	O2'-C2'	-2.82	1.37	1.43	
2	B	2001	FAD	C6A-N6A	2.79	1.44	1.34	
6	B	2005	NAD	O3B-C3B	2.67	1.49	1.43	
6	B	2005	NAD	C2A-N1A	2.65	1.38	1.33	
2	A	2001	FAD	C2B-C1B	-2.60	1.49	1.53	
2	B	2001	FAD	C10-N1	2.55	1.38	1.33	
6	A	2005	NAD	C5A-C4A	-2.55	1.34	1.40	
6	А	2005	NAD	O7N-C7N	-2.53	1.19	1.24	



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	А	2005	NAD	C2A-N1A	2.50	1.38	1.33
2	В	2001	FAD	O4'-C4'	-2.42	1.38	1.43
2	В	2001	FAD	PA-O5B	-2.42	1.49	1.59
2	А	2001	FAD	PA-O5B	-2.41	1.49	1.59
3	В	2002	HYP	CA-C	2.39	1.58	1.52
2	А	2001	FAD	C2A-N3A	2.38	1.35	1.32
2	А	2001	FAD	O4B-C4B	-2.32	1.39	1.45
2	В	2001	FAD	O4B-C4B	-2.30	1.39	1.45
3	А	2002	HYP	CA-C	2.28	1.58	1.52
6	А	2005	NAD	O2B-C2B	-2.23	1.37	1.43
2	А	2001	FAD	O4'-C4'	-2.18	1.38	1.43
2	А	2001	FAD	PA-O2A	-2.09	1.45	1.55
2	А	2001	FAD	C10-N1	2.04	1.37	1.33
6	В	2005	NAD	O7N-C7N	-2.03	1.20	1.24
6	В	2005	NAD	C5A-C4A	-2.03	1.35	1.40
2	В	2001	FAD	O3'-C3'	-2.03	1.38	1.43

Continued from previous page...

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
6	А	2005	NAD	C1B-N9A-C4A	-7.78	112.98	126.64
6	В	2005	NAD	C1B-N9A-C4A	-6.61	115.02	126.64
6	В	2005	NAD	N3A-C2A-N1A	-5.54	120.01	128.68
6	А	2005	NAD	N3A-C2A-N1A	-5.37	120.28	128.68
2	А	2001	FAD	N3A-C2A-N1A	-5.35	120.31	128.68
2	В	2001	FAD	N3A-C2A-N1A	-5.30	120.39	128.68
2	А	2001	FAD	C4-C4X-N5	3.64	123.42	118.23
2	А	2001	FAD	C4-N3-C2	-3.38	119.40	125.64
6	В	2005	NAD	O4B-C1B-C2B	-2.99	102.55	106.93
6	В	2005	NAD	O4D-C1D-C2D	-2.96	102.61	106.93
6	В	2005	NAD	C3N-C2N-N1N	-2.91	117.59	120.43
2	А	2001	FAD	C4X-C4-N3	2.86	120.46	113.19
6	А	2005	NAD	C3N-C2N-N1N	-2.60	117.88	120.43
6	А	2005	NAD	O4B-C1B-C2B	-2.60	103.13	106.93
2	А	2001	FAD	C4X-C10-N10	2.47	120.09	116.48
2	В	2001	FAD	C4-N3-C2	-2.46	121.10	125.64
2	А	2001	FAD	C10-C4X-N5	-2.45	119.67	124.86
2	В	2001	FAD	C4-C4X-N5	2.38	121.61	118.23
2	A	2001	FAD	C4X-C10-N1	-2.35	119.28	124.73
2	В	2001	FAD	C4X-C4-N3	2.34	119.13	113.19
2	В	2001	FAD	O4-C4-C4X	-2.29	120.52	126.60
6	А	2005	NAD	O4D-C1D-C2D	-2.26	103.62	106.93



6

А

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	2002	HYP	O-C-CA	-2.19	115.05	122.26
2	А	2001	FAD	O4-C4-C4X	-2.17	120.84	126.60
2	В	2001	FAD	O5'-P-O1P	2.16	117.49	109.07
2	А	2001	FAD	C2A-N1A-C6A	2.15	122.43	118.75
2	В	2001	FAD	C4X-C10-N10	2.14	119.61	116.48
2	А	2001	FAD	O5'-P-O1P	2.14	117.43	109.07
2	В	2001	FAD	C9-C9A-N10	-2.12	118.98	121.84
2	В	2001	FAD	C4A-C5A-N7A	-2.05	107.26	109.40
6	А	2005	NAD	C2N-C3N-C4N	2.03	120.56	118.26

C5B-O5B-PA-O1A

Continued from previous page...

There are no chirality outliers.

Mol	Chain	\mathbf{Res}	Type	Atoms
2	А	2001	FAD	C2'-C3'-C4'-O4'
2	А	2001	FAD	C2'-C3'-C4'-C5'
2	А	2001	FAD	O3'-C3'-C4'-O4'
2	А	2001	FAD	O3'-C3'-C4'-C5'
2	В	2001	FAD	C2'-C3'-C4'-O4'
2	В	2001	FAD	C2'-C3'-C4'-C5'
2	В	2001	FAD	O3'-C3'-C4'-O4'
2	В	2001	FAD	O3'-C3'-C4'-C5'

2005

All (30) torsion outliers are listed below:

4	А	2003	PGE	O1-C1-C2-O2
4	В	2003	PGE	O1-C1-C2-O2
4	В	2003	PGE	O3-C5-C6-O4
9	В	2004	PEG	O1-C1-C2-O2
6	А	2005	NAD	C3D-C4D-C5D-O5D
9	В	2004	PEG	O2-C3-C4-O4
6	А	2005	NAD	C4D-C5D-O5D-PN
2	А	2001	FAD	P-O3P-PA-O5B
2	В	2001	FAD	P-O3P-PA-O5B
6	В	2005	NAD	C4D-C5D-O5D-PN
2	А	2001	FAD	C5B-O5B-PA-O3P
2	В	2001	FAD	C5B-O5B-PA-O3P
2	В	2001	FAD	C3B-C4B-C5B-O5B
6	А	2005	NAD	O4D-C4D-C5D-O5D
3	В	2002	HYP	O-C-CA-CB
9	В	2004	PEG	C4-C3-O2-C2
4	В	2003	PGE	C4-C3-O2-C2
6	А	2005	NAD	C5B-O5B-PA-O3
2	А	2001	FAD	C3B-C4B-C5B-O5B

NAD



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type	Atoms
4	А	2003	PGE	C4-C3-O2-C2
2	В	2001	FAD	C4'-C5'-O5'-P

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	2001	FAD	4	0
6	А	2005	NAD	2	0
6	В	2005	NAD	2	0
2	А	2001	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RS	\mathbf{RZ} >	>2	$OWAB(Å^2)$	Q < 0.9
1	А	1218/1235~(98%)	0.03	29 (2%)	59	64	11, 19, 36, 68	0
1	В	1217/1235~(98%)	0.08	44 (3%)	42	49	9, 18, 37, 74	0
All	All	2435/2470 (98%)	0.06	73 (2%)	50	57	9, 18, 37, 74	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	81	GLY	7.6
1	А	80 SER		6.9
1	В	80	SER	5.9
1	А	127	ASP	5.6
1	В	1231	ALA	4.8
1	В	14	ALA	4.6
1	А	82	SER	4.4
1	А	133	HIS	4.3
1	А	132	SER	4.3
1	А	506	PRO	4.2
1	В	81	GLY	4.1
1	В	918	THR	4.1
1	В	506	PRO	4.0
1	В	79	HIS	3.8
1	А	134	LEU	3.8
1	А	14	ALA	3.6
1	В	917	GLU	3.5
1	А	526	VAL	3.5
1	В	912	ILE	3.5
1	В	82	SER	3.4
1	A	79	HIS	3.4
1	В	1140	ASN	3.3
1	В	915	ALA	3.3
1	В	133	HIS	3.1



6X9D

Mol	Chain	Res	Type	RSRZ
1	В	438	ASP	3.0
1	В	15	PRO	3.0
1	А	122	ARG	3.0
1	А	130	TRP	2.9
1	В	60	ILE	2.9
1	А	120	LEU	2.8
1	А	1231	ALA	2.8
1	А	75	LEU	2.8
1	А	115	ALA	2.8
1	В	1233	GLY	2.8
1	А	116	THR	2.7
1	В	510	ILE	2.7
1	А	123	ASP	2.7
1	А	571	ALA	2.6
1	А	129	ASN	2.6
1	А	83	GLY	2.6
1	В	1230	MET	2.6
1	В	115	ALA	2.6
1	А	121	ILE	2.6
1	В	436	GLY	2.5
1	В	491	LEU	2.5
1	А	582	THR	2.5
1	В	508	VAL	2.5
1	В	914	LEU	2.5
1	В	919	GLY	2.5
1	В	223	LEU	2.5
1	В	916	SER	2.4
1	В	493	ASN	2.4
1	В	1059	GLY	2.4
1	В	129	ASN	2.4
1	В	75	LEU	2.4
1	В	1229	LEU	2.4
1	В	77	GLY	2.4
1	В	132	SER	2.3
1	В	800	PRO	2.3
1	В	19	ALA	2.3
1	В	1136	ALA	2.3
1	А	523	ALA	2.2
1	А	493	ASN	2.2
1	В	120	LEU	2.2
1	А	84	VAL	2.2
1	В	84	VAL	2.2



Mol	Chain	Res	Type	RSRZ
1	В	513	LEU	2.1
1	А	1233	GLY	2.1
1	В	514	ILE	2.1
1	В	439	PHE	2.0
1	В	1056	TRP	2.0
1	А	583	GLY	2.0
1	В	1232	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	PEG	В	2004	7/7	0.71	0.14	34,40,46,48	0
4	PGE	А	2003	10/10	0.88	0.12	33,43,52,54	0
4	PGE	В	2003	10/10	0.89	0.10	36,43,48,51	0
7	SO4	А	2008	5/5	0.90	0.13	29,33,38,42	5
3	HYP	А	2002	9/9	0.90	0.12	17,24,32,33	0
7	SO4	А	2007	5/5	0.91	0.18	29,31,35,46	5
3	HYP	В	2002	9/9	0.93	0.09	15,16,20,22	0
5	FMT	А	2004	3/3	0.93	0.09	29,29,29,30	0
6	NAD	А	2005	44/44	0.93	0.09	13,21,26,28	0
2	FAD	В	2001	53/53	0.94	0.09	13,17,31,33	0
8	MG	А	2010	1/1	0.94	0.10	29,29,29,29	0
2	FAD	А	2001	53/53	0.94	0.09	12,17,24,27	0
7	SO4	В	2007	5/5	0.95	0.22	55, 56, 57, 58	5
7	SO4	А	2009	5/5	0.96	0.15	30,37,42,47	5
6	NAD	В	2005	44/44	0.96	0.09	10,14,16,25	0
8	MG	B	2008	1/1	0.98	0.14	25,25,25,25	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
7	SO4	А	2006	5/5	0.99	0.05	$14,\!15,\!16,\!19$	0
7	SO4	В	2006	5/5	0.99	0.06	12,13,16,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.















6.5 Other polymers (i)

There are no such residues in this entry.

