

Full wwPDB X-ray Structure Validation Report (i)

Sep 19, 2023 – 08:08 PM JST

PDB ID : 7X9D

Title: DNMT3B in complex with harmine

Authors: Cho, C.-C.; Yuan, H.S.

Deposited on : 2022-03-15

Resolution : 3.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	1447 (3.10-3.06)
Clashscore	141614	1546 (3.10-3.06)
Ramachandran outliers	138981	1487 (3.10-3.06)
Sidechain outliers	138945	1486 (3.10-3.06)
RSRZ outliers	127900	1416 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	286	85%		12%	·
1	D	286	85%		12%	
2	В	204	72%	22%		6%
2	С	204	82%	1.	4%	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 3B.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	277	Total	С	N	О	S	0	0	0
1	11	211	2233	1432	394	394	13			
1	D	277	Total	С	N	О	S	0	0	0
1	ש	211	2233	1432	394	394	13	U	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	568	GLY	-	expression tag	UNP Q9UBC3
A	569	HIS	-	expression tag	UNP Q9UBC3
A	570	MET	-	expression tag	UNP Q9UBC3
D	568	GLY	-	expression tag	UNP Q9UBC3
D	569	HIS	-	expression tag	UNP Q9UBC3
D	570	MET	-	expression tag	UNP Q9UBC3

• Molecule 2 is a protein called DNA (cytosine-5)-methyltransferase 3-like.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
9	В	191	Total	С	N	О	S	0	0	0
	Ъ	191	1580	1032	270	274	4	0	0	0
2	С	197	Total	С	N	О	S	0	0	0
		197	1623	1057	278	283	5	U	U	U


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	176	GLY	-	expression tag	UNP Q9UJW3
В	177	HIS	-	expression tag	UNP Q9UJW3
С	176	GLY	-	expression tag	UNP Q9UJW3
С	177	HIS	-	expression tag	UNP Q9UJW3

• Molecule 3 is 7-METHOXY-1-METHYL-9H-BETA-CARBOLINE (three-letter code: HRM)



(formula: $\mathrm{C}_{13}\mathrm{H}_{12}\mathrm{N}_2\mathrm{O})$ (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 16			0	0
3	D	1	Total 16			0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA (cytosine-5)-methyltransferase 3B









4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	72.02Å 236.16Å 230.65Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 - 3.08	Depositor
rtesolution (A)	29.94 - 3.08	EDS
% Data completeness	78.9 (29.94-3.08)	Depositor
(in resolution range)	78.9 (29.94-3.08)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.13 (at 3.06Å)	Xtriage
Refinement program	PHENIX 1.19.2	Depositor
D D.	0.213 , 0.245	Depositor
R, R_{free}	0.212 , 0.243	DCC
R_{free} test set	1986 reflections (6.85%)	wwPDB-VP
Wilson B-factor (Å ²)	67.4	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 45.5	EDS
L-test for twinning ²	$ < L >=0.53, < L^2>=0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7701	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HRM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles		
IVIOI	Chain	RMSZ # Z > 5		RMSZ	# Z > 5	
1	A	0.25	0/2284	0.49	0/3081	
1	D	0.25	0/2284	0.49	0/3081	
2	В	0.25	0/1630	0.48	0/2217	
2	С	0.25	0/1673	0.51	0/2273	
All	All	0.25	0/7871	0.49	0/10652	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2233	0	2249	22	0
1	D	2233	0	2249	18	0
2	В	1580	0	1564	26	0
2	С	1623	0	1609	17	0
3	A	16	0	12	2	0
3	D	16	0	12	1	0
All	All	7701	0	7695	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A.,	A	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:699:VAL:HG12	1:D:701:ALA:H	1.56	0.68
2:B:204:THR:HG22	2:B:209:LEU:HB2	1.79	0.64
1:D:760:LEU:HA	1:D:804:GLU:HG2	1.80	0.63
2:C:283:MET:HG3	2:C:325:SER:HB2	1.80	0.62
1:A:586:THR:HG23	1:A:836:VAL:HG12	1.85	0.58
2:B:206:LEU:O	2:B:372:ARG:NH1	2.37	0.58
1:A:760:LEU:HA	1:A:804:GLU:HG2	1.86	0.57
1:A:627:ASP:HA	3:A:901:HRM:HAB2	1.86	0.56
1:A:685:LYS:HD2	1:A:685:LYS:H	1.71	0.56
2:B:194:SER:OG	2:B:197:GLU:O	2.21	0.56
2:B:285:VAL:HG13	2:B:323:VAL:HG12	1.88	0.56
2:C:247:PRO:HG3	2:C:359:TRP:CD1	2.43	0.54
1:A:628:VAL:HB	3:A:901:HRM:HAB3	1.90	0.53
1:A:693:PHE:HB3	1:A:739:LEU:HD23	1.91	0.53
2:B:192:VAL:HG23	2:B:240:LEU:HB3	1.91	0.53
2:B:288:LEU:HD21	2:B:343:LEU:HD22	1.90	0.53
2:B:305:GLU:HG3	2:B:306:PRO:HD2	1.91	0.52
1:A:689:ASP:OD1	1:A:689:ASP:N	2.39	0.52
2:C:183:PRO:O	2:C:187:ARG:HG3	2.10	0.52
2:B:290:LEU:HD22	2:B:294:ASP:HB3	1.92	0.51
2:B:306:PRO:HB3	2:B:324:TRP:CE2	2.45	0.51
2:C:247:PRO:HG3	2:C:359:TRP:HD1	1.76	0.51
2:B:325:SER:HB3	2:B:330:ILE:HD13	1.90	0.51
2:B:245:THR:HG22	2:B:286:ASP:HA	1.94	0.50
1:A:731:ARG:NH2	1:A:831:GLY:O	2.45	0.50
1:D:590:VAL:HG22	1:D:836:VAL:HG23	1.93	0.49
2:B:200:LYS:O	2:B:204:THR:HG23	2.12	0.49
1:A:700:VAL:HG23	1:A:734:TYR:CZ	2.47	0.49
2:C:238:PHE:O	2:C:278:ARG:NH1	2.42	0.49
1:A:804:GLU:O	1:A:808:ILE:HG13	2.14	0.48
2:B:311:ASP:HB2	2:B:363:LEU:HD21	1.96	0.48
2:B:307:VAL:HB	2:B:323:VAL:HG22	1.95	0.48
1:D:798:ASP:OD1	1:D:799:VAL:N	2.46	0.47
1:D:588:TYR:CE1	1:D:598:VAL:HG21	2.49	0.47
2:B:264:HIS:O	2:B:268:GLN:HG2	2.15	0.47
2:C:248:LEU:HD21	2:C:347:ALA:HA	1.96	0.47
1:A:578:LEU:HB3	1:A:645:VAL:HG22	1.97	0.46
1:A:674:GLU:OE2	2:B:300:ARG:NH2	2.48	0.46
1:A:726:VAL:HG12	1:A:748:ALA:HB2	1.98	0.46
2:C:206:LEU:HD11	2:C:365:LYS:HG3	1.97	0.46

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
2:C:287:ASN:HD22	2:C:321:VAL:HG12	1.81	0.46
1:A:655:SER:HA	1:A:701:ALA:O	2.16	0.45
2:B:294:ASP:HA	2:B:297:VAL:HG22	1.98	0.45
2:C:196:PHE:O	2:C:222:VAL:HG12	2.17	0.45
2:C:210:GLU:HB3	2:C:378:PHE:CD2	2.52	0.45
2:B:317:LEU:HG	2:B:346:LEU:HD12	1.97	0.45
1:D:700:VAL:HG23	1:D:734:TYR:CZ	2.52	0.45
1:D:613:VAL:HG12	1:D:818:VAL:HB	1.98	0.44
1:A:836:VAL:N	1:A:837:PRO:HD2	2.32	0.44
1:D:612:ALA:O	1:D:616:VAL:HG12	2.17	0.44
1:A:631:ILE:HA	1:A:635:ASN:HD22	1.83	0.44
1:A:739:LEU:HB2	1:A:742:MET:HG3	2.00	0.44
2:C:325:SER:HB3	2:C:330:ILE:HG12	1.99	0.44
1:A:661:ARG:HG2	1:A:663:GLY:H	1.83	0.43
2:B:183:PRO:HB2	2:B:185:TRP:CD1	2.53	0.43
1:A:590:VAL:HG21	1:A:839:ILE:HB	2.01	0.43
2:C:195:LEU:HD21	2:C:266:LEU:HD22	2.01	0.43
2:C:260:LEU:HD11	2:C:298:ALA:HB2	2.01	0.43
2:B:350:LYS:HA	2:B:350:LYS:HD2	1.80	0.43
1:D:836:VAL:N	1:D:837:PRO:HD2	2.33	0.43
2:B:195:LEU:C	2:B:197:GLU:H	2.22	0.43
1:D:597:LYS:HE3	1:D:853:GLU:HG3	2.01	0.42
1:A:750:LYS:HD3	1:A:750:LYS:HA	1.64	0.42
2:C:300:ARG:NH1	1:D:674:GLU:OE1	2.52	0.42
2:B:287:ASN:HD22	2:B:321:VAL:HG12	1.85	0.42
1:D:739:LEU:HB2	1:D:742:MET:HG3	2.02	0.42
2:B:196:PHE:HB2	2:B:244:ALA:O	2.20	0.42
1:A:801:TRP:CE2	1:D:819:SER:HA	2.55	0.42
2:B:196:PHE:HE2	2:B:263:PHE:HB2	1.86	0.41
1:D:724:ILE:HD12	1:D:730:HIS:CD2	2.55	0.41
1:D:646:ILE:HG21	1:D:843:PHE:HZ	1.85	0.41
1:D:723:ALA:HB3	1:D:731:ARG:HB3	2.03	0.41
1:A:801:TRP:NE1	1:D:819:SER:HA	2.35	0.41
2:B:307:VAL:O	2:B:322:ARG:HA	2.21	0.41
1:D:628:VAL:HB	3:D:901:HRM:HAB3	2.03	0.40
2:C:206:LEU:O	2:C:372:ARG:NH2	2.55	0.40
2:B:292:LYS:H	2:B:292:LYS:HD2	1.86	0.40
2:C:198:ASP:OD1	2:C:199:ILE:N	2.54	0.40
2:C:264:HIS:O	2:C:268:GLN:HG2	2.20	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	$273/286 \ (96\%)$	260 (95%)	13 (5%)	0	100	100
1	D	273/286~(96%)	263 (96%)	10 (4%)	0	100	100
2	В	185/204 (91%)	180 (97%)	5 (3%)	0	100	100
2	C	191/204 (94%)	182 (95%)	9 (5%)	0	100	100
All	All	922/980 (94%)	885 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	242/249 (97%)	239 (99%)	3 (1%)	71	87	
1	D	242/249 (97%)	239 (99%)	3 (1%)	71	87	
2	В	175/185 (95%)	173 (99%)	2 (1%)	73	88	
2	C	180/185 (97%)	178 (99%)	2 (1%)	73	88	
All	All	839/868 (97%)	829 (99%)	10 (1%)	71	87	

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	708	ARG
1	A	720	MET

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	819	SER
2	В	359	TRP
2	В	377	TYR
2	С	234	GLU
2	С	372	ARG
1	D	570	MET
1	D	676	TYR
1	D	689	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mal True Chain Das I		Link	Bond lengths			Bond angles			
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HRM	D	901	-	18,18,18	1.34	2 (11%)	21,26,26	1.25	2 (9%)
3	HRM	A	901	-	18,18,18	1.33	3 (16%)	21,26,26	1.36	4 (19%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HRM	D	901	-	-	2/2/2/2	0/3/3/3
3	HRM	A	901	-	-	2/2/2/2	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
3	D	901	HRM	CAB-CAL	3.04	1.52	1.50
3	A	901	HRM	CAB-CAL	2.72	1.52	1.50
3	A	901	HRM	CAE-CAC	2.62	1.40	1.36
3	D	901	HRM	CAE-CAC	2.49	1.40	1.36
3	A	901	HRM	CAO-CAP	-2.06	1.37	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	901	HRM	CAB-CAL-CAP	-2.79	117.09	121.74
3	A	901	HRM	CAE-CAC-NAH	-2.77	121.08	123.81
3	D	901	HRM	CAE-CAC-NAH	-2.75	121.11	123.81
3	A	901	HRM	CAE-CAO-CAN	2.21	137.99	133.15
3	A	901	HRM	CAA-OAJ-CAK	-2.06	113.03	117.51
3	D	901	HRM	CAB-CAL-CAP	-2.06	118.31	121.74

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	HRM	CAD-CAK-OAJ-CAA
3	A	901	HRM	CAG-CAK-OAJ-CAA
3	D	901	HRM	CAG-CAK-OAJ-CAA
3	D	901	HRM	CAD-CAK-OAJ-CAA


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	901	HRM	1	0
3	A	901	HRM	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	277/286~(96%)	-0.11	9 (3%) 47 25	31, 53, 101, 131	0
1	D	$277/286 \ (96\%)$	-0.05	9 (3%) 47 25	31, 52, 104, 125	0
2	В	191/204 (93%)	0.02	8 (4%) 36 18	36, 66, 110, 141	0
2	С	197/204 (96%)	0.14	10 (5%) 28 12	35, 64, 116, 139	0
All	All	942/980 (96%)	-0.01	36 (3%) 40 20	31, 58, 108, 141	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	С	275	GLY	5.4
2	С	274	PRO	4.0
1	A	657	VAL	3.5
2	В	180	GLU	3.5
2	С	185	TRP	3.4
1	D	778	SER	3.3
2	С	352	SER	3.2
1	A	787	GLN	3.1
2	С	180	GLU	3.0
1	D	659	PRO	2.9
1	A	687	GLY	2.9
2	В	251	THR	2.9
2	С	181	THR	2.9
2	В	182	VAL	2.7
1	A	786	ASN	2.6
1	D	652	ASN	2.6
1	D	656	ASN	2.6
1	A	660	ALA	2.5
1	A	853	GLU	2.5
1	D	657	VAL	2.5
2	С	182	VAL	2.5

Continued on next page...



Continued from previous page...

Mol	Chain	Res Type		RSRZ	
1	D	853	GLU	2.5	
2	В	185	TRP	2.4	
2	С	217	GLN	2.4	
1	D	660	ALA	2.4	
2	В	276	SER	2.3	
1	D	658	ASN	2.3	
2	В	181	THR	2.3	
2	С	249	GLY	2.2	
1	A	659	PRO	2.2	
2	В	275	GLY	2.2	
1	A	688	ASP	2.1	
1	A	777	LYS	2.1	
1	D	686	GLU	2.1	
2	В	376	LYS	2.0	
2	С	212	GLY	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	HRM	A	901	16/16	0.83	0.28	47,70,91,104	0
3	HRM	D	901	16/16	0.89	0.22	54,69,94,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

