



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 10:08 PM JST

PDB ID : 7X9E  
Title : Crystal structure of the 76E1 Fab in complex with a SARS-CoV-2 spike peptide  
Authors : Chen, X.; Zhang, T.; Ding, J.; Sun, X.; Sun, B.  
Deposited on : 2022-03-15  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
1	C	221	
2	B	216	
2	D	216	
3	E	25	
3	F	25	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 76E1 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	211	1577	995	267	309	6	0	0	0
1	C	211	1576	993	267	310	6	0	0	0

- Molecule 2 is a protein called 76E1 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	211	1575	987	265	317	6	0	0	0
2	D	210	1564	981	263	314	6	0	0	0

- Molecule 3 is a protein called Spike peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	13	112	73	19	20	0	0	0
3	F	15	126	83	21	22	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	18	Total	O	0	0
			18	18		
4	B	28	Total	O	0	0
			28	28		
4	C	25	Total	O	0	0
			25	25		
4	D	20	Total	O	0	0
			20	20		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	E	1	Total	O	0	0
			1	1		






- Molecule 3: Spike peptide



- Molecule 3: Spike peptide



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	130.36Å 84.52Å 87.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.88 – 2.60 44.40 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (38.88-2.60) 99.0 (44.40-2.60)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.195 , 0.243 0.195 , 0.243	Depositor DCC
$R_{free}$ test set	1499 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 28.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.021 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6622	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/1611	0.65	0/2192
1	C	0.45	0/1609	0.65	0/2188
2	B	0.51	0/1614	0.62	0/2200
2	D	0.53	0/1603	0.61	0/2185
3	E	0.43	0/113	0.60	0/148
3	F	0.42	0/128	0.56	0/169
All	All	0.48	0/6678	0.63	0/9082

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	97	ASN	Peptide
2	D	97	ASN	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1577	0	1544	21	0
1	C	1576	0	1542	22	0
2	B	1575	0	1532	19	0
2	D	1564	0	1519	11	0
3	E	112	0	115	4	0
3	F	126	0	132	0	0
4	A	18	0	0	1	0
4	B	28	0	0	2	0
4	C	25	0	0	1	0
4	D	20	0	0	0	0
4	E	1	0	0	0	0
All	All	6622	0	6384	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2:VAL:N	4:C:301:HOH:O	2.25	0.70
1:C:67:ARG:NH2	1:C:90:ASP:OD2	2.24	0.69
1:A:22:CYS:HB3	1:A:79:LEU:HB3	1.77	0.67
1:A:67:ARG:NH2	1:A:90:ASP:OD2	2.29	0.65
1:C:181:GLN:HG2	2:D:164:GLU:HG3	1.79	0.65
1:A:44:GLY:N	4:A:301:HOH:O	2.30	0.64
1:A:131:VAL:O	1:A:219:LYS:HD2	2.00	0.62
2:D:85:GLU:HB2	2:D:109:VAL:HG23	1.81	0.61
1:A:101:ILE:HD12	2:B:34:PHE:HB3	1.82	0.61
1:C:34:MET:HB3	1:C:79:LEU:HD22	1.82	0.60
2:B:51:TYR:HD1	2:B:52:GLU:HG3	1.66	0.59
1:C:83:MET:HB3	1:C:86:LEU:HD21	1.86	0.56
1:A:101:ILE:HG13	1:A:102:VAL:N	2.19	0.56
1:A:62:ASP:HA	1:A:65:LYS:HE3	1.87	0.56
1:A:28:SER:OG	1:A:31:ASP:OD2	2.21	0.56
2:B:39:ARG:NH1	4:B:302:HOH:O	2.35	0.56
1:C:129:PRO:HB3	1:C:155:TYR:HB3	1.88	0.55
1:A:83:MET:HE2	1:A:86:LEU:HD21	1.88	0.55
2:D:193:ARG:NH2	2:D:193:ARG:HB2	2.22	0.55
1:A:63:SER:O	1:A:67:ARG:NH1	2.41	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:73:ASP:OD1	1:C:75:SER:OG	2.27	0.53
1:A:100:VAL:HG12	1:A:111:ASP:HB2	1.91	0.52
1:A:172:GLY:O	1:A:192:VAL:HA	2.10	0.51
1:C:179:VAL:HG12	2:D:166:THR:OG1	2.11	0.51
2:D:96:ARG:C	2:D:98:ASN:HB2	2.32	0.50
2:D:63:ARG:HD2	2:D:78:SER:O	2.11	0.50
2:D:168:PRO:HA	2:D:177:ALA:O	2.12	0.49
2:B:153:LYS:HD2	2:B:198:GLN:NE2	2.25	0.49
1:A:102:VAL:HG21	3:E:818:ILE:HG12	1.93	0.49
1:C:67:ARG:HH22	1:C:90:ASP:CG	2.14	0.47
1:C:89:GLU:N	1:C:89:GLU:OE1	2.47	0.47
3:E:814:LYS:O	3:E:818:ILE:HG23	2.14	0.47
2:D:114:LYS:HD3	2:D:145:PRO:HD3	1.97	0.47
2:B:26:SER:O	2:B:31:SER:OG	2.31	0.47
2:D:166:THR:HG23	2:D:167:THR:O	2.15	0.47
2:D:37:TRP:CE2	2:D:75:LEU:HB2	2.51	0.46
1:C:62:ASP:HA	1:C:65:LYS:HE3	1.98	0.46
2:B:81:GLN:O	2:B:109:VAL:HG21	2.15	0.46
1:C:68:PHE:HB3	1:C:81:LEU:HD11	1.98	0.46
2:B:37:TRP:CZ3	2:B:90:CYS:HB3	2.51	0.45
2:B:51:TYR:CD1	2:B:52:GLU:HG3	2.50	0.45
1:C:65:LYS:HE3	1:C:65:LYS:HB2	1.61	0.45
1:C:172:GLY:O	1:C:192:VAL:HA	2.16	0.45
2:B:127:GLU:OE1	2:B:127:GLU:N	2.43	0.45
2:B:151:ALA:HB3	2:B:198:GLN:HB2	1.99	0.45
1:C:100:VAL:CG2	1:C:103:ALA:HB2	2.47	0.45
2:B:40:GLN:NE2	4:B:307:HOH:O	2.49	0.45
3:E:821:LEU:HD12	3:E:821:LEU:HA	1.59	0.45
2:B:207:GLU:HG3	2:B:208:LYS:N	2.32	0.45
1:C:99:LEU:HD21	1:C:113:TRP:CZ2	2.52	0.44
1:A:180:LEU:HD13	1:A:186:TYR:CZ	2.53	0.44
2:B:170:LYS:HE3	2:B:176:TYR:OH	2.17	0.44
1:C:89:GLU:H	1:C:89:GLU:CD	2.20	0.43
2:B:153:LYS:HB2	2:B:196:SER:HB2	2.00	0.43
2:B:108:THR:HG21	2:B:145:PRO:HB3	1.99	0.43
1:C:153:LYS:HE3	1:C:154:ASP:OD2	2.19	0.43
2:D:193:ARG:HB2	2:D:193:ARG:CZ	2.49	0.42
1:C:169:LEU:HD21	1:C:192:VAL:HG21	2.02	0.42
1:A:38:ARG:HA	1:A:93:LEU:O	2.19	0.42
1:C:127:LYS:HD3	1:C:185:LEU:HD13	2.00	0.42
1:A:102:VAL:HG11	3:E:818:ILE:HG13	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:96:ARG:O	2:B:98:ASN:HB2	2.20	0.41
2:B:160:LYS:HE2	2:B:160:LYS:HB2	1.94	0.41
1:A:131:VAL:HA	1:A:151:LEU:O	2.21	0.41
1:A:153:LYS:NZ	2:B:133:LYS:HG2	2.35	0.41
2:B:96:ARG:C	2:B:98:ASN:HB2	2.41	0.41
1:C:12:VAL:O	1:C:121:VAL:HA	2.19	0.41
1:C:179:VAL:O	1:C:186:TYR:HA	2.20	0.41
1:A:53:GLY:HA2	1:A:72:ARG:NH1	2.37	0.40
1:A:54:ASP:OD2	1:A:56:ARG:HD2	2.21	0.40
1:A:4:LEU:HD22	1:A:22:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/221 (93%)	197 (96%)	8 (4%)	0	100	100
1	C	205/221 (93%)	198 (97%)	7 (3%)	0	100	100
2	B	209/216 (97%)	196 (94%)	13 (6%)	0	100	100
2	D	208/216 (96%)	197 (95%)	11 (5%)	0	100	100
3	E	11/25 (44%)	11 (100%)	0	0	100	100
3	F	13/25 (52%)	12 (92%)	1 (8%)	0	100	100
All	All	851/924 (92%)	811 (95%)	40 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/186 (96%)	160 (90%)	18 (10%)	7	14
1	C	178/186 (96%)	157 (88%)	21 (12%)	5	9
2	B	178/183 (97%)	162 (91%)	16 (9%)	9	18
2	D	176/183 (96%)	155 (88%)	21 (12%)	5	9
3	E	13/22 (59%)	8 (62%)	5 (38%)	0	0
3	F	15/22 (68%)	15 (100%)	0	100	100
All	All	738/782 (94%)	657 (89%)	81 (11%)	6	11

All (81) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LEU
1	A	23	GLU
1	A	59	SER
1	A	101	ILE
1	A	115	GLN
1	A	125	SER
1	A	145	THR
1	A	148	LEU
1	A	150	CYS
1	A	159	PRO
1	A	170	THR
1	A	171	SER
1	A	174	HIS
1	A	181	GLN
1	A	197	SER
1	A	202	GLN
1	A	206	CYS
1	A	211	LYS
2	B	8	LEU
2	B	13	SER
2	B	42	PRO
2	B	62	VAL

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
2	B	80	LEU
2	B	83	GLU
2	B	99	LEU
2	B	114	LYS
2	B	148	VAL
2	B	157	SER
2	B	160	LYS
2	B	169	SER
2	B	174	ASN
2	B	183	SER
2	B	204	SER
2	B	207	GLU
1	C	11	VAL
1	C	13	GLN
1	C	18	LEU
1	C	55	THR
1	C	56	ARG
1	C	63	SER
1	C	88	SER
1	C	89	GLU
1	C	96	CYS
1	C	115	GLN
1	C	121	VAL
1	C	150	CYS
1	C	158	GLU
1	C	174	HIS
1	C	182	SER
1	C	191	VAL
1	C	192	VAL
1	C	193	THR
1	C	196	SER
1	C	206	CYS
1	C	219	LYS
2	D	4	LEU
2	D	17	SER
2	D	18	VAL
2	D	27	SER
2	D	28	ASP
2	D	67	SER
2	D	74	SER
2	D	80	LEU
2	D	96	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	98	ASN
2	D	107	LEU
2	D	125	SER
2	D	133	LYS
2	D	148	VAL
2	D	149	THR
2	D	159	VAL
2	D	166	THR
2	D	179	SER
2	D	190	LYS
2	D	204	SER
2	D	208	LYS
3	E	814	LYS
3	E	816	SER
3	E	818	ILE
3	E	821	LEU
3	E	825	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	171	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/221 (95%)	-0.15	3 (1%) 75 71	17, 32, 59, 82	0
1	C	211/221 (95%)	-0.42	0 100 100	18, 30, 48, 64	0
2	B	211/216 (97%)	-0.33	0 100 100	19, 32, 46, 60	0
2	D	210/216 (97%)	-0.25	2 (0%) 82 80	19, 38, 55, 67	0
3	E	13/25 (52%)	-0.14	1 (7%) 13 10	31, 36, 50, 52	0
3	F	15/25 (60%)	-0.24	0 100 100	33, 39, 54, 62	0
All	All	871/924 (94%)	-0.29	6 (0%) 87 86	17, 33, 53, 82	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	199	LEU	4.4
2	D	34	PHE	3.2
2	D	119	VAL	2.9
1	A	200	GLY	2.9
1	A	201	THR	2.1
3	E	817	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.