



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2023 – 11:08 am GMT

PDB ID : 2XA8
Title : Crystal structure of the Fab domain of omalizumab at 2.41Å
Authors : Huang, C.H.; Hung, F.H.A.; Lim, C.; Chang, T.W.; Ma, C.
Deposited on : 2010-03-30
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	219	
2	L	218	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OMALIZUMAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	219	1643	1038	279	321	5	0	0	0

- Molecule 2 is a protein called OMALIZUMAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	218	1682	1048	278	350	6	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	25	Total	O	0	0
			25	25		
3	L	47	Total	O	0	0
			47	47		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: OMALIZUMAB HEAVY CHAIN



- Molecule 2: OMALIZUMAB LIGHT CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.60Å 73.85Å 141.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.93 – 2.42 36.93 – 2.42	Depositor EDS
% Data completeness (in resolution range)	97.7 (36.93-2.42) 97.8 (36.93-2.42)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.42Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.6.3_473)	Depositor
R, R_{free}	0.221 , 0.269 0.211 , 0.261	Depositor DCC
R_{free} test set	1311 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3397	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.33	0/1687	0.51	0/2301
2	L	0.34	0/1719	0.52	0/2333
All	All	0.33	0/3406	0.51	0/4634

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	98	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1643	0	1594	41	0
2	L	1682	0	1605	46	0
3	H	25	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	47	0	0	1	0
All	All	3397	0	3199	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:164:SER:H	1:H:205:ASN:HD21	1.25	0.84
1:H:163:ASN:HD21	1:H:203:ILE:H	1.26	0.80
2:L:137:VAL:HG22	2:L:182:THR:HG22	1.65	0.79
2:L:83:GLN:HB3	2:L:85:GLU:OE2	1.83	0.79
1:H:163:ASN:ND2	1:H:203:ILE:H	1.83	0.76
2:L:15:VAL:O	3:L:2003:HOH:O	2.03	0.74
2:L:40:TYR:HE2	2:L:93:GLN:HE21	1.35	0.74
2:L:6:GLN:H	2:L:104:GLN:HE22	1.37	0.73
1:H:134:PRO:HG3	1:H:146:LEU:HB3	1.72	0.70
1:H:40:GLN:HE22	2:L:42:GLN:HE22	1.38	0.69
2:L:41:GLN:HB2	2:L:51:LEU:HD11	1.75	0.67
1:H:141:GLY:O	1:H:143:THR:N	2.29	0.65
2:L:104:GLN:HE21	2:L:104:GLN:H	1.45	0.64
2:L:1:ASP:O	2:L:1:ASP:CG	2.35	0.64
1:H:89:GLU:HG2	1:H:89:GLU:O	1.96	0.64
2:L:202:HIS:CD2	2:L:204:GLY:H	2.15	0.63
1:H:68:ILE:HG23	1:H:83:MET:HG2	1.80	0.63
1:H:164:SER:H	1:H:205:ASN:ND2	1.96	0.62
2:L:117:PRO:HD3	2:L:202:HIS:CD2	2.34	0.62
2:L:20:THR:HG22	2:L:78:THR:HB	1.80	0.62
1:H:18:LEU:HD23	1:H:18:LEU:O	2.00	0.61
2:L:85:GLU:CD	2:L:85:GLU:H	2.05	0.60
1:H:62:PRO:CD	2:L:99:PRO:HG3	2.32	0.59
2:L:38:ASN:HD22	2:L:53:TYR:HA	1.68	0.59
1:H:29:ILE:HG13	1:H:77:ASN:ND2	2.18	0.58
1:H:11:LEU:HD12	1:H:124:THR:HG22	1.86	0.57
1:H:64:VAL:O	1:H:68:ILE:HD11	2.03	0.57
1:H:53:THR:HG22	1:H:54:TYR:N	2.19	0.57
2:L:127:GLU:O	2:L:130:LYS:HG2	2.05	0.56
2:L:179:LEU:C	2:L:179:LEU:HD23	2.27	0.55
2:L:123:PRO:HB3	2:L:218:CYS:HB2	1.88	0.54
1:H:209:LYS:HA	1:H:209:LYS:HE2	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:62:PRO:HD3	2:L:99:PRO:HG3	1.90	0.52
2:L:104:GLN:H	2:L:104:GLN:NE2	2.05	0.52
1:H:65:LYS:HG3	1:H:66:GLY:H	1.75	0.51
1:H:29:ILE:CG1	1:H:77:ASN:HA	2.41	0.51
2:L:6:GLN:HA	2:L:22:THR:O	2.09	0.51
1:H:18:LEU:CD2	1:H:83:MET:HB2	2.41	0.51
1:H:28:SER:HA	1:H:77:ASN:HD21	1.75	0.51
1:H:18:LEU:HD22	1:H:83:MET:HB2	1.93	0.50
1:H:43:GLY:O	1:H:44:LYS:CB	2.60	0.50
2:L:147:GLU:H	2:L:147:GLU:CD	2.16	0.49
1:H:132:LEU:HB2	1:H:147:GLY:C	2.33	0.49
1:H:43:GLY:O	1:H:44:LYS:HB2	2.13	0.49
1:H:98:ARG:HD3	3:H:2009:HOH:O	2.11	0.49
2:L:109:GLU:HG2	2:L:110:ILE:N	2.28	0.49
2:L:65:ARG:HG3	2:L:79:ILE:HG23	1.94	0.48
1:H:65:LYS:HG3	1:H:66:GLY:N	2.29	0.48
2:L:1:ASP:N	2:L:2:ILE:HA	2.29	0.48
1:H:18:LEU:HD23	1:H:83:MET:H	1.79	0.47
2:L:184:THR:O	2:L:185:LEU:HD23	2.13	0.47
2:L:195:VAL:HG22	2:L:214:ASN:HD21	1.79	0.47
2:L:144:TYR:CG	2:L:145:PRO:HA	2.49	0.47
2:L:195:VAL:HG22	2:L:214:ASN:ND2	2.29	0.47
1:H:67:ARG:O	1:H:83:MET:HA	2.14	0.47
2:L:76:THR:HG22	2:L:78:THR:HG22	1.97	0.46
1:H:53:THR:HG22	1:H:54:TYR:H	1.80	0.46
2:L:83:GLN:O	2:L:86:ASP:HB2	2.14	0.46
2:L:44:PRO:CG	2:L:169:GLU:HG3	2.46	0.45
2:L:191:GLU:HA	2:L:191:GLU:OE1	2.17	0.45
2:L:84:PRO:HD2	2:L:85:GLU:OE2	2.17	0.45
1:H:40:GLN:HE22	2:L:42:GLN:NE2	2.13	0.44
2:L:189:ASP:O	2:L:193:HIS:HD2	2.00	0.44
1:H:29:ILE:HG12	1:H:77:ASN:HA	1.98	0.43
1:H:175:PRO:HB2	2:L:166:SER:OG	2.18	0.43
2:L:116:ALA:HA	2:L:117:PRO:HD3	1.90	0.43
1:H:35:TRP:CZ3	1:H:98:ARG:HB2	2.54	0.42
2:L:149:LYS:HE3	2:L:151:GLN:HG3	2.00	0.42
1:H:4:LEU:HB3	1:H:22:CYS:SG	2.60	0.42
1:H:127:PRO:HB3	1:H:153:TYR:HB3	2.01	0.42
2:L:107:LYS:HD2	2:L:107:LYS:HA	1.87	0.42
2:L:111:LYS:HA	2:L:144:TYR:OH	2.20	0.42
1:H:18:LEU:HD23	1:H:18:LEU:C	2.40	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:53:THR:CG2	1:H:54:TYR:N	2.84	0.41
1:H:3:GLN:HG2	1:H:25:SER:HB2	2.03	0.41
1:H:163:ASN:HD21	1:H:203:ILE:N	2.06	0.41
2:L:39:TRP:CD2	2:L:77:LEU:HB2	2.56	0.41
2:L:145:PRO:HB2	2:L:147:GLU:OE2	2.21	0.41
2:L:190:TYR:CZ	2:L:215:ARG:HG3	2.56	0.41
2:L:162:ASN:HD22	2:L:162:ASN:HA	1.59	0.40
2:L:39:TRP:CE2	2:L:77:LEU:HB2	2.56	0.40
1:H:6:GLU:H	1:H:113:GLN:HE22	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	217/219 (99%)	200 (92%)	14 (6%)	3 (1%)	11	14
2	L	216/218 (99%)	204 (94%)	10 (5%)	2 (1%)	17	24
All	All	433/437 (99%)	404 (93%)	24 (6%)	5 (1%)	13	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	44	LYS
2	L	99	PRO
1	H	65	LYS
1	H	142	GLY
2	L	98	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	183/183 (100%)	163 (89%)	20 (11%)	6	8
2	L	192/192 (100%)	184 (96%)	8 (4%)	30	46
All	All	375/375 (100%)	347 (92%)	28 (8%)	13	20

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	1	GLU
1	H	3	GLN
1	H	11	LEU
1	H	13	GLN
1	H	18	LEU
1	H	20	LEU
1	H	25	SER
1	H	29	ILE
1	H	44	LYS
1	H	67	ARG
1	H	68	ILE
1	H	86	LEU
1	H	89	GLU
1	H	132	LEU
1	H	143	THR
1	H	158	VAL
1	H	159	THR
1	H	199	THR
1	H	201	THR
1	H	205	ASN
2	L	1	ASP
2	L	15	VAL
2	L	73	THR
2	L	78	THR
2	L	93	GLN
2	L	104	GLN
2	L	107	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	162	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	3	GLN
1	H	40	GLN
1	H	77	ASN
1	H	163	ASN
1	H	179	GLN
1	H	200	GLN
1	H	205	ASN
2	L	38	ASN
2	L	93	GLN
2	L	104	GLN
2	L	128	GLN
2	L	162	ASN
2	L	193	HIS
2	L	202	HIS
2	L	203	GLN
2	L	214	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	219/219 (100%)	0.37	17 (7%) 13 11	26, 42, 76, 117	0
2	L	218/218 (100%)	0.15	7 (3%) 47 45	26, 40, 66, 129	0
All	All	437/437 (100%)	0.26	24 (5%) 25 23	26, 41, 71, 129	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	218	CYS	6.1
1	H	65	LYS	5.2
1	H	140	SER	4.9
2	L	1	ASP	4.1
2	L	31	TYR	3.9
1	H	13	GLN	3.5
1	H	12	VAL	3.0
1	H	119	VAL	3.0
2	L	217	GLU	2.9
1	H	17	SER	2.8
1	H	76	LYS	2.6
1	H	15	GLY	2.6
2	L	130	LYS	2.4
1	H	85	SER	2.4
1	H	136	SER	2.3
1	H	1	GLU	2.3
1	H	84	ASN	2.3
2	L	32	ASP	2.2
1	H	44	LYS	2.1
1	H	141	GLY	2.1
1	H	87	ARG	2.1
1	H	139	THR	2.0
2	L	206	SER	2.0
1	H	11	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.