

Full wwPDB NMR Structure Validation Report (i)

Aug 21, 2022 – 08:17 AM EDT

PDB ID : 2XBD

Title : INTERNAL XYLAN BINDING DOMAIN FROM CELLULOMONAS FIMI

XYLANASE D, NMR, MINIMIZED AVERAGE STRUCTURE

Authors: Simpson, P.J.; Bolam, D.N.; Cooper, A.; Ciruela, A.; Hazlewood, G.P.;

Gilbert, H.J.; Williamson, M.P.

Deposited on : 1998-10-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.29

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	87	100%



2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1192 atoms, of which 577 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called XYLANASE D.

Mol	Chain	Residues		Atoms								
1	Λ	97	Total	С	Н	N	О	S	0			
1	A	01	1192	368	577	110	134	3	U			



4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: XYLANASE D

C	Ch	a	in	۱.	A	•]	LO	0%	6																							
T247	G248	C249	S250	V251	T252	T254	R255	A256	E257	õ	W259	S260	D261	R262	F263	Ñ	26		Y267		V269	S270	G271	S272	2	A274	W275	\sim	2	N278	N	Ò	ຸ ເ	N282	8 8	Ńο	T286	1 0	ı Ö	A289	õi	Ši.	Ši.	Ši .	Ši i	53	G297	N d	T301	R302	T303	m	T305	m
N307	808	8	31	N311	T312	G314	3 1	T316	31.	M318	K319	N320	32	S322	S323	T324	T325	P326	32	32	32	33	A331	က်	2333																													



Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: HYBRID DISTANCE GEOMETRY/ SIM-ULATED ANNEALING.

Of the 50 calculated structures, 1 were deposited, based on the following criterion: MINIMIZED AVERAGE STRUCTURE, CALCULATED FROM 38 LOWEST ENERGY STRUCTURES.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version					
X-PLOR	refinement	3.1					
MSI FELIX	structure solution	97.0					

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes				
1	A	0	0	0	0				
All	All	0	0	0	-				

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	-	-	-
All	All	0	-	-	-	-

There are no Ramachandran outliers.



6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

