



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 22, 2023 – 06:50 PM JST

PDB ID : 7XB3
Title : Crystal structure of SARS-Cov-2 main protease D48N mutant
Authors : Hu, X.H.; Li, J.; Zhang, J.
Deposited on : 2022-03-20
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replicase polyprotein 1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	2199	1396	371	412	20	0	0	0
1	B	286	2137	1359	359	399	20	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	ASN	ASP	engineered mutation	UNP P0DTC1
B	48	ASN	ASP	engineered mutation	UNP P0DTC1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	48	Total	O	0	0
			48	48		
2	B	31	Total	O	0	0
			31	31		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Replicase polyprotein 1a



- Molecule 1: Replicase polyprotein 1a



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.05Å 102.94Å 103.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.31 – 2.08 56.89 – 2.08	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.31-2.08) 93.6 (56.89-2.08)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.08Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.214 , 0.242 0.213 , 0.242	Depositor DCC
R_{free} test set	2004 reflections (4.48%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtrriage
Anisotropy	0.546	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.010 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4415	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	2/2248 (0.1%)	0.60	1/3068 (0.0%)
1	B	0.44	1/2184 (0.0%)	0.56	0/2978
All	All	0.46	3/4432 (0.1%)	0.58	1/6046 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	CYS	CB-SG	-10.60	1.64	1.82
1	B	128	CYS	CB-SG	-8.20	1.68	1.82
1	A	145	CYS	CB-SG	-5.31	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	LEU	CA-CB-CG	-5.03	103.73	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2199	0	2093	17	0
1	B	2137	0	2035	25	0
2	A	48	0	0	0	0
2	B	31	0	0	0	0
All	All	4415	0	4128	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:MET:HE1	1:B:285:ALA:O	1.88	0.72
1:A:139:SER:HB2	1:B:4:ARG:HB2	1.72	0.72
1:B:141:LEU:H	1:B:141:LEU:HD23	1.60	0.66
1:A:169:THR:HG23	1:A:171:VAL:HG22	1.78	0.64
1:B:276:MET:CE	1:B:285:ALA:C	2.68	0.61
1:A:140:PHE:HD2	1:A:172:HIS:CD2	2.19	0.61
1:B:276:MET:HE2	1:B:285:ALA:C	2.24	0.58
1:A:4:ARG:HH12	1:B:5:LYS:NZ	2.03	0.57
1:B:188:ARG:NH2	1:B:190:THR:OG1	2.39	0.56
1:B:276:MET:CE	1:B:285:ALA:O	2.56	0.53
1:A:108:PRO:HB3	1:A:132:PRO:HA	1.91	0.53
1:A:140:PHE:HD2	1:A:172:HIS:CG	2.27	0.53
1:A:165:MET:HE1	1:A:187:ASP:HA	1.91	0.52
1:A:288:GLU:HG2	1:A:291:PHE:CE2	2.46	0.51
1:B:12:LYS:NZ	1:B:155:ASP:OD2	2.44	0.51
1:B:276:MET:HE1	1:B:286:LEU:HA	1.92	0.51
1:B:86:VAL:HG13	1:B:179:GLY:HA2	1.93	0.51
1:A:4:ARG:HB2	1:B:139:SER:HB3	1.93	0.50
1:B:5:LYS:HG2	1:B:291:PHE:CE2	2.46	0.50
1:A:86:VAL:HG13	1:A:179:GLY:HA2	1.92	0.50
1:A:4:ARG:HH12	1:B:5:LYS:HZ2	1.58	0.49
1:B:266:ALA:O	1:B:270:GLU:HG2	2.13	0.49
1:A:95:ASN:HB3	1:A:98:THR:OG1	2.13	0.49
1:A:288:GLU:HG2	1:A:291:PHE:HE2	1.78	0.49
1:B:276:MET:HE1	1:B:285:ALA:C	2.31	0.49
1:B:140:PHE:HB2	1:B:172:HIS:CE1	2.49	0.48
1:B:219:PHE:O	1:B:267:SER:OG	2.28	0.48
1:B:108:PRO:HB3	1:B:132:PRO:HA	1.98	0.46
1:A:4:ARG:N	1:B:139:SER:HB2	2.30	0.46
1:B:276:MET:O	1:B:279:ARG:HB2	2.19	0.43
1:B:57:LEU:HD22	1:B:57:LEU:H	1.83	0.42
1:B:113:SER:O	1:B:149:GLY:HA2	2.20	0.41
1:A:207:TRP:NE1	1:A:282:LEU:HD12	2.36	0.41
1:A:169:THR:CG2	1:A:171:VAL:HG22	2.49	0.41
1:B:227:LEU:HD12	1:B:227:LEU:HA	1.91	0.41
1:A:279:ARG:HE	1:A:279:ARG:HB3	1.75	0.40
1:B:114:VAL:HG11	1:B:140:PHE:CZ	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/306 (96%)	284 (97%)	8 (3%)	1 (0%)	41	39
1	B	282/306 (92%)	278 (99%)	4 (1%)	0	100	100
All	All	575/612 (94%)	562 (98%)	12 (2%)	1 (0%)	47	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/263 (89%)	229 (98%)	4 (2%)	60	65
1	B	226/263 (86%)	217 (96%)	9 (4%)	31	31
All	All	459/526 (87%)	446 (97%)	13 (3%)	43	46

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	128	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	165	MET
1	A	216	ASP
1	B	81	SER
1	B	121	SER
1	B	140	PHE
1	B	141	LEU
1	B	181	PHE
1	B	216	ASP
1	B	267	SER
1	B	276	MET
1	B	277	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	214	ASN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/306 (96%)	0.32	8 (2%) 54 59	26, 38, 59, 69	0
1	B	286/306 (93%)	0.59	22 (7%) 13 16	27, 47, 71, 82	0
All	All	581/612 (94%)	0.46	30 (5%) 27 31	26, 41, 64, 82	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	140	PHE	7.3
1	B	140	PHE	4.8
1	B	56	ASP	4.1
1	A	141	LEU	3.9
1	B	58	LEU	3.9
1	A	142	ASN	3.9
1	B	154	TYR	3.6
1	B	286	LEU	3.6
1	B	278	GLY	3.1
1	A	291	PHE	3.1
1	B	276	MET	2.9
1	B	155	ASP	2.8
1	B	57	LEU	2.8
1	B	60	ARG	2.8
1	B	62	SER	2.7
1	B	191	ALA	2.7
1	B	139	SER	2.6
1	B	188	ARG	2.6
1	B	232	LEU	2.6
1	A	282	LEU	2.6
1	B	141	LEU	2.5
1	B	128	CYS	2.5
1	A	213	ILE	2.5
1	A	281	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	189	GLN	2.4
1	B	61	LYS	2.4
1	B	294	PHE	2.3
1	B	235	MET	2.2
1	A	136	ILE	2.1
1	B	297	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.