



wwPDB EM Validation Summary Report ⓘ

Nov 29, 2022 – 10:53 PM JST

PDB ID : 7XBX
EMDB ID : EMD-33108
Title : Cryo-EM structure of the human chemokine receptor CX3CR1 in complex with CX3CL1 and Gi1
Authors : Lu, M.; Zhao, W.; Han, S.; Zhu, Y.; Wu, B.; Zhao, Q.
Deposited on : 2022-03-22
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	354	
2	D	346	
3	E	71	
4	R	463	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12644 atoms, of which 6028 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	C	216	3128	1038	1506	264	307	13	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	47	CYS	SER	engineered mutation	UNP P63096
C	202	THR	GLY	engineered mutation	UNP P63096
C	203	ALA	GLY	engineered mutation	UNP P63096
C	245	ALA	GLU	engineered mutation	UNP P63096
C	326	SER	ALA	engineered mutation	UNP P63096

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	D	311	4293	1402	2050	378	446	17	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	expression tag	UNP P62873
D	-4	HIS	-	expression tag	UNP P62873
D	-3	HIS	-	expression tag	UNP P62873
D	-2	HIS	-	expression tag	UNP P62873
D	-1	HIS	-	expression tag	UNP P62873
D	0	HIS	-	expression tag	UNP P62873

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	E	33	414	141	192	36	44	1	0	0

- Molecule 4 is a protein called Processed fractalkine, CX3C chemokine receptor 1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
4	R	349	4753	1625	2280	402	432	14	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	35	CYS	GLY	engineered mutation	UNP P78423
R	77	SER	-	linker	UNP P78423
R	78	GLY	-	linker	UNP P78423
R	79	SER	-	linker	UNP P78423
R	80	GLY	-	linker	UNP P78423
R	81	SER	-	linker	UNP P78423
R	82	GLY	-	linker	UNP P78423
R	83	SER	-	linker	UNP P78423
R	84	GLY	-	linker	UNP P78423
R	85	SER	-	linker	UNP P78423
R	86	GLY	-	linker	UNP P78423
R	87	SER	-	linker	UNP P78423
R	88	GLY	-	linker	UNP P78423
R	89	SER	-	linker	UNP P78423
R	90	GLY	-	linker	UNP P78423
R	91	SER	-	linker	UNP P78423
R	92	GLY	-	linker	UNP P78423
R	93	SER	-	linker	UNP P78423
R	94	GLY	-	linker	UNP P78423
R	95	SER	-	linker	UNP P78423
R	96	GLY	-	linker	UNP P78423
R	97	SER	-	linker	UNP P78423
R	98	GLY	-	linker	UNP P78423
R	99	SER	-	linker	UNP P78423
R	100	GLY	-	linker	UNP P78423
R	101	SER	-	linker	UNP P78423
R	102	GLY	-	linker	UNP P78423
R	103	SER	-	linker	UNP P78423
R	222	LEU	ILE	engineered mutation	UNP P49238
R	278	CYS	LEU	engineered mutation	UNP P49238
R	323	SER	CYS	engineered mutation	UNP P49238

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
R	352	VAL	MET	engineered mutation	UNP P49238
R	418	GLU	-	expression tag	UNP P49238
R	419	PHE	-	expression tag	UNP P49238
R	420	LEU	-	expression tag	UNP P49238
R	421	GLU	-	expression tag	UNP P49238
R	422	VAL	-	expression tag	UNP P49238
R	423	LEU	-	expression tag	UNP P49238
R	424	PHE	-	expression tag	UNP P49238
R	425	GLN	-	expression tag	UNP P49238
R	426	GLY	-	expression tag	UNP P49238
R	427	PRO	-	expression tag	UNP P49238
R	428	TRP	-	expression tag	UNP P49238
R	429	SER	-	expression tag	UNP P49238
R	430	HIS	-	expression tag	UNP P49238
R	431	PRO	-	expression tag	UNP P49238
R	432	GLN	-	expression tag	UNP P49238
R	433	PHE	-	expression tag	UNP P49238
R	434	GLU	-	expression tag	UNP P49238
R	435	LYS	-	expression tag	UNP P49238
R	436	GLY	-	expression tag	UNP P49238
R	437	GLY	-	expression tag	UNP P49238
R	438	GLY	-	expression tag	UNP P49238
R	439	SER	-	expression tag	UNP P49238
R	440	GLY	-	expression tag	UNP P49238
R	441	GLY	-	expression tag	UNP P49238
R	442	GLY	-	expression tag	UNP P49238
R	443	SER	-	expression tag	UNP P49238
R	444	GLY	-	expression tag	UNP P49238
R	445	GLY	-	expression tag	UNP P49238
R	446	SER	-	expression tag	UNP P49238
R	447	ALA	-	expression tag	UNP P49238
R	448	TRP	-	expression tag	UNP P49238
R	449	SER	-	expression tag	UNP P49238
R	450	HIS	-	expression tag	UNP P49238
R	451	PRO	-	expression tag	UNP P49238
R	452	GLN	-	expression tag	UNP P49238
R	453	PHE	-	expression tag	UNP P49238
R	454	GLU	-	expression tag	UNP P49238
R	455	LYS	-	expression tag	UNP P49238
R	456	ASP	-	expression tag	UNP P49238
R	457	TYR	-	expression tag	UNP P49238
R	458	LYS	-	expression tag	UNP P49238

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	459	ASP	-	expression tag	UNP P49238
R	460	ASP	-	expression tag	UNP P49238
R	461	ASP	-	expression tag	UNP P49238
R	462	ASP	-	expression tag	UNP P49238
R	463	LYS	-	expression tag	UNP P49238

- Molecule 5 is CHOLESTEROL (three-letter code: CLR) (formula: $C_{27}H_{46}O$).



Mol	Chain	Residues	Atoms			AltConf
5	R	1	Total	C	O	0
			56	54	2	
5	R	1	Total	C	O	0
			56	54	2	

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	490779	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	2.1875	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.142	Depositor
Minimum map value	-0.104	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.018	Depositor
Map size (\AA)	267.52, 267.52, 267.52	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.045, 1.045, 1.045	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.24	0/1649	0.41	0/2235
2	D	0.24	0/2289	0.45	0/3130
3	E	0.24	0/228	0.35	0/317
4	R	0.24	0/2518	0.41	0/3462
All	All	0.24	0/6684	0.42	0/9144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1622	1506	1517	14	0
2	D	2243	2050	2052	16	0
3	E	222	192	192	0	0
4	R	2473	2280	2281	47	0
5	R	56	0	90	6	0
All	All	6616	6028	6132	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:R:223:THR:HG21	5:R:501:CLR:H262	1.57	0.87
4:R:230:TYR:O	4:R:234:VAL:HG12	1.85	0.76
4:R:213:ILE:HD12	4:R:260:ILE:HD13	1.69	0.72
4:R:278:CYS:SG	4:R:279:GLY:N	2.65	0.69
4:R:60:ASP:OD1	4:R:61:ALA:N	2.26	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	210/354 (59%)	206 (98%)	4 (2%)	0	100	100
2	D	309/346 (89%)	303 (98%)	6 (2%)	0	100	100
3	E	31/71 (44%)	31 (100%)	0	0	100	100
4	R	343/463 (74%)	333 (97%)	10 (3%)	0	100	100
All	All	893/1234 (72%)	873 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	164/306 (54%)	163 (99%)	1 (1%)	86	94
2	D	229/289 (79%)	228 (100%)	1 (0%)	91	95
3	E	20/58 (34%)	20 (100%)	0	100	100
4	R	228/393 (58%)	228 (100%)	0	100	100
All	All	641/1046 (61%)	639 (100%)	2 (0%)	92	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	257	LYS
2	D	230	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	R	294	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PCA	R	1	4	7,8,9	2.21	2 (28%)	9,10,12	2.06	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PCA	R	1	4	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	R	1	PCA	CD-N	4.67	1.46	1.34
4	R	1	PCA	CA-N	3.35	1.50	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	R	1	PCA	OE-CD-CG	-3.09	121.37	126.76
4	R	1	PCA	CA-N-CD	-2.99	103.35	113.58
4	R	1	PCA	CB-CA-N	2.49	110.45	103.30
4	R	1	PCA	CG-CD-N	2.46	114.76	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	CLR	R	501	-	31,31,31	1.76	5 (16%)	48,48,48	1.45	8 (16%)
5	CLR	R	502	-	31,31,31	1.76	5 (16%)	48,48,48	1.36	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CLR	R	501	-	-	5/10/68/68	0/4/4/4
5	CLR	R	502	-	-	3/10/68/68	0/4/4/4

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	R	501	CLR	C6-C5	6.24	1.46	1.33
5	R	502	CLR	C6-C5	6.24	1.46	1.33
5	R	502	CLR	O1-C3	-4.33	1.30	1.43
5	R	501	CLR	O1-C3	-4.33	1.30	1.43
5	R	502	CLR	C4-C3	3.35	1.58	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	R	501	CLR	C4-C5-C10	3.37	120.90	116.42
5	R	501	CLR	C12-C11-C9	3.31	118.85	113.11
5	R	502	CLR	C4-C5-C10	3.16	120.62	116.42
5	R	502	CLR	C12-C11-C9	2.97	118.27	113.11
5	R	501	CLR	C2-C3-C4	-2.88	106.36	110.31

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	R	501	CLR	C21-C20-C22-C23
5	R	501	CLR	C22-C23-C24-C25
5	R	501	CLR	C20-C22-C23-C24
5	R	501	CLR	C23-C24-C25-C26
5	R	501	CLR	C23-C24-C25-C27

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	R	501	CLR	4	0
5	R	502	CLR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33108. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 101




Y Index: 144




Z Index: 126


6.3.2 Raw map



X Index: 132



Y Index: 153




Z Index: 126

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.018. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 47 nm³; this corresponds to an approximate mass of 42 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.93	5.98	4.02

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.93 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33108 and PDB model 7XBX. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.018 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.018).










9.4 Atom inclusion [i](#)



At the recommended contour level, 65% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.018) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6524	 0.3610
C	 0.6982	 0.4080
D	 0.7252	 0.3830
E	 0.6290	 0.3100
R	 0.5420	 0.3150

