



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 5, 2024 – 08:59 AM EST

PDB ID : 1XDQ  
Title : Structural and Biochemical Identification of a Novel Bacterial Oxidoreductase  
Authors : Loschi, L.; Brokx, S.J.; Hills, T.L.; Zhang, G.; Bertero, M.G.; Lovering, A.L.;  
Weiner, J.H.; Strynadka, N.C.  
Deposited on : 2004-09-07  
Resolution : 2.55 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 % 63% 20% 5% 12%
1	B	298	 4% 61% 23% • 12%
1	C	298	 2% 59% 25% • 11%
1	D	298	 3% 58% 22% 7% 12%
1	E	298	 4% 62% 20% 6% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
5	MTE	A	301	X	-	-	-
5	MTE	B	302	X	-	-	-
5	MTE	C	303	X	-	-	-
5	MTE	D	304	X	-	-	-
5	MTE	E	305	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10771 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacterial Sulfite Oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	Total 2085	C 1351	N 353	O 375	S 6	0	0	0
1	B	263	Total 2093	C 1357	N 354	O 376	S 6	0	0	0
1	C	264	Total 2102	C 1363	N 356	O 377	S 6	0	0	0
1	D	262	Total 2085	C 1351	N 353	O 375	S 6	0	0	0
1	E	262	Total 2085	C 1351	N 353	O 375	S 6	0	0	0

There are 45 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	275	ASP	ALA	SEE REMARK 999	UNP P76342
A	291	LEU	-	expression tag	UNP P76342
A	292	GLU	-	expression tag	UNP P76342
A	293	HIS	-	expression tag	UNP P76342
A	294	HIS	-	expression tag	UNP P76342
A	295	HIS	-	expression tag	UNP P76342
A	296	HIS	-	expression tag	UNP P76342
A	297	HIS	-	expression tag	UNP P76342
A	298	HIS	-	expression tag	UNP P76342
B	275	ASP	ALA	SEE REMARK 999	UNP P76342
B	291	LEU	-	expression tag	UNP P76342
B	292	GLU	-	expression tag	UNP P76342
B	293	HIS	-	expression tag	UNP P76342
B	294	HIS	-	expression tag	UNP P76342
B	295	HIS	-	expression tag	UNP P76342
B	296	HIS	-	expression tag	UNP P76342
B	297	HIS	-	expression tag	UNP P76342
B	298	HIS	-	expression tag	UNP P76342
C	275	ASP	ALA	SEE REMARK 999	UNP P76342

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	291	LEU	-	expression tag	UNP P76342
C	292	GLU	-	expression tag	UNP P76342
C	293	HIS	-	expression tag	UNP P76342
C	294	HIS	-	expression tag	UNP P76342
C	295	HIS	-	expression tag	UNP P76342
C	296	HIS	-	expression tag	UNP P76342
C	297	HIS	-	expression tag	UNP P76342
C	298	HIS	-	expression tag	UNP P76342
D	275	ASP	ALA	SEE REMARK 999	UNP P76342
D	291	LEU	-	expression tag	UNP P76342
D	292	GLU	-	expression tag	UNP P76342
D	293	HIS	-	expression tag	UNP P76342
D	294	HIS	-	expression tag	UNP P76342
D	295	HIS	-	expression tag	UNP P76342
D	296	HIS	-	expression tag	UNP P76342
D	297	HIS	-	expression tag	UNP P76342
D	298	HIS	-	expression tag	UNP P76342
E	275	ASP	ALA	SEE REMARK 999	UNP P76342
E	291	LEU	-	expression tag	UNP P76342
E	292	GLU	-	expression tag	UNP P76342
E	293	HIS	-	expression tag	UNP P76342
E	294	HIS	-	expression tag	UNP P76342
E	295	HIS	-	expression tag	UNP P76342
E	296	HIS	-	expression tag	UNP P76342
E	297	HIS	-	expression tag	UNP P76342
E	298	HIS	-	expression tag	UNP P76342

- Molecule 2 is UREA (three-letter code: URE) (formula: CH<sub>4</sub>N<sub>2</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			4	1	2	1		
2	B	1	Total	C	N	O	0	0
			4	1	2	1		
2	C	1	Total	C	N	O	0	0
			4	1	2	1		
2	D	1	Total	C	N	O	0	0
			4	1	2	1		
2	E	1	Total	C	N	O	0	0
			4	1	2	1		


- Molecule 3 is MOLYBDENUM ATOM (three-letter code: MO) (formula: Mo).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mo	0	0
			1	1		
3	B	1	Total	Mo	0	0
			1	1		
3	C	1	Total	Mo	0	0
			1	1		
3	D	1	Total	Mo	0	0
			1	1		
3	E	1	Total	Mo	0	0
			1	1		

- Molecule 4 is OXYGEN ATOM (three-letter code: O) (formula: O).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 1 1	0	0
4	B	1	Total O 1 1	0	0
4	C	1	Total O 1 1	0	0
4	D	1	Total O 1 1	0	0
4	E	1	Total O 1 1	0	0

- Molecule 5 is PHOSPHONIC ACIDMONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,7,8A, 9,10,10A-HEXAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: MTE) (formula:  $C_{10}H_{14}N_5O_6PS_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O P S 24 10 5 6 1 2	0	0
5	B	1	Total C N O P S 24 10 5 6 1 2	0	0
5	C	1	Total C N O P S 24 10 5 6 1 2	0	0
5	D	1	Total C N O P S 24 10 5 6 1 2	0	0
5	E	1	Total C N O P S 24 10 5 6 1 2	0	0

- Molecule 6 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	49	Total 49	O 49	0	0
6	B	26	Total 26	O 26	0	0
6	C	28	Total 28	O 28	0	0
6	D	36	Total 36	O 36	0	0
6	E	32	Total 32	O 32	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacterial Sulfite Oxidase



- Molecule 1: Bacterial Sulfite Oxidase



- Molecule 1: Bacterial Sulfite Oxidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.34Å 165.06Å 181.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.75 – 2.55 39.75 – 2.54	Depositor EDS
% Data completeness (in resolution range)	95.3 (39.75-2.55) 94.6 (39.75-2.54)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.16 (at 2.54Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.230 , 0.274 0.222 , 0.265	Depositor DCC
$R_{free}$ test set	3654 reflections (7.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.2	Xtrriage
Anisotropy	0.565	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10771	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O, URE, MTE, MO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2150	0.80	2/2932 (0.1%)
1	B	0.50	0/2158	0.80	2/2943 (0.1%)
1	C	0.51	0/2167	0.81	5/2954 (0.2%)
1	D	0.51	0/2150	0.80	3/2932 (0.1%)
1	E	0.49	0/2150	0.76	1/2932 (0.0%)
All	All	0.51	0/10775	0.80	13/14693 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	96	ARG	NE-CZ-NH2	7.48	124.04	120.30
1	D	101	ARG	NE-CZ-NH1	-6.35	117.13	120.30
1	D	96	ARG	NE-CZ-NH1	-5.80	117.40	120.30
1	A	101	ARG	NE-CZ-NH1	-5.57	117.51	120.30
1	C	101	ARG	NE-CZ-NH1	-5.38	117.61	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	158	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2085	0	2051	57	0
1	B	2093	0	2062	65	0
1	C	2102	0	2075	72	0
1	D	2085	0	2051	75	0
1	E	2085	0	2051	59	0
2	A	4	0	4	0	0
2	B	4	0	4	0	0
2	C	4	0	4	0	0
2	D	4	0	4	0	0
2	E	4	0	4	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
5	A	24	0	9	1	0
5	B	24	0	9	3	0
5	C	24	0	9	3	0
5	D	24	0	9	2	0
5	E	24	0	9	2	0
6	A	49	0	0	2	1
6	B	26	0	0	0	0
6	C	28	0	0	1	0
6	D	36	0	0	4	0
6	E	32	0	0	2	0
All	All	10771	0	10355	326	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ALA:HA	1:A:188:GLN:HG2	1.27	1.14
1:D:57:ALA:HA	1:D:188:GLN:HG2	1.30	1.11
1:B:57:ALA:HA	1:B:188:GLN:HG2	1.29	1.07
1:E:57:ALA:HA	1:E:188:GLN:HG2	1.35	1.06
1:C:57:ALA:HA	1:C:188:GLN:HG2	1.38	1.05

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:646:HOH:O	6:A:646:HOH:O[6_445]	2.10	0.10

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/298 (87%)	245 (95%)	12 (5%)	1 (0%)	34	46
1	B	259/298 (87%)	240 (93%)	18 (7%)	1 (0%)	34	46
1	C	260/298 (87%)	244 (94%)	14 (5%)	2 (1%)	19	27
1	D	258/298 (87%)	242 (94%)	14 (5%)	2 (1%)	19	27
1	E	258/298 (87%)	241 (93%)	15 (6%)	2 (1%)	19	27
All	All	1293/1490 (87%)	1212 (94%)	73 (6%)	8 (1%)	25	34

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	143	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	143	GLN
1	C	143	GLN
1	D	143	GLN
1	E	143	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/249 (88%)	197 (90%)	21 (10%)	8	9
1	B	219/249 (88%)	200 (91%)	19 (9%)	10	12
1	C	220/249 (88%)	200 (91%)	20 (9%)	9	11
1	D	218/249 (88%)	195 (89%)	23 (11%)	6	7
1	E	218/249 (88%)	196 (90%)	22 (10%)	7	8
All	All	1093/1245 (88%)	988 (90%)	105 (10%)	8	9

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	264	ARG
1	D	175	LEU
1	E	214	LEU
1	C	286	LEU
1	D	88	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	148	GLN
1	D	148	GLN
1	E	224	ASN
1	E	148	GLN
1	B	148	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 10 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	URE	B	402	-	3,3,3	0.86	0	3,3,3	1.95	1 (33%)
2	URE	A	401	-	3,3,3	0.97	0	3,3,3	2.03	2 (66%)
5	MTE	B	302	3	21,26,26	4.19	14 (66%)	21,40,40	6.18	13 (61%)
5	MTE	D	304	3	21,26,26	4.71	15 (71%)	21,40,40	5.96	12 (57%)
5	MTE	E	305	3	21,26,26	4.76	14 (66%)	21,40,40	6.17	14 (66%)
2	URE	C	403	-	3,3,3	0.81	0	3,3,3	2.01	2 (66%)
5	MTE	C	303	3	21,26,26	4.82	15 (71%)	21,40,40	5.89	11 (52%)
5	MTE	A	301	3	21,26,26	4.60	15 (71%)	21,40,40	6.22	12 (57%)
2	URE	D	404	-	3,3,3	1.14	0	3,3,3	1.95	2 (66%)
2	URE	E	405	-	3,3,3	0.85	0	3,3,3	2.01	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MTE	D	304	3	1/1/6/8	2/6/34/34	0/3/3/3
5	MTE	E	305	3	1/1/6/8	2/6/34/34	0/3/3/3
5	MTE	B	302	3	1/1/6/8	2/6/34/34	0/3/3/3
5	MTE	A	301	3	1/1/6/8	2/6/34/34	0/3/3/3
5	MTE	C	303	3	1/1/6/8	2/6/34/34	0/3/3/3

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	304	MTE	C7-C6	13.10	1.64	1.53
5	C	303	MTE	C7-C6	11.55	1.62	1.53
5	E	305	MTE	C7-C6	10.72	1.62	1.53
5	A	301	MTE	C7-C6	10.64	1.62	1.53
5	C	303	MTE	C4'-C3'	-9.72	1.38	1.52

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	301	MTE	O3'-C7-C6	-24.52	92.61	108.96
5	B	302	MTE	O3'-C7-C6	-24.43	92.67	108.96
5	E	305	MTE	O3'-C7-C6	-24.04	92.93	108.96
5	D	304	MTE	O3'-C7-C6	-23.34	93.39	108.96
5	C	303	MTE	O3'-C7-C6	-22.87	93.71	108.96

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	301	MTE	C6
5	B	302	MTE	C6
5	C	303	MTE	C6
5	D	304	MTE	C6
5	E	305	MTE	C6

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	301	MTE	C4'-O4'-P-O2P
5	A	301	MTE	C4'-O4'-P-O3P
5	B	302	MTE	C4'-O4'-P-O2P
5	B	302	MTE	C4'-O4'-P-O3P
5	C	303	MTE	C4'-O4'-P-O2P



There are no ring outliers.


5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	302	MTE	3	0
5	D	304	MTE	2	0
5	E	305	MTE	2	0
5	C	303	MTE	3	0
5	A	301	MTE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/298 (87%)	-0.12	2 (0%) 86 89	29, 50, 74, 105	0
1	B	263/298 (88%)	0.07	12 (4%) 32 39	34, 55, 80, 99	0
1	C	264/298 (88%)	0.13	6 (2%) 60 67	36, 56, 80, 102	0
1	D	262/298 (87%)	0.15	8 (3%) 49 56	36, 56, 81, 98	0
1	E	262/298 (87%)	0.19	11 (4%) 36 42	38, 58, 82, 101	0
All	All	1313/1490 (88%)	0.08	39 (2%) 50 57	29, 55, 81, 105	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	31	ASN	3.6
1	B	31	ASN	3.6
1	D	89	ARG	3.5
1	A	150	ARG	3.2
1	E	27	ALA	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	URE	A	401	4/4	0.74	0.26	49,62,62,66	0
2	URE	E	405	4/4	0.74	0.39	51,53,61,63	0
2	URE	D	404	4/4	0.83	0.44	40,43,53,55	0
2	URE	B	402	4/4	0.83	0.21	52,58,63,63	0
2	URE	C	403	4/4	0.84	0.34	49,56,62,64	0
5	MTE	B	302	24/24	0.96	0.16	26,43,50,55	0
5	MTE	C	303	24/24	0.97	0.18	41,45,51,54	0
5	MTE	D	304	24/24	0.97	0.18	37,47,52,54	0
5	MTE	E	305	24/24	0.97	0.19	42,48,54,57	0
5	MTE	A	301	24/24	0.98	0.15	34,41,44,51	0
3	MO	E	505	1/1	0.99	0.14	49,49,49,49	0
3	MO	D	504	1/1	0.99	0.15	41,41,41,41	0
4	O	C	603	1/1	1.00	0.14	42,42,42,42	0
4	O	D	604	1/1	1.00	0.16	31,31,31,31	0
4	O	E	605	1/1	1.00	0.15	32,32,32,32	0
3	MO	C	503	1/1	1.00	0.14	41,41,41,41	0
3	MO	A	501	1/1	1.00	0.13	40,40,40,40	0
3	MO	B	502	1/1	1.00	0.13	41,41,41,41	0
4	O	A	601	1/1	1.00	0.12	35,35,35,35	0
4	O	B	602	1/1	1.00	0.14	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around MTE B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MTE C 303:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around MTE D 304:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.