



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 07:29 PM BST

PDB ID : 4XGO  
Title : Crystal structure of leucine-rich repeat domain of APL1B  
Authors : Williams, M.; Summers, B.; Baxter, R.H.G.  
Deposited on : 2015-01-01  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	 5% 91% . .
1	B	356	 10% 88% . 8%
2	C	2	 100%
2	D	2	 100%

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 6299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anopheles Plasmodium-responsive Leucine-rich repeat protein 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	2746	1762	460	511	13	0	7	0
1	B	328	2654	1698	448	493	15	0	9	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	MET	LEU	conflict	UNP F2YBL9
A	176	ALA	VAL	conflict	UNP F2YBL9
A	225	VAL	THR	conflict	UNP F2YBL9
A	236	LYS	ILE	conflict	UNP F2YBL9
A	239	TYR	ILE	conflict	UNP F2YBL9
A	240	GLN	HIS	conflict	UNP F2YBL9
A	244	LYS	THR	conflict	UNP F2YBL9
A	245	MET	ILE	conflict	UNP F2YBL9
A	371	HIS	-	expression tag	UNP F2YBL9
A	372	HIS	-	expression tag	UNP F2YBL9
A	373	HIS	-	expression tag	UNP F2YBL9
A	374	HIS	-	expression tag	UNP F2YBL9
A	375	HIS	-	expression tag	UNP F2YBL9
A	376	HIS	-	expression tag	UNP F2YBL9
B	120	MET	LEU	conflict	UNP F2YBL9
B	176	ALA	VAL	conflict	UNP F2YBL9
B	225	VAL	THR	conflict	UNP F2YBL9
B	236	LYS	ILE	conflict	UNP F2YBL9
B	239	TYR	ILE	conflict	UNP F2YBL9
B	240	GLN	HIS	conflict	UNP F2YBL9
B	244	LYS	THR	conflict	UNP F2YBL9
B	245	MET	ILE	conflict	UNP F2YBL9
B	371	HIS	-	expression tag	UNP F2YBL9
B	372	HIS	-	expression tag	UNP F2YBL9

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	373	HIS	-	expression tag	UNP F2YBL9
B	374	HIS	-	expression tag	UNP F2YBL9
B	375	HIS	-	expression tag	UNP F2YBL9
B	376	HIS	-	expression tag	UNP F2YBL9

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	28	16	2	10	0	0	0
2	D	2	28	16	2	10	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
3	A	1	5	4	1	0	0
3	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
3	B	1	5	4	1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0
4	B	1	4	2	2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Cl 1 1	0	0
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Na 1 1	0	0


- Molecule 8 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
8	A	432	Total 438	O 438	0	6
8	B	340	Total 343	O 343	0	5


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Anopheles Plasmodium-responsive Leucine-rich repeat protein 1B




- Molecule 1: Anopheles Plasmodium-responsive Leucine-rich repeat protein 1B



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.42Å 74.96Å 214.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	107.18 – 1.74 40.33 – 1.74	Depositor EDS
% Data completeness (in resolution range)	89.9 (107.18-1.74) 90.0 (40.33-1.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 1.74Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.187 , 0.220 0.193 , 0.225	Depositor DCC
$R_{free}$ test set	4796 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.9	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 52.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6299	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, CL, NA, PO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2830	0.63	0/3861
1	B	0.37	0/2728	0.59	0/3719
All	All	0.40	0/5558	0.61	0/7580

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2746	0	2695	10	0
1	B	2654	0	2583	9	0
2	C	28	0	25	0	0
2	D	28	0	25	1	0
3	A	5	0	0	0	0
3	B	10	0	0	0	0
4	A	4	0	3	0	0
4	B	4	0	3	1	0
5	A	36	0	48	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	1	0	0	0	0
8	A	438	0	0	2	0
8	B	343	0	0	0	0
All	All	6299	0	5382	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:THR:HA	1:A:211:ASN:HD22	1.66	0.60
1:B:214:ASP:HA	4:B:1003:ACT:H3	1.83	0.59
1:A:59:ASP:OD1	1:B:29:HIS:NE2	2.40	0.54
1:B:162:LEU:HD13	1:B:165:VAL:HG21	1.91	0.53
1:B:147[B]:SER:OG	2:D:1:NAG:H62	2.15	0.47
1:A:59:ASP:OD2	1:B:31:THR:HG22	2.16	0.45
1:A:27:SER:HA	1:B:54:GLN:HB3	2.00	0.42
1:B:208:GLN:HA	1:B:230:SER:O	2.20	0.42
1:A:264:TYR:O	1:A:287:ASN:ND2	2.54	0.41
1:A:99:ASP:OD1	1:A:121:SER:OG	2.32	0.41
1:B:351:ILE:O	1:B:352:ASP:CB	2.68	0.41
1:A:45:VAL:O	1:A:45:VAL:HG12	2.21	0.41
1:A:76:PHE:CE1	1:A:100:ARG:HG3	2.56	0.41
1:B:56:GLU:OE2	1:B:80:ARG:NE	2.54	0.41
1:A:317:ASN:ND2	8:A:1517:HOH:O	2.32	0.40
5:A:1010:GOL:O1	8:A:1530:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/356 (97%)	319 (92%)	26 (8%)	0	100	100
1	B	331/356 (93%)	306 (92%)	25 (8%)	0	100	100
All	All	676/712 (95%)	625 (92%)	51 (8%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/331 (94%)	308 (99%)	3 (1%)	76	63
1	B	298/331 (90%)	297 (100%)	1 (0%)	92	89
All	All	609/662 (92%)	605 (99%)	4 (1%)	84	75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	TYR
1	A	214	ASP
1	A	247	ARG
1	B	136	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	287	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	D	1	1,2	14,14,15	0.67	0	17,19,21	0.82	0
2	NAG	D	2	2	14,14,15	0.52	0	17,19,21	0.83	1 (5%)
2	NAG	C	2	2	14,14,15	0.52	0	17,19,21	1.09	1 (5%)
2	NAG	C	1	1,2	14,14,15	0.66	0	17,19,21	1.16	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	O5-C1-C2	-2.64	107.13	111.29
2	C	2	NAG	C3-C4-C5	2.26	114.27	110.24
2	D	2	NAG	C1-O5-C5	2.21	115.19	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O5-C5-C6-O6

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	1,2	14,14,15	0.66	0	17,19,21	1.16	1 (5%)
2	NAG	C	2	2	14,14,15	0.52	0	17,19,21	1.09	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.67	0	17,19,21	0.82	0
2	NAG	D	2	2	14,14,15	0.52	0	17,19,21	0.83	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	O5-C1-C2	-2.64	107.13	111.29
2	C	2	NAG	C3-C4-C5	2.26	114.27	110.24
2	D	2	NAG	C1-O5-C5	2.21	115.19	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	1006	-	5,5,5	0.48	0	5,5,5	0.33	0
5	GOL	A	1009	-	5,5,5	0.38	0	5,5,5	0.18	0
5	GOL	A	1007	-	5,5,5	0.29	0	5,5,5	0.39	0
5	GOL	A	1010	-	5,5,5	0.32	0	5,5,5	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	1003	-	4,4,4	0.94	0	6,6,6	0.70	0
3	PO4	B	1004	-	4,4,4	0.85	0	6,6,6	0.47	0
3	PO4	B	1005	-	4,4,4	0.85	0	6,6,6	0.40	0
4	ACT	B	1003	-	1,3,3	0.55	0	0,3,3	0.00	-
5	GOL	A	1005	-	5,5,5	0.49	0	5,5,5	0.22	0
5	GOL	A	1008	-	5,5,5	0.34	0	5,5,5	0.31	0
4	ACT	A	1004	-	1,3,3	0.16	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	1006	-	-	2/4/4/4	-
5	GOL	A	1009	-	-	4/4/4/4	-
5	GOL	A	1007	-	-	2/4/4/4	-
5	GOL	A	1010	-	-	2/4/4/4	-
5	GOL	A	1005	-	-	0/4/4/4	-
5	GOL	A	1008	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1006	GOL	O1-C1-C2-O2
5	A	1006	GOL	O1-C1-C2-C3
5	A	1009	GOL	O1-C1-C2-C3
5	A	1009	GOL	C1-C2-C3-O3
5	A	1009	GOL	O2-C2-C3-O3
5	A	1007	GOL	C1-C2-C3-O3
5	A	1010	GOL	O1-C1-C2-C3
5	A	1008	GOL	O1-C1-C2-C3
5	A	1010	GOL	O1-C1-C2-O2
5	A	1008	GOL	C1-C2-C3-O3
5	A	1009	GOL	O1-C1-C2-O2
5	A	1008	GOL	O2-C2-C3-O3
5	A	1008	GOL	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	A	1007	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1010	GOL	1	0
4	B	1003	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	340/356 (95%)	0.16	19 (5%) 24 29	9, 20, 41, 57	0
1	B	328/356 (92%)	0.63	37 (11%) 5 6	11, 27, 70, 93	0
All	All	668/712 (93%)	0.39	56 (8%) 11 14	9, 23, 60, 93	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	311	THR	9.9
1	B	337	VAL	7.9
1	B	336	ASN	6.4
1	B	313	HIS	6.3
1	B	315	LEU	6.3
1	B	312	SER	5.6
1	B	314	THR	5.3
1	A	25	TYR	5.2
1	B	343	HIS	4.7
1	B	334	PHE	4.7
1	B	270	THR	4.5
1	B	296	TYR	4.2
1	A	311	THR	4.2
1	B	287	ASN	4.2
1	B	355	LEU	3.9
1	A	336	ASN	3.8
1	B	351	ILE	3.7
1	B	342	VAL	3.7
1	A	314	THR	3.4
1	A	65	TYR	3.3
1	B	357	HIS	3.3
1	A	335	ARG	3.1
1	B	292	ASP	2.9
1	A	338	ALA	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	246	GLN	2.9
1	B	309	LEU	2.8
1	B	335	ARG	2.7
1	A	313	HIS	2.7
1	B	354	HIS	2.7
1	A	341	ALA	2.5
1	A	337	VAL	2.5
1	B	295	GLN	2.5
1	B	333	LEU	2.4
1	B	348	HIS	2.4
1	B	353	TYR	2.4
1	B	33	PHE	2.4
1	A	343	HIS	2.4
1	A	340	PRO	2.3
1	B	288	GLN	2.3
1	B	240	GLN	2.3
1	B	291	PHE	2.3
1	A	312	SER	2.2
1	B	318	LEU	2.2
1	A	30	LEU	2.2
1	B	67	HIS	2.2
1	B	298	TYR	2.2
1	B	360	CYS	2.2
1	A	339	GLN	2.2
1	B	356	GLU	2.2
1	B	290	GLN	2.1
1	B	364	SER	2.1
1	A	267	PRO	2.1
1	A	292	ASP	2.1
1	B	29	HIS	2.0
1	B	347	GLN	2.0
1	A	334	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.80	0.23	42,49,57,58	0

*Continued on next page...*

Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	D	2	14/15	0.85	0.17	41,47,51,55	0
2	NAG	C	1	14/15	0.90	0.12	17,21,33,37	0
2	NAG	D	1	14/15	0.95	0.09	19,25,28,32	0


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.80	0.23	42,49,57,58	0
2	NAG	D	2	14/15	0.85	0.17	41,47,51,55	0
2	NAG	C	1	14/15	0.90	0.12	17,21,33,37	0
2	NAG	D	1	14/15	0.95	0.09	19,25,28,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	1008	6/6	0.60	0.14	53,55,56,56	0
5	GOL	A	1009	6/6	0.74	0.19	50,50,51,52	0
5	GOL	A	1006	6/6	0.76	0.18	38,41,42,44	0
4	ACT	B	1003	4/4	0.80	0.14	41,41,41,44	0
4	ACT	A	1004	4/4	0.80	0.28	42,44,45,46	0
5	GOL	A	1007	6/6	0.82	0.13	35,42,45,46	0
3	PO4	B	1004	5/5	0.82	0.15	69,71,73,75	0
6	CL	A	1011	1/1	0.83	0.11	54,54,54,54	0
5	GOL	A	1010	6/6	0.84	0.22	36,39,43,47	0
7	NA	B	1007	1/1	0.89	0.17	27,27,27,27	0
5	GOL	A	1005	6/6	0.90	0.11	30,34,34,34	0
3	PO4	B	1005	5/5	0.90	0.29	57,59,62,63	0
6	CL	B	1006	1/1	0.96	0.17	64,64,64,64	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	A	1003	5/5	0.98	0.06	23,23,23,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.