

# wwPDB X-ray Structure Validation Summary Report (i)

Nov 13, 2023 – 02:33 PM JST

PDB ID : 5XG9

Title : Crystal Structure of PEG-bound SH3 domain of Myosin IB from Entamoeba

histolytica

Authors: Gautam, G.; Gourinath, S.

Deposited on : 2017-04-13

Resolution : 1.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	66	83%	5% •	11%
1	В	66	77%	11% •	11%
1	С	66	83%	6%	11%
1	D	66	85%	5%	11%
1	Е	66	83%	6%	11%
1	F	66	80%	8%	12%





 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain					
1	G	66	86%	11%				
1	Н	66	86%	11%				



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4519 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Unconventional myosin IB.

Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf	Trace
1	В	59	Total	С	N	О	S	0	0	0
1	Б	39	472	306	73	92	1	0	U	U
1	A	59	Total	С	N	О	S	0	0	0
1	A	39	475	309	73	92	1	0	U	U
1	С	59	Total	С	N	О	S	0	0	0
1		39	472	306	73	92	1	0	U	U
1	D	59	Total C N	N	О	S	0	1	0	
1	D	39	481	311	74	95	1	0	1	
1	Е	59	Total	С	N	О	S	0	0	0
1	15	39	475	309	73	92	1	0	0	U
1	F	58	Total	С	N	Ο	S	0	0	0
1	I.	30	467	303	72	91	1	0	0	U
1	G	59	Total	С	N	О	S	0	0	0
1	G	99	472	306	73	92	1		U	U
1	Н	59	Total	С	N	О	S	0	0	0
1	11	J9	472	306	73	92	1		0	

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-1	ALA	-	expression tag	UNP C4LUC7
В	0	SER	-	expression tag	UNP C4LUC7
В	1	MET	-	expression tag	UNP C4LUC7
В	57	LEU	-	expression tag	UNP C4LUC7
В	58	GLU	-	expression tag	UNP C4LUC7
В	59	HIS	-	expression tag	UNP C4LUC7
В	60	HIS	-	expression tag	UNP C4LUC7
В	61	HIS	-	expression tag	UNP C4LUC7
В	62	HIS	-	expression tag	UNP C4LUC7
В	63	HIS	-	expression tag	UNP C4LUC7
В	64	HIS	-	expression tag	UNP C4LUC7
A	-1	ALA	-	expression tag	UNP C4LUC7
A	0	SER	-	expression tag	UNP C4LUC7

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP C4LUC7
A	57	LEU	-	expression tag	UNP C4LUC7
A	58	GLU	-	expression tag	UNP C4LUC7
A	59	HIS	-	expression tag	UNP C4LUC7
A	60	HIS	-	expression tag	UNP C4LUC7
A	61	HIS	-	expression tag	UNP C4LUC7
A	62	HIS	-	expression tag	UNP C4LUC7
A	63	HIS	-	expression tag	UNP C4LUC7
A	64	HIS	-	expression tag	UNP C4LUC7
С	-1	ALA	-	expression tag	UNP C4LUC7
С	0	SER	-	expression tag	UNP C4LUC7
С	1	MET	-	expression tag	UNP C4LUC7
С	57	LEU	-	expression tag	UNP C4LUC7
С	58	GLU	-	expression tag	UNP C4LUC7
С	59	HIS	-	expression tag	UNP C4LUC7
С	60	HIS	-	expression tag	UNP C4LUC7
С	61	HIS	-	expression tag	UNP C4LUC7
С	62	HIS	-	expression tag	UNP C4LUC7
С	63	HIS	-	expression tag	UNP C4LUC7
С	64	HIS	-	expression tag	UNP C4LUC7
D	-1	ALA	-	expression tag	UNP C4LUC7
D	0	SER	_	expression tag	UNP C4LUC7
D	1	MET	-	expression tag	UNP C4LUC7
D	57	LEU	-	expression tag	UNP C4LUC7
D	58	GLU	_	expression tag	UNP C4LUC7
D	59	HIS	-	expression tag	UNP C4LUC7
D	60	HIS	-	expression tag	UNP C4LUC7
D	61	HIS	-	expression tag	UNP C4LUC7
D	62	HIS	-	expression tag	UNP C4LUC7
D	63	HIS	-	expression tag	UNP C4LUC7
D	64	HIS	-	expression tag	UNP C4LUC7
Е	-1	ALA	-	expression tag	UNP C4LUC7
Е	0	SER	-	expression tag	UNP C4LUC7
Е	1	MET	-	expression tag	UNP C4LUC7
Е	57	LEU	-	expression tag	UNP C4LUC7
Е	58	GLU	-	expression tag	UNP C4LUC7
Е	59	HIS	-	expression tag	UNP C4LUC7
Е	60	HIS	-	expression tag	UNP C4LUC7
E	61	HIS	-	expression tag	UNP C4LUC7
Е	62	HIS	-	expression tag	UNP C4LUC7
Е	63	HIS	-	expression tag	UNP C4LUC7
E	64	HIS	_	expression tag	UNP C4LUC7

Continued on next page...




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	-1	ALA	-	expression tag	UNP C4LUC7
F	0	SER	_	expression tag	UNP C4LUC7
F	1	MET	-	expression tag	UNP C4LUC7
F	57	LEU	_	expression tag	UNP C4LUC7
F	58	GLU	-	expression tag	UNP C4LUC7
F	59	HIS	-	expression tag	UNP C4LUC7
F	60	HIS	-	expression tag	UNP C4LUC7
F	61	HIS	-	expression tag	UNP C4LUC7
F	62	HIS	-	expression tag	UNP C4LUC7
F	63	HIS	-	expression tag	UNP C4LUC7
F	64	HIS	-	expression tag	UNP C4LUC7
G	-1	ALA	-	expression tag	UNP C4LUC7
G	0	SER	-	expression tag	UNP C4LUC7
G	1	MET	-	expression tag	UNP C4LUC7
G	57	LEU	-	expression tag	UNP C4LUC7
G	58	GLU	-	expression tag	UNP C4LUC7
G	59	HIS	-	expression tag	UNP C4LUC7
G	60	HIS	-	expression tag	UNP C4LUC7
G	61	HIS	-	expression tag	UNP C4LUC7
G	62	HIS	-	expression tag	UNP C4LUC7
G	63	HIS	-	expression tag	UNP C4LUC7
G	64	HIS	-	expression tag	UNP C4LUC7
Н	-1	ALA	-	expression tag	UNP C4LUC7
Н	0	SER	-	expression tag	UNP C4LUC7
Н	1	MET	-	expression tag	UNP C4LUC7
Н	57	LEU	-	expression tag	UNP C4LUC7
Н	58	GLU	-	expression tag	UNP C4LUC7
Н	59	HIS	-	expression tag	UNP C4LUC7
Н	60	HIS	-	expression tag	UNP C4LUC7
Н	61	HIS	-	expression tag	UNP C4LUC7
Н	62	HIS	-	expression tag	UNP C4LUC7
Н	63	HIS	-	expression tag	UNP C4LUC7
Н	64	HIS	-	expression tag	UNP C4LUC7


 $\bullet$  Molecule 2 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $\mathrm{C_{10}H_{22}O_6}).$ 





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	В	1	Total 0	C 8	O 5	0	0


• Molecule 3 is  $2,5,8,11,14,17,20,23,26,29,32,35,38,41,44,47,50,53,56,59,62,65,68,71,74,77,80-HEPTACOSAOXADOOCTACONTAN-82-OL (three-letter code: PEU) (formula: <math>C_{55}H_{112}O_{28}$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	В	1	Total C O	0	0	
	D	1	24 16 8		Ü	
3	Λ	1	Total C O	0	0	
9	Λ	1	21 14 7		U	




• Molecule 4 is 1-(2-METHOXY-ETHOXY)-2-{2-[2-(2-METHOXY-ETHOXY]-ETHOXY}-ETHOXY}-ETHOXY}-ETHOXY



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 18 12 6	0	0
4	С	1	Total C O 17 11 6	0	0
4	С	1	Total C O 14 9 5	0	0
4	С	1	Total C O 6 3 3	0	0
4	D	1	Total C O 7 4 3	0	0
4	G	1	Total C O 18 12 6	0	0

 $\bullet$  Molecule 5 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	С	1	Total O S 5 4 1	0	0
5	G	1	Total O S 5 4 1	0	0

# • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	82	Total O 82 82	0	0
6	A	83	Total O 83 83	0	0
6	С	84	Total O 84 84	0	0
6	D	71	Total O 71 71	0	0
6	E	61	Total O 61 61	0	0
6	F	59	Total O 59 59	0	0
6	G	83	Total O 83 83	0	0
6	Н	62	Total O 62 62	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Unconventional myosin IB



• Molecule 1: Unconventional myosin IB







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	106.46Å 79.61Å 88.48Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $122.65^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	74.50 - 1.78	Depositor
rtesolution (A)	31.60 - 1.78	EDS
% Data completeness	98.3 (74.50-1.78)	Depositor
(in resolution range)	98.3 (31.60-1.78)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.52 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
D D.	0.183 , 0.228	Depositor
$R, R_{free}$	0.193 , $0.237$	DCC
$R_{free}$ test set	2892 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.5	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34 , 41.6	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4519	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEU, PG6, SO4, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	Bond lengths		nd angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	1.03	0/487	0.94	1/662 (0.2%)
1	В	1.07	1/484 (0.2%)	0.96	1/658 (0.2%)
1	С	0.99	1/484 (0.2%)	0.92	0/658
1	D	1.00	0/493	0.86	0/670
1	Е	0.94	0/487	0.85	0/662
1	F	0.96	0/479	0.82	0/651
1	G	0.96	0/484	0.94	1/658 (0.2%)
1	Н	0.98	0/484	0.93	1/658 (0.2%)
All	All	0.99	$2/3882 \ (0.1\%)$	0.90	4/5277 (0.1%)

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
1	В	31	GLU	CD-OE2	-5.38	1.19	1.25
1	С	18	GLU	CD-OE1	-5.10	1.20	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	1	MET	CG-SD-CE	6.74	110.99	100.20
1	G	25	ASP	CB-CG-OD1	6.62	124.26	118.30
1	Н	33	ASP	CB-CG-OD2	-6.12	112.79	118.30
1	A	16	ASP	CB-CG-OD2	-5.35	113.48	118.30

There are no chirality outliers.

There are no planarity outliers.



# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	475	0	464	2	0
1	В	472	0	455	8	0
1	С	472	0	455	3	0
1	D	481	0	460	3	1
1	Ε	475	0	464	4	0
1	F	467	0	453	4	0
1	G	472	0	455	0	0
1	Н	472	0	455	1	0
2	В	13	0	17	1	1
3	A	21	0	26	0	0
3	В	24	0	30	3	1
4	A	18	0	26	0	0
4	С	37	0	42	0	0
4	D	7	0	8	0	0
4	G	18	0	26	0	0
5	С	5	0	0	0	0
5	G	5	0	0	0	0
6	A	83	0	0	0	0
6	В	82	0	0	2	0
6	С	84	0	0	3	0
6	D	71	0	0	3	0
6	Ε	61	0	0	3	0
6	F	59	0	0	3	0
6	G	83	0	0	0	1
6	Н	62	0	0	1	0
All	All	4519	0	3836	24	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
2:B:101:1PE:H232	6:B:251:HOH:O	1.85	0.77
1:F:38:LYS:NZ	6:F:101:HOH:O	2.12	0.75

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
1:D:6:LYS:HD3	6:D:218:HOH:O	1.87	0.74
1:B:33:ASP:OD2	3:B:102:PEU:HAM2	1.88	0.73
1:E:33:ASP:HB2	6:E:142:HOH:O	1.92	0.69

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
6:G:283:HOH:O	6:G:283:HOH:O[2_556]	1.33	0.87
2:B:101:1PE:OH6	2:B:101:1PE:OH6[2_555]	2.10	0.10
1:D:33:ASP:OD2	3:B:102:PEU:OAK[4_445]	2.16	0.04

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	57/66~(86%)	55 (96%)	2 (4%)	0	100	100
1	В	57/66~(86%)	54 (95%)	3 (5%)	0	100	100
1	С	57/66~(86%)	54 (95%)	3 (5%)	0	100	100
1	D	58/66~(88%)	57 (98%)	1 (2%)	0	100	100
1	E	57/66~(86%)	56 (98%)	1 (2%)	0	100	100
1	F	56/66~(85%)	54 (96%)	2 (4%)	0	100	100
1	G	57/66~(86%)	57 (100%)	0	0	100	100
1	Н	57/66 (86%)	57 (100%)	0	0	100	100
All	All	456/528~(86%)	444 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	50/57 (88%)	49 (98%)	1 (2%)	55	40
1	В	49/57 (86%)	49 (100%)	0	100	100
1	C	49/57 (86%)	49 (100%)	0	100	100
1	D	50/57 (88%)	50 (100%)	0	100	100
1	${ m E}$	50/57 (88%)	49 (98%)	1 (2%)	55	40
1	F	49/57 (86%)	49 (100%)	0	100	100
1	G	49/57 (86%)	48 (98%)	1 (2%)	55	40
1	Н	49/57 (86%)	49 (100%)	0	100	100
All	All	395/456 (87%)	392 (99%)	3 (1%)	81	76

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASP
1	Е	16	ASP
1	G	0	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	eles
Mol	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PG6	С	102	-	15,15,17	0.68	0	13,13,16	0.59	0
4	PG6	С	104	-	4,4,17	0.42	0	3,3,16	0.55	0
4	PG6	D	101	-	6,6,17	12.74	1 (16%)	5,5,16	1.27	1 (20%)
4	PG6	A	101	-	17,17,17	0.53	0	16,16,16	0.52	0
5	SO4	G	101	-	4,4,4	0.33	0	6,6,6	0.05	0
3	PEU	В	102	_	23,23,82	0.47	0	22,22,81	0.27	0
3	PEU	A	102	_	20,20,82	0.61	0	19,19,81	1.02	0
2	1PE	В	101	-	12,12,15	0.44	0	11,11,14	0.28	0
4	PG6	С	103	_	13,13,17	0.40	0	12,12,16	0.29	0
5	SO4	С	101	-	4,4,4	0.32	0	6,6,6	0.05	0
4	PG6	G	102	-	17,17,17	0.83	0	16,16,16	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG6	С	102	-	-	9/11/11/15	-
4	PG6	С	104	-	-	1/2/2/15	-
4	PG6	D	101	-	-	3/4/4/15	-
4	PG6	A	101	-	-	7/15/15/15	-
3	PEU	В	102	-	-	14/21/21/80	-
3	PEU	A	102	-	-	10/18/18/80	-
2	1PE	В	101	-	-	4/10/10/13	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG6	С	103	-	-	5/11/11/15	-
4	PG6	G	102	-	-	9/15/15/15	-

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
4	D	101	PG6	O4-C7	31.18	3.03	1.42

#### All (1) bond angle outliers are listed below:

$\mathbf{Mol}$	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
4	D	101	PG6	O4-C7-C6	2.77	127.87	111.81

There are no chirality outliers.

5 of 62 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	102	PEU	CAJ-CAI-OAH-CAG
4	G	102	PG6	O3-C6-C7-O4
2	В	101	1PE	OH4-C13-C23-OH3
3	A	102	PEU	OAN-CAO-CAP-OAQ
4	С	102	PG6	O1-C2-C3-O2

There are no ring outliers.


2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	102	PEU	3	1
2	В	101	1PE	1	1


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient



equivalents in the CSD to analyse the geometry.







# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	59/66 (89%)	-0.21	2 (3%) 45 44	17, 23, 38, 47	0
1	В	59/66 (89%)	-0.32	0 100 100	14, 19, 29, 38	0
1	С	59/66 (89%)	-0.27	1 (1%) 70 70	16, 22, 42, 50	0
1	D	59/66 (89%)	-0.40	0 100 100	17, 23, 39, 51	0
1	E	59/66 (89%)	-0.12	1 (1%) 70 70	16, 26, 42, 48	0
1	F	58/66 (87%)	-0.20	1 (1%) 70 70	18, 26, 40, 52	0
1	G	59/66 (89%)	-0.21	1 (1%) 70 70	16, 22, 45, 49	0
1	Н	59/66 (89%)	-0.18	1 (1%) 70 70	19, 26, 38, 50	0
All	All	471/528 (89%)	-0.24	7 (1%) 73 73	14, 24, 41, 52	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	57	LEU	5.4
1	Н	57	LEU	5.4
1	С	57	LEU	4.6
1	A	57	LEU	3.5
1	A	34	GLU	2.8

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	PG6	G	102	18/18	0.78	0.14	45,49,58,59	0
4	PG6	С	102	17/18	0.81	0.12	38,46,53,54	0
4	PG6	A	101	18/18	0.84	0.12	39,46,60,62	0
3	PEU	A	102	21/83	0.86	0.16	33,38,48,48	0
3	PEU	В	102	24/83	0.86	0.14	28,42,55,72	0
4	PG6	С	103	14/18	0.89	0.14	38,46,57,71	0
4	PG6	D	101	7/18	0.90	0.11	42,46,49,57	0
2	1PE	В	101	13/16	0.91	0.12	29,34,41,44	0
5	SO4	G	101	5/5	0.93	0.26	49,58,69,70	0
4	PG6	С	104	6/18	0.94	0.10	28,39,39,40	0
5	SO4	С	101	5/5	0.95	0.17	47,50,54,56	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




# Electron density around PG6 G 102:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around PG6 C 102:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










# Electron density around PG6 A 101: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)









# 









# 6.5 Other polymers (i)

There are no such residues in this entry.

