



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 22, 2023 – 10:47 PM JST

PDB ID : 7XGW  
Title : Apo structure of LW domain from Trypanosoma brucei  
Authors : Liao, S.; Gao, J.; Chen, M.; Tu, X.  
Deposited on : 2022-04-06  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	68% 10% 23%
1	B	133	70% 5% 26%
1	C	133	3% 63% 10% 26%
1	D	133	% 62% 11% 27%
1	E	133	68% 6% 26%
1	F	133	% 68% 9% 23%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 4519 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription elongation factor s-II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	F	103	773	489	130	152	2	0	0	0
1	A	103	771	489	129	151	2	0	0	0
1	B	99	731	466	120	143	2	0	0	0
1	C	98	712	456	118	136	2	0	0	0
1	D	97	723	459	120	142	2	0	0	0
1	E	99	731	464	120	145	2	0	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	227	MET	-	initiating methionine	UNP Q586Y0
F	228	GLY	-	expression tag	UNP Q586Y0
F	229	HIS	-	expression tag	UNP Q586Y0
F	230	HIS	-	expression tag	UNP Q586Y0
F	231	HIS	-	expression tag	UNP Q586Y0
F	232	HIS	-	expression tag	UNP Q586Y0
F	233	HIS	-	expression tag	UNP Q586Y0
F	234	HIS	-	expression tag	UNP Q586Y0
F	235	MET	-	expression tag	UNP Q586Y0
A	227	MET	-	initiating methionine	UNP Q586Y0
A	228	GLY	-	expression tag	UNP Q586Y0
A	229	HIS	-	expression tag	UNP Q586Y0
A	230	HIS	-	expression tag	UNP Q586Y0
A	231	HIS	-	expression tag	UNP Q586Y0
A	232	HIS	-	expression tag	UNP Q586Y0
A	233	HIS	-	expression tag	UNP Q586Y0
A	234	HIS	-	expression tag	UNP Q586Y0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	MET	-	expression tag	UNP Q586Y0
B	227	MET	-	initiating methionine	UNP Q586Y0
B	228	GLY	-	expression tag	UNP Q586Y0
B	229	HIS	-	expression tag	UNP Q586Y0
B	230	HIS	-	expression tag	UNP Q586Y0
B	231	HIS	-	expression tag	UNP Q586Y0
B	232	HIS	-	expression tag	UNP Q586Y0
B	233	HIS	-	expression tag	UNP Q586Y0
B	234	HIS	-	expression tag	UNP Q586Y0
B	235	MET	-	expression tag	UNP Q586Y0
C	227	MET	-	initiating methionine	UNP Q586Y0
C	228	GLY	-	expression tag	UNP Q586Y0
C	229	HIS	-	expression tag	UNP Q586Y0
C	230	HIS	-	expression tag	UNP Q586Y0
C	231	HIS	-	expression tag	UNP Q586Y0
C	232	HIS	-	expression tag	UNP Q586Y0
C	233	HIS	-	expression tag	UNP Q586Y0
C	234	HIS	-	expression tag	UNP Q586Y0
C	235	MET	-	expression tag	UNP Q586Y0
D	227	MET	-	initiating methionine	UNP Q586Y0
D	228	GLY	-	expression tag	UNP Q586Y0
D	229	HIS	-	expression tag	UNP Q586Y0
D	230	HIS	-	expression tag	UNP Q586Y0
D	231	HIS	-	expression tag	UNP Q586Y0
D	232	HIS	-	expression tag	UNP Q586Y0
D	233	HIS	-	expression tag	UNP Q586Y0
D	234	HIS	-	expression tag	UNP Q586Y0
D	235	MET	-	expression tag	UNP Q586Y0
E	227	MET	-	initiating methionine	UNP Q586Y0
E	228	GLY	-	expression tag	UNP Q586Y0
E	229	HIS	-	expression tag	UNP Q586Y0
E	230	HIS	-	expression tag	UNP Q586Y0
E	231	HIS	-	expression tag	UNP Q586Y0
E	232	HIS	-	expression tag	UNP Q586Y0
E	233	HIS	-	expression tag	UNP Q586Y0
E	234	HIS	-	expression tag	UNP Q586Y0
E	235	MET	-	expression tag	UNP Q586Y0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	F	15	Total O 15 15	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	A	15	Total 15	O 15	0	0
2	B	13	Total 13	O 13	0	0
2	C	13	Total 13	O 13	0	0
2	D	11	Total 11	O 11	0	0
2	E	11	Total 11	O 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription elongation factor s-II




- Molecule 1: Transcription elongation factor s-II




- Molecule 1: Transcription elongation factor s-II



- Molecule 1: Transcription elongation factor s-II



- Molecule 1: Transcription elongation factor s-II



ALA  
LYS  
ARG  
ALA  
LEU  
GLY  
THR

- Molecule 1: Transcription elongation factor s-II

Chain E:  68% 6% 26%

MET  
GLY  
HIS  
HIS  
HIS  
HIS  
HIS  
HIS  
MET  
SER  
SER  
ALA  
ASP  
ASP  
ASP  
VAL  
VAL  
TYR  
VAL  
VAL  
ASP  
MET  
LEU  
ARG  
LYS  
SER  
ARG  
HIS  
D254  
N276  
GLY  
D278  
V279  
R289  
L304  
I308  
R330  
F331  
E346  
A353  
LYS  
ARG  
ALA  
LEU  
GLY  
THR

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.72Å 111.72Å 93.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.27 – 2.25 42.27 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.4 (42.27-2.25) 99.4 (42.27-2.25)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	59.94 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.199 , 0.230 0.200 , 0.227	Depositor DCC
$R_{free}$ test set	1992 reflections (6.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.9	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.177 for -h,-k,l	Xtrriage
Reported twinning fraction	0.220 for -h,-k,l	Depositor
Outliers	1 of 32323 reflections (0.003%)	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4519	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.27	0/782	0.47	0/1066
1	B	0.35	0/742	0.46	0/1014
1	C	0.42	0/722	0.51	0/984
1	D	0.46	1/733 (0.1%)	0.50	0/996
1	E	0.32	0/741	0.50	0/1012
1	F	0.25	0/784	0.41	0/1065
All	All	0.35	1/4504 (0.0%)	0.47	0/6137

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	324	PHE	C-N	7.66	1.48	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	771	0	754	10	0
1	B	731	0	700	2	0
1	C	712	0	679	13	0
1	D	723	0	697	9	0
1	E	731	0	692	4	0
1	F	773	0	753	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	15	0	0	0	0
2	B	13	0	0	0	0
2	C	13	0	0	1	0
2	D	11	0	0	0	0
2	E	11	0	0	0	0
2	F	15	0	0	0	0
All	All	4519	0	4275	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:280:VAL:HB	1:D:323:GLY:HA3	1.56	0.87
1:C:268:THR:HG23	1:C:318:GLN:HE22	1.49	0.77
1:C:353:ALA:O	1:D:330:ARG:NH2	2.20	0.73
1:F:268:THR:HG23	1:F:318:GLN:HE22	1.56	0.70
1:D:280:VAL:CB	1:D:323:GLY:HA3	2.25	0.66
1:C:317:GLY:HA2	1:C:320:LEU:HD12	1.79	0.63
1:C:255:VAL:HG11	1:C:300:ALA:HB1	1.84	0.58
1:F:268:THR:HG23	1:F:318:GLN:NE2	2.18	0.58
1:D:266:GLU:OE2	1:D:289:ARG:NH2	2.39	0.55
1:C:341:ARG:NH2	2:C:401:HOH:O	2.41	0.53
1:E:276:ASN:N	1:E:279:VAL:O	2.42	0.53
1:A:308:ILE:HG13	1:A:309:GLY:N	2.27	0.50
1:C:308:ILE:HG12	1:D:291:VAL:HG12	1.93	0.49
1:A:255:VAL:HG21	1:A:300:ALA:HB1	1.95	0.48
1:C:320:LEU:HD11	1:C:336:LEU:HD12	1.96	0.47
1:E:304:LEU:O	1:E:308:ILE:HD12	2.15	0.47
1:A:303:THR:HG23	1:A:306:GLN:H	1.81	0.46
1:F:258:ALA:O	1:F:262:THR:HG23	2.16	0.46
1:A:269:GLU:CG	1:A:289:ARG:HH12	2.30	0.45
1:A:266:GLU:O	1:A:269:GLU:HG2	2.17	0.45
1:C:307:LEU:HD23	1:C:307:LEU:HA	1.87	0.45
1:B:263:THR:HG23	1:B:293:LEU:HD22	1.98	0.44
1:A:270:THR:HA	1:A:273:GLU:HB2	2.00	0.44
1:D:267:LEU:HD22	1:D:319:PHE:CE2	2.53	0.44
1:D:340:PHE:HD1	1:D:343:LEU:HD12	1.83	0.44
1:B:275:LEU:HA	1:B:275:LEU:HD23	1.73	0.43
1:C:268:THR:HG23	1:C:318:GLN:NE2	2.26	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:265:LYS:O	1:F:265:LYS:HD3	2.18	0.43
1:A:337:ASP:HB3	1:E:346:GLU:HB3	2.01	0.43
1:C:308:ILE:HD11	1:D:295:SER:OG	2.21	0.41
1:A:269:GLU:HG2	1:A:289:ARG:HH12	1.86	0.41
1:C:271:TYR:HE1	1:C:324:PHE:HE2	1.68	0.41
1:F:303:THR:CG2	1:F:306:GLN:H	2.34	0.41
1:F:325:PRO:HA	1:F:326:PRO:HD3	1.96	0.41
1:F:311:LYS:HE3	1:F:311:LYS:HB3	1.68	0.41
1:A:303:THR:CG2	1:A:306:GLN:H	2.33	0.41
1:D:280:VAL:HB	1:D:323:GLY:CA	2.40	0.40
1:C:271:TYR:CE1	1:C:324:PHE:CE2	3.09	0.40
1:A:331:PHE:CE2	1:E:308:ILE:HD13	2.56	0.40
1:F:257:ASN:OD1	1:F:306:GLN:HG2	2.22	0.40
1:C:271:TYR:HE1	1:C:324:PHE:CE2	2.39	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	101/133 (76%)	99 (98%)	2 (2%)	0	100	100
1	B	97/133 (73%)	94 (97%)	3 (3%)	0	100	100
1	C	94/133 (71%)	94 (100%)	0	0	100	100
1	D	93/133 (70%)	92 (99%)	1 (1%)	0	100	100
1	E	95/133 (71%)	95 (100%)	0	0	100	100
1	F	101/133 (76%)	99 (98%)	2 (2%)	0	100	100
All	All	581/798 (73%)	573 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/115 (71%)	82 (100%)	0	100	100
1	B	76/115 (66%)	73 (96%)	3 (4%)	32	38
1	C	71/115 (62%)	69 (97%)	2 (3%)	43	52
1	D	76/115 (66%)	74 (97%)	2 (3%)	46	55
1	E	76/115 (66%)	73 (96%)	3 (4%)	32	38
1	F	81/115 (70%)	80 (99%)	1 (1%)	71	80
All	All	462/690 (67%)	451 (98%)	11 (2%)	49	58

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	355	ARG
1	B	305	LYS
1	B	321	SER
1	B	331	PHE
1	C	272	MET
1	C	320	LEU
1	D	305	LYS
1	D	331	PHE
1	E	289	ARG
1	E	330	ARG
1	E	331	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	318	GLN
1	C	286	ASN
1	C	318	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	103/133 (77%)	-0.03	0	100   100	32, 48, 96, 165	0
1	B	99/133 (74%)	-0.09	0	100   100	35, 51, 79, 102	0
1	C	98/133 (73%)	0.21	4 (4%)	37   40	32, 55, 94, 107	0
1	D	97/133 (72%)	0.00	1 (1%)	82   84	34, 54, 85, 106	0
1	E	99/133 (74%)	-0.08	0	100   100	32, 46, 82, 104	0
1	F	103/133 (77%)	-0.14	1 (0%)	82   84	37, 53, 79, 93	0
All	All	599/798 (75%)	-0.02	6 (1%)	82   84	32, 52, 87, 165	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	282	VAL	3.6
1	C	285	ALA	3.2
1	D	282	VAL	3.0
1	F	254	ASP	2.7
1	C	325	PRO	2.2
1	C	283	SER	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.