

wwPDB X-ray Structure Validation Summary Report (i)

Jun 22, 2024 – 08:42 PM EDT

PDB ID	:	4XK9
Title	:	Crystal structure of A-AChBP in complex with pinnatoxin G
Authors	:	Bourne, Y.; Sulzenbacher, G.; Marchot, P.
Deposited on		
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	228	88%	• 8%
	11	220	5%	• 070
1	В	228	88%	6% 7%
1	С	228	86%	5% 8%
1	D	228	84%	6% • 9%
1	Е	228	4%	• 7%



Mol	Chain	Length	Quality of chain		
1	Б	000	5%		
	F	228	88%	5%	7%
1	C	000	7%		
	G	228	5%	• •	9%
1	Н	228		= 0/	70/
	11	228	88%	5%	7%
1	Т	228	89%		7%
	1		4%	•	/ /0
1	J	228	88%	5%	7%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 18268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace				
1	А	210	Total	С	Ν	0	S	0	1	0				
	A	210	1680	1061	279	331	9	0	1	0				
1	В	213	Total	С	Ν	Ο	S	0	2	0				
	D	215	1702	1075	282	336	9	0	2	0				
1	С	209	Total	С	Ν	Ο	\mathbf{S}	0	1	0				
1	U	205	1672	1055	278	330	9	0	I	0				
1	D	207	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
1	D	201	1657	1046	274	328	9	0	1					
1	Е	212	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
	Ц	Ц		Ц	Ц		1691	1068	279	335	9	<u> </u>	1	0
1	F	211	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
	T,	211	1692	1068	281	334	9	0	1	0				
1	G	208	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
	ŭ	200	1661	1049	274	329	9	0	T	0				
1	Н	212	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
	11	212	1693	1070	279	335	9	0	T	0				
1	Ι	211	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0				
	1	211	1686	1065	278	334	9		1	0				
1	J	212	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	2	0				
	0	212	1699	1073	282	335	9		<u> </u>	0				

• Molecule 1 is a protein called Soluble acetylcholine receptor.

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-8	ASP	-	expression tag	UNP Q8WSF8
А	-7	TYR	-	expression tag	UNP Q8WSF8
А	-6	LYS	-	expression tag	UNP Q8WSF8
А	-5	ASP	-	expression tag	UNP Q8WSF8
А	-4	ASP	-	expression tag	UNP Q8WSF8
А	-3	ASP	-	expression tag	UNP Q8WSF8
А	-2	ASP	-	expression tag	UNP Q8WSF8
А	-1	LYS	-	expression tag	UNP Q8WSF8
А	0	LEU	-	expression tag	UNP Q8WSF8




Chain	Residue	Modelled	Actual	Comment	Reference
В	-8	ASP	-	expression tag	UNP Q8WSF8
В	-7	TYR	-	expression tag	UNP Q8WSF8
В	-6	LYS	-	expression tag	UNP Q8WSF8
В	-5	ASP	-	expression tag	UNP Q8WSF8
В	-4	ASP	-	expression tag	UNP Q8WSF8
В	-3	ASP	_	expression tag	UNP Q8WSF8
В	-2	ASP	-	expression tag	UNP Q8WSF8
В	-1	LYS	_	expression tag	UNP Q8WSF8
В	0	LEU	_	expression tag	UNP Q8WSF8
С	-8	ASP	_	expression tag	UNP Q8WSF8
С	-7	TYR	-	expression tag	UNP Q8WSF8
С	-6	LYS	-	expression tag	UNP Q8WSF8
С	-5	ASP	-	expression tag	UNP Q8WSF8
С	-4	ASP	-	expression tag	UNP Q8WSF8
С	-3	ASP	-	expression tag	UNP Q8WSF8
С	-2	ASP	-	expression tag	UNP Q8WSF8
С	-1	LYS	-	expression tag	UNP Q8WSF8
С	0	LEU	_	expression tag	UNP Q8WSF8
D	-8	ASP	_	expression tag	UNP Q8WSF8
D	-7	TYR	_	expression tag	UNP Q8WSF8
D	-6	LYS	_	expression tag	UNP Q8WSF8
D	-5	ASP	_	expression tag	UNP Q8WSF8
D	-4	ASP	_	expression tag	UNP Q8WSF8
D	-3	ASP	_	expression tag	UNP Q8WSF8
D	-2	ASP	_	expression tag	UNP Q8WSF8
D	-1	LYS	_	expression tag	UNP Q8WSF8
D	0	LEU	_	expression tag	UNP Q8WSF8
Е	-8	ASP	_	expression tag	UNP Q8WSF8
Е	-7	TYR	_	expression tag	UNP Q8WSF8
Е	-6	LYS	_	expression tag	UNP Q8WSF8
Е	-5	ASP	_	expression tag	UNP Q8WSF8
Е	-4	ASP	_	expression tag	UNP Q8WSF8
Е	-3	ASP	_	expression tag	UNP Q8WSF8
Е	-2	ASP	_	expression tag	UNP Q8WSF8
Е	-1	LYS	_	expression tag	UNP Q8WSF8
Е	0	LEU	_	expression tag	UNP Q8WSF8
F	-8	ASP	_	expression tag	UNP Q8WSF8
F	-7	TYR	-	expression tag	UNP Q8WSF8
F	-6	LYS	_	expression tag	UNP Q8WSF8
F	-5	ASP	_	expression tag	UNP Q8WSF8
		ASP		expression tag	UNP Q8WSF8
\mathbf{F}	-4	ADI	-	CAPICOSION UNE	



Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference	
F	-2	ASP	-	expression tag	UNP Q8WSF8	
F	-1	LYS	-	expression tag	UNP Q8WSF8	
F	0	LEU	-	expression tag	UNP Q8WSF8	
G	-8	ASP	-	expression tag	UNP Q8WSF8	
G	-7	TYR	-	expression tag	UNP Q8WSF8	
G	-6	LYS	-	expression tag	UNP Q8WSF8	
G	-5	ASP	-	expression tag	UNP Q8WSF8	
G	-4	ASP	-	expression tag	UNP Q8WSF8	
G	-3	ASP	-	expression tag	UNP Q8WSF8	
G	-2	ASP	-	expression tag	UNP Q8WSF8	
G	-1	LYS	-	expression tag	UNP Q8WSF8	
G	0	LEU	-	expression tag	UNP Q8WSF8	
Η	-8	ASP	-	expression tag	UNP Q8WSF8	
Н	-7	TYR	-	expression tag	UNP Q8WSF8	
H	-6	LYS	-	expression tag	UNP Q8WSF8	
Н	-5	ASP	-	expression tag	UNP Q8WSF8	
Н	-4	ASP	-	expression tag	UNP Q8WSF8	
Н	-3	ASP	-	expression tag	UNP Q8WSF8	
Н	-2	ASP	-	expression tag	UNP Q8WSF8	
Н	-1	LYS	-	expression tag	UNP Q8WSF8	
Н	0	LEU	-	expression tag	UNP Q8WSF8	
Ι	-8	ASP	-	expression tag	UNP Q8WSF8	
Ι	-7	TYR	-	expression tag	UNP Q8WSF8	
Ι	-6	LYS	-	expression tag	UNP Q8WSF8	
Ι	-5	ASP	-	expression tag	UNP Q8WSF8	
Ι	-4	ASP	-	expression tag	UNP Q8WSF8	
Ι	-3	ASP	-	expression tag	UNP Q8WSF8	
Ι	-2	ASP	-	expression tag	UNP Q8WSF8	
Ι	-1	LYS	-	expression tag	UNP Q8WSF8	
Ι	0	LEU	-	expression tag	UNP Q8WSF8	
J	-8	ASP	-	expression tag	UNP Q8WSF8	
J	-7	TYR	_	expression tag	UNP Q8WSF8	
J	-6	LYS	-	expression tag	UNP Q8WSF8	
J	-5	ASP	-	expression tag	UNP Q8WSF8	
J	-4	ASP	-	expression tag	UNP Q8WSF8	
J	-3	ASP	-	expression tag	UNP Q8WSF8	
J	-2	ASP	-	expression tag	UNP Q8WSF8	
J	-1	LYS	-	expression tag	UNP Q8WSF8	
J	0	LEU	-	expression tag	UNP Q8WSF8	

• Molecule 2 is Pinnatoxin G (three-letter code: 41J) (formula: $C_{42}H_{63}NO_7$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf				
2	А	1	Total C N O	0	0				
		-	50 42 1 7	Ŭ	Ŭ				
2	В	1	Total C N O	0	0				
	D	1	50 42 1 7	0					
2	С	1	Total C N O	0	0				
2	U	1	50 42 1 7	0	0				
2	D	1	Total C N O	0	0				
	D	L L	50 42 1 7	0	0				
2	Е	1	Total C N O	0	0				
	Ľ	Ľ				Ŧ	50 42 1 7	0	0
2	F	1	Total C N O	0	0				
	Г	1	50 42 1 7	0	0				
2	G	1	Total C N O	0	0				
	G	1	50 42 1 7	0	0				
2	Н	1	Total C N O	0	0				
	п	1	50 42 1 7	0	0				
2	Ι	1	Total C N O	0	0				
		1	50 42 1 7	U	0				
2	J	1	Total C N O	0	0				
	J	L	50 42 1 7	U	0				

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Cl 1 1	0	0
3	В	1	Total Cl 1 1	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	1	Total Cl 1 1	0	0
3	Ε	1	Total Cl 1 1	0	0
3	G	1	Total Cl 1 1	0	0
3	Н	1	Total Cl 1 1	0	0
3	J	1	Total Cl 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	93	Total O 93 93	0	0
4	В	88	Total O 88 88	0	0
4	С	91	Total O 91 91	0	0
4	D	79	Total O 79 79	0	0
4	Е	108	Total O 108 108	0	0
4	F	81	Total O 81 81	0	0
4	G	118	Total O 118 118	0	0
4	Н	102	Total O 102 102	0	0
4	Ι	74	Total O 74 74	0	0
4	J	94	Total O 94 94	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 88% 8% ASP LYS ASP ASP ASP ASP • Molecule 1: Soluble acetylcholine receptor Chain B: 88% 6% 7% • Molecule 1: Soluble acetylcholine receptor Chain C: 86% 5% 8% ALA GLY ASN GLY ASP LYS ASP ASP ASP ASP ASP LYS • Molecule 1: Soluble acetylcholine receptor Chain D: 84% 9% 6% · GLY PHE PHE ASN ASN LEU LEU ASP • Molecule 1: Soluble acetylcholine receptor Chain E: 89% 7% ASP LYS ASP
- Molecule 1: Soluble acetylcholine receptor







4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	184.84Å 142.92Å 142.13Å	Depositor
a, b, c, α , β , γ	90.00° 126.48° 90.00°	Depositor
Resolution (Å)	34.00 - 2.20	Depositor
Resolution (A)	33.62 - 2.20	EDS
% Data completeness	99.6 (34.00-2.20)	Depositor
(in resolution range)	99.6 (33.62-2.20)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.07	Depositor
$< I/\sigma(I) > 1$	$2.36 (at 2.20 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
D D.	0.199 , 0.211	Depositor
R, R_{free}	0.211 , 0.224	DCC
R_{free} test set	7513 reflections (5.02%)	wwPDB-VP
Wilson B-factor $(Å^2)$	31.8	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, 39.4	EDS
L-test for twinning ²	$ L > = 0.47, < L^2 > = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18268	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 30.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2007e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 41J, $\rm CL$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	B	ond angles
IVI0I	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.48	0/1724	0.79	3/2351~(0.1%)
1	В	0.49	0/1749	0.75	1/2386~(0.0%)
1	С	0.50	0/1716	0.81	1/2340~(0.0%)
1	D	0.58	3/1700~(0.2%)	0.87	7/2318~(0.3%)
1	Ε	0.50	0/1735	0.77	2/2366~(0.1%)
1	F	0.46	0/1736	0.78	4/2366~(0.2%)
1	G	0.51	0/1705	0.80	4/2326~(0.2%)
1	Н	0.50	0/1737	0.76	1/2368~(0.0%)
1	Ι	0.50	0/1730	0.81	4/2359~(0.2%)
1	J	0.47	0/1746	0.75	0/2380
All	All	0.50	3/17278~(0.0%)	0.79	27/23560~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	D	0	1
1	F	0	1
1	Н	0	1
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	206	GLU	CD-OE2	-7.70	1.17	1.25
1	D	206	GLU	CD-OE1	7.02	1.33	1.25
1	D	193	GLU	CD-OE1	-6.69	1.18	1.25



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	С	27	ASP	CB-CG-OD2	10.29	127.56	118.30
1	G	207	ARG	NE-CZ-NH1	9.52	125.06	120.30
1	D	206	GLU	CG-CD-OE1	8.34	134.99	118.30
1	D	206	GLU	CG-CD-OE2	-7.96	102.37	118.30
1	В	-2	ASP	CB-CG-OD1	7.74	125.26	118.30

The worst 5 of 27 bond angle outliers are listed below:

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	190	CYS	Peptide
1	D	190	CYS	Peptide
1	F	190	CYS	Peptide
1	Н	190	CYS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1680	0	1615	5	0
1	В	1702	0	1632	14	0
1	С	1672	0	1604	10	0
1	D	1657	0	1592	8	0
1	Е	1691	0	1621	5	0
1	F	1692	0	1630	7	0
1	G	1661	0	1591	8	0
1	Н	1693	0	1628	6	0
1	Ι	1686	0	1619	4	0
1	J	1699	0	1634	10	0
2	А	50	0	63	0	0
2	В	50	0	63	0	0
2	С	50	0	63	0	0
2	D	50	0	63	0	0
2	Е	50	0	63	0	0
2	F	50	0	63	1	0
2	G	50	0	63	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Н	50	0	63	0	0
2	Ι	50	0	63	0	0
2	J	50	0	63	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	Ε	1	0	0	0	0
3	G	1	0	0	0	0
3	Н	1	0	0	0	0
3	J	1	0	0	0	0
4	А	93	0	0	0	0
4	В	88	0	0	0	0
4	С	91	0	0	0	0
4	D	79	0	0	0	0
4	Ε	108	0	0	0	0
4	F	81	0	0	0	0
4	G	118	0	0	1	0
4	Н	102	0	0	1	0
4	Ι	74	0	0	0	0
4	J	94	0	0	0	0
All	All	18268	0	16796	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 60 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170:ALA:O	1:C:207:ARG:NH1	1.90	1.05
1:J:97[B]:ARG:HH11	1:J:97[B]:ARG:HG2	1.41	0.85
1:J:97[B]:ARG:HH11	1:J:97[B]:ARG:CG	1.97	0.77
1:J:170:ALA:O	1:J:207[B]:ARG:NH1	2.20	0.75
1:F:7:MET:CE	1:G:18:PRO:HB2	2.21	0.70

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	209/228~(92%)	206 (99%)	3~(1%)	0	100	100
1	В	213/228~(93%)	211 (99%)	2(1%)	0	100	100
1	\mathbf{C}	208/228~(91%)	205~(99%)	3~(1%)	0	100	100
1	D	206/228~(90%)	203~(98%)	3~(2%)	0	100	100
1	Ε	211/228~(92%)	208~(99%)	3~(1%)	0	100	100
1	F	210/228~(92%)	207~(99%)	3~(1%)	0	100	100
1	G	207/228~(91%)	204 (99%)	3~(1%)	0	100	100
1	Η	211/228~(92%)	207~(98%)	4(2%)	0	100	100
1	Ι	210/228~(92%)	207~(99%)	3~(1%)	0	100	100
1	J	212/228~(93%)	209~(99%)	3~(1%)	0	100	100
All	All	2097/2280~(92%)	2067 (99%)	30 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	191/206~(93%)	191 (100%)	0	100	100
1	В	193/206~(94%)	193 (100%)	0	100	100
1	С	190/206~(92%)	189 (100%)	1 (0%)	88	94
1	D	189/206~(92%)	188 (100%)	1 (0%)	88	94



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Ε	192/206~(93%)	190~(99%)	2(1%)	76 86
1	F	193/206~(94%)	193 (100%)	0	100 100
1	G	189/206~(92%)	188 (100%)	1 (0%)	88 94
1	Н	193/206~(94%)	192 (100%)	1 (0%)	88 94
1	Ι	192/206~(93%)	192 (100%)	0	100 100
1	J	193/206~(94%)	191 (99%)	2(1%)	76 86
All	All	1915/2060~(93%)	1907 (100%)	8 (0%)	91 96

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	193	GLU
1	J	74	ASN
1	G	193	GLU
1	Е	193	GLU
1	Н	193	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	105	GLN
1	J	74	ASN
1	G	187	HIS
1	Е	105	GLN
1	Н	15	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 7 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	41J	F	301	-	$51,\!57,\!57$	1.66	3 (5%)	50,91,91	1.57	7 (14%)
2	41J	Н	301	-	$51,\!57,\!57$	1.63	2 (3%)	50,91,91	1.65	6 (12%)
2	41J	G	301	-	$51,\!57,\!57$	1.60	2 (3%)	50,91,91	1.61	5 (10%)
2	41J	Е	301	-	$51,\!57,\!57$	1.64	2 (3%)	50,91,91	1.56	7 (14%)
2	41J	В	301	-	$51,\!57,\!57$	1.63	2 (3%)	50,91,91	1.63	8 (16%)
2	41J	J	301	-	$51,\!57,\!57$	1.59	2 (3%)	50,91,91	1.53	8 (16%)
2	41J	Ι	301	-	$51,\!57,\!57$	1.69	5 (9%)	50,91,91	1.72	7 (14%)
2	41J	С	301	-	$51,\!57,\!57$	1.64	4 (7%)	50,91,91	1.70	<u>6 (12%)</u>
2	41J	D	301	-	$51,\!57,\!57$	1.64	3 (5%)	50,91,91	1.66	7 (14%)
2	41J	А	301	-	$51,\!57,\!57$	1.64	2 (3%)	50,91,91	1.64	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	41J	F	301	-	-	1/18/129/129	-
2	41J	Н	301	-	-	0/18/129/129	-
2	41J	G	301	-	-	0/18/129/129	-
2	41J	Е	301	-	-	1/18/129/129	-
2	41J	В	301	-	-	3/18/129/129	-
2	41J	J	301	-	-	0/18/129/129	-
2	41J	Ι	301	-	-	0/18/129/129	-
2	41J	С	301	-	-	0/18/129/129	-
2	41J	D	301	-	-	0/18/129/129	-
2	41J	А	301	-	-	2/18/129/129	-

The worst 5 of 27 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(\text{\AA})$	Ideal(Å)
2	Ε	301	41J	C35-C36	8.17	1.52	1.33
2	Ι	301	41J	C35-C36	8.11	1.52	1.33
2	G	301	41J	C35-C36	8.02	1.52	1.33
2	F	301	41J	C35-C36	8.00	1.52	1.33
2	D	301	41J	C35-C36	7.99	1.52	1.33

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	301	41J	O16-C15-C12	6.17	116.69	109.07
2	Н	301	41J	O16-C15-C12	5.95	116.41	109.07
2	В	301	41J	O16-C15-C12	5.72	116.13	109.07
2	F	301	41J	O16-C15-C12	5.52	115.89	109.07
2	Ι	301	41J	O16-C15-C12	5.48	115.84	109.07

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	301	41J	C38-C36-C37-C48
2	А	301	41J	O16-C17-C18-C21
2	Е	301	41J	C38-C36-C37-C48
2	F	301	41J	C38-C36-C37-C48
2	В	301	41J	C35-C36-C37-C48


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	41J	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient must be highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	210/228~(92%)	-0.01	10 (4%) 30 29	25, 44, 98, 138	0
1	В	213/228~(93%)	0.12	11 (5%) 27 26	28, 46, 86, 103	0
1	С	209/228~(91%)	-0.01	9 (4%) 35 33	27, 46, 86, 113	0
1	D	207/228~(90%)	0.30	17 (8%) 11 10	26, 50, 90, 122	0
1	Е	212/228~(92%)	-0.13	9 (4%) 36 34	24, 39, 75, 106	0
1	F	211/228~(92%)	0.18	11 (5%) 27 26	25, 50, 95, 132	0
1	G	208/228~(91%)	0.07	15 (7%) 15 14	24, 40, 87, 128	0
1	Н	212/228~(92%)	-0.06	12 (5%) 23 22	26, 45, 78, 109	0
1	Ι	211/228~(92%)	0.21	16 (7%) 13 12	27, 51, 96, 140	0
1	J	212/228~(92%)	0.00	9 (4%) 36 34	26, 43, 82, 106	0
All	All	2105/2280 (92%)	0.07	119 (5%) 23 22	24, 46, 89, 140	0

The worst 5 of 119 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	0	LEU	7.0
1	Ι	-1	LYS	7.0
1	G	1	HIS	6.0
1	D	208	ARG	6.0
1	Ι	-2	ASP	5.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	А	302	1/1	0.90	0.11	48,48,48,48	0
2	41J	D	301	50/50	0.94	0.10	28,31,33,34	0
2	41J	Ι	301	50/50	0.94	0.10	28,32,33,34	0
2	41J	В	301	50/50	0.94	0.11	27,29,31,32	0
3	CL	С	302	1/1	0.94	0.11	45,45,45,45	0
2	41J	Н	301	50/50	0.95	0.09	20,22,23,24	0
2	41J	F	301	50/50	0.95	0.09	23,28,29,31	0
2	41J	С	301	50/50	0.96	0.09	21,23,26,27	0
2	41J	J	301	50/50	0.96	0.09	23,24,27,28	0
2	41J	G	301	50/50	0.96	0.09	22,24,26,26	0
2	41J	А	301	50/50	0.96	0.10	20,22,23,24	0
3	CL	G	302	1/1	0.96	0.05	44,44,44,44	0
2	41J	Е	301	50/50	0.97	0.07	22,24,25,26	0
3	CL	J	302	1/1	0.97	0.05	40,40,40,40	0
3	CL	В	302	1/1	0.98	0.04	39,39,39,39	0
3	CL	Е	302	1/1	0.98	0.03	34,34,34,34	0
3	CL	Н	302	1/1	0.99	0.04	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


























6.5 Other polymers (i)

There are no such residues in this entry.

