



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 03:41 PM JST

PDB ID : 5XMV
Title : Plasmodium vivax SHMT bound with PLP-glycine and GS362
Authors : Chitnumsub, P.; Jaruwat, A.; Leartsakulpanich, U.; Schwertz, G.; Diederich, F.
Deposited on : 2017-05-16
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	
1	B	442	
1	C	442	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 10915 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	B	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	C	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			

- Molecule 2 is N-GLYCINE-[3-HYDROXY-2-METHYL-5-PHOSPHONOXYMETHYL-PYRIDIN-4-YL-METHANE] (three-letter code: PLG) (formula: C₁₀H₁₅N₂O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	B	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	C	1	Total	C	N	O	P	0	0
			20	10	2	7	1		

- Molecule 3 is (4 {S})-6-azanyl-4-[3-(2-chlorophenyl)-5-(trifluoromethyl)phenyl]-3-methyl-4-propan-2-yl-2 {H}-pyrano[2,3-c]pyrazole-5- carbonitrile (three-letter code: 8AU) (formula: C₂₄H₂₀ClF₃N₄O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	Cl	F	N	O	0	0
			33	24	1	3	4	1		
3	B	1	Total	C	Cl	F	N	O	0	0
			33	24	1	3	4	1		
3	C	1	Total	C	Cl	F	N	O	0	0
			33	24	1	3	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	120	Total	O	0	0
			120	120		
5	B	133	Total	O	0	0
			133	133		
5	C	127	Total	O	0	0
			127	127		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase




- Molecule 1: Serine hydroxymethyltransferase



- Molecule 1: Serine hydroxymethyltransferase





4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.67Å 58.84Å 234.90Å 90.00° 90.14° 90.00°	Depositor
Resolution (Å)	30.00 – 2.16 29.19 – 2.16	Depositor EDS
% Data completeness (in resolution range)	93.3 (30.00-2.16) 92.7 (29.19-2.16)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.242 , 0.306 0.244 , 0.301	Depositor DCC
R_{free} test set	6878 reflections (9.78%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtrriage
Anisotropy	0.052	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 12.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.018 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.018 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.459 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.448 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.015 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10915	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PLG, 8AU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/3521	0.72	3/4754 (0.1%)
1	B	0.56	0/3521	0.70	1/4754 (0.0%)
1	C	0.57	0/3521	0.72	2/4754 (0.0%)
All	All	0.57	0/10563	0.71	6/14262 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	C	240	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	240	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	C	240	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	240	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	208	ASP	CB-CG-OD1	5.06	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3477	16	0
1	B	3458	0	3477	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3458	0	3477	17	0
2	A	20	0	11	0	0
2	B	20	0	12	1	0
2	C	20	0	12	1	0
3	A	33	0	0	0	0
3	B	33	0	0	0	0
3	C	33	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	120	0	0	0	0
5	B	133	0	0	0	0
5	C	127	0	0	1	0
All	All	10915	0	10466	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:GLU:OE2	1:A:26:ARG:NH2	2.29	0.66
1:C:229:ASP:OD1	1:C:253:ARG:HD2	1.99	0.61
1:B:385:LYS:HE3	1:B:385:LYS:H	1.69	0.58
1:A:385:LYS:H	1:A:385:LYS:HE3	1.70	0.56
1:C:234:THR:HB	1:C:236:HIS:CE1	2.43	0.53
1:B:109:LEU:HD21	1:B:179:ILE:HD11	1.90	0.53
1:B:113:VAL:CG1	1:B:177:VAL:HG23	2.41	0.51
1:C:153:TYR:HB3	1:C:169:MET:HE2	1.93	0.51
1:B:385:LYS:H	1:B:385:LYS:CE	2.24	0.51
1:C:174:LYS:HD2	1:C:201:VAL:HG13	1.92	0.51
1:A:7:LEU:HA	1:A:10:ILE:CG2	2.42	0.49
1:C:85:PHE:HB3	1:C:225:PHE:CE2	2.47	0.49
1:C:7:LEU:HA	1:C:10:ILE:HG22	1.95	0.49
1:B:315:GLN:O	1:B:334:LYS:HE2	2.12	0.49
1:A:385:LYS:H	1:A:385:LYS:CE	2.26	0.49
1:C:109:LEU:HD22	1:C:119:ILE:HG21	1.95	0.49
1:A:234:THR:HB	1:A:236:HIS:CE1	2.47	0.49
1:A:193:GLN:O	1:A:197:ILE:HG13	2.12	0.48
1:A:113:VAL:CG1	1:A:177:VAL:HG23	2.42	0.48
1:B:229:ASP:OD1	1:B:253:ARG:HD2	2.14	0.48
1:C:153:TYR:HB3	1:C:169:MET:CE	2.43	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:363:ASP:HB3	1:C:365:VAL:HG22	1.96	0.47
1:B:251:LYS:HD2	1:B:258:GLU:OE1	2.14	0.47
1:B:321:ASN:HD21	1:C:424:GLN:HE22	1.62	0.47
1:A:7:LEU:HA	1:A:10:ILE:HG22	1.96	0.47
1:B:3:ASN:ND2	1:B:10:ILE:HD12	2.29	0.46
1:A:109:LEU:O	1:A:113:VAL:HG22	2.16	0.46
1:B:167:ARG:HD3	1:B:197:ILE:HD13	1.98	0.46
1:C:235:THR:O	1:C:241:GLY:N	2.45	0.45
1:B:398:ILE:O	1:B:402:VAL:HG23	2.17	0.45
1:C:184:SER:HB3	1:C:371:ARG:HD3	1.99	0.45
1:B:161:VAL:HG23	1:B:185:TYR:CZ	2.52	0.45
1:C:181:GLY:HA3	1:C:208:ASP:O	2.16	0.45
1:B:22:GLU:OE2	1:B:26:ARG:NH2	2.51	0.44
1:B:195:ARG:HB2	1:B:227:HIS:HB3	2.00	0.44
1:B:319:VAL:HG12	1:B:320:THR:HG23	2.00	0.44
1:B:234:THR:HB	1:B:236:HIS:CE1	2.52	0.43
1:C:249:PHE:HB3	1:C:261:ILE:HD12	2.01	0.43
2:C:501:PLG:H4A1	5:C:666:HOH:O	2.19	0.43
1:A:127:GLY:O	1:A:182:TYR:HB3	2.19	0.42
1:A:136:ASP:OD1	1:A:137:GLU:N	2.52	0.42
1:C:398:ILE:O	1:C:402:VAL:HG23	2.20	0.42
1:A:400:ILE:HG21	1:A:425:LEU:HD21	2.02	0.42
1:A:3:ASN:ND2	1:A:10:ILE:HD13	2.33	0.42
1:A:209:ILE:HG13	1:A:233:THR:HB	2.02	0.42
1:B:62:ARG:HB3	1:B:64:TYR:O	2.20	0.42
1:B:343:GLN:OE1	1:B:354:ASN:HA	2.20	0.42
1:C:125:CYS:HG	1:C:364:CYS:HG	1.66	0.42
1:B:396:ARG:CZ	1:B:428:LEU:HD13	2.50	0.42
1:B:181:GLY:HA3	1:B:208:ASP:O	2.20	0.42
1:C:120:MET:SD	1:C:169:MET:HE3	2.60	0.41
1:A:10:ILE:HG21	1:B:41:ALA:CB	2.50	0.41
1:B:39:ASN:OD1	1:B:42:VAL:HG23	2.20	0.41
1:A:398:ILE:O	1:A:402:VAL:HG23	2.21	0.41
1:B:274:HIS:O	1:B:277:LYS:HB2	2.21	0.41
1:B:97:GLN:N	1:B:98:PRO:CD	2.84	0.40
1:B:209:ILE:HD12	1:B:213:SER:HA	2.03	0.40
1:B:34:SER:OG	2:B:501:PLG:O	2.31	0.40
1:B:95:ASN:OD1	1:B:98:PRO:HD3	2.22	0.40
1:B:91:GLU:HG3	1:B:252:LYS:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/442 (100%)	427 (97%)	12 (3%)	1 (0%)	47	46
1	B	440/442 (100%)	425 (97%)	14 (3%)	1 (0%)	47	46
1	C	440/442 (100%)	429 (98%)	10 (2%)	1 (0%)	47	46
All	All	1320/1326 (100%)	1281 (97%)	36 (3%)	3 (0%)	47	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	LYS
1	C	237	LYS
1	B	237	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/381 (100%)	354 (93%)	27 (7%)	14	9
1	B	381/381 (100%)	361 (95%)	20 (5%)	23	19
1	C	381/381 (100%)	355 (93%)	26 (7%)	16	10
All	All	1143/1143 (100%)	1070 (94%)	73 (6%)	17	12

All (73) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	GLU
1	A	54	TYR
1	A	62	ARG
1	A	79	LYS
1	A	90	GLU
1	A	100	SER
1	A	138	LYS
1	A	151	LYS
1	A	167	ARG
1	A	172	SER
1	A	176	LYS
1	A	183	THR
1	A	193	GLN
1	A	196	GLN
1	A	250	ASN
1	A	254	ASN
1	A	261	ILE
1	A	327	LEU
1	A	352	SER
1	A	356	ASN
1	A	360	SER
1	A	364	CYS
1	A	385	LYS
1	A	405	GLN
1	A	413	VAL
1	A	417	LYS
1	A	430	GLN
1	B	54	TYR
1	B	62	ARG
1	B	100	SER
1	B	138	LYS
1	B	139	LYS
1	B	167	ARG
1	B	183	THR
1	B	197	ILE
1	B	250	ASN
1	B	252	LYS
1	B	254	ASN
1	B	295	GLU
1	B	327	LEU
1	B	334	LYS
1	B	352	SER
1	B	364	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	385	LYS
1	B	413	VAL
1	B	427	GLN
1	B	430	GLN
1	C	21	GLU
1	C	54	TYR
1	C	62	ARG
1	C	89	ASP
1	C	100	SER
1	C	136	ASP
1	C	138	LYS
1	C	167	ARG
1	C	169	MET
1	C	176	LYS
1	C	183	THR
1	C	193	GLN
1	C	196	GLN
1	C	206	PHE
1	C	250	ASN
1	C	254	ASN
1	C	327	LEU
1	C	347	ASN
1	C	352	SER
1	C	355	LYS
1	C	357	THR
1	C	364	CYS
1	C	385	LYS
1	C	405	GLN
1	C	427	GLN
1	C	439	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	299	GLN
1	A	347	ASN
1	A	427	GLN
1	B	250	ASN
1	B	347	ASN
1	C	9	GLN
1	C	123	HIS
1	C	250	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	347	ASN
1	C	424	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLG	C	501	-	20,20,20	2.76	3 (15%)	25,28,28	2.49	7 (28%)
2	PLG	A	501	-	20,20,20	2.75	4 (20%)	25,28,28	2.27	9 (36%)
3	8AU	C	502	-	31,36,36	2.67	4 (12%)	43,56,56	1.60	8 (18%)
2	PLG	B	501	-	20,20,20	2.79	3 (15%)	25,28,28	2.81	9 (36%)
3	8AU	A	502	-	31,36,36	2.67	5 (16%)	43,56,56	1.63	9 (20%)
3	8AU	B	502	-	31,36,36	2.55	4 (12%)	43,56,56	1.57	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLG	C	501	-	-	6/12/12/12	0/1/1/1
2	PLG	A	501	-	-	4/12/12/12	0/1/1/1
3	8AU	C	502	-	-	0/22/44/44	0/4/4/4
2	PLG	B	501	-	-	5/12/12/12	0/1/1/1
3	8AU	A	502	-	-	1/22/44/44	0/4/4/4
3	8AU	B	502	-	-	1/22/44/44	0/4/4/4

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	8AU	C8-C3	-11.09	1.27	1.42
3	C	502	8AU	C8-C3	-10.82	1.27	1.42
3	B	502	8AU	C8-C3	-10.42	1.28	1.42
2	A	501	PLG	C3-C2	9.19	1.50	1.40
2	B	501	PLG	C3-C2	9.03	1.49	1.40
2	C	501	PLG	C3-C2	8.78	1.49	1.40
3	C	502	8AU	O1-C2	7.80	1.45	1.36
3	B	502	8AU	O1-C2	7.03	1.44	1.36
3	A	502	8AU	O1-C2	6.79	1.44	1.36
2	C	501	PLG	C5-C4	5.92	1.48	1.40
2	B	501	PLG	C5-C4	5.76	1.48	1.40
2	A	501	PLG	C5-C4	5.66	1.48	1.40
2	B	501	PLG	C3-C4	5.03	1.47	1.40
2	C	501	PLG	C3-C4	5.00	1.47	1.40
3	A	502	8AU	C22-C25	4.87	1.48	1.39
2	A	501	PLG	C3-C4	4.82	1.47	1.40
3	B	502	8AU	C22-C25	4.45	1.47	1.39
3	C	502	8AU	C22-C25	4.22	1.47	1.39
3	A	502	8AU	N12-N11	2.18	1.41	1.37
3	B	502	8AU	N12-N11	2.17	1.41	1.37
3	A	502	8AU	C25-CL3	2.12	1.78	1.73
3	C	502	8AU	N12-N11	2.02	1.41	1.37
2	A	501	PLG	OXT-C	-2.02	1.24	1.30

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	PLG	C4A-N-CA	11.43	124.51	112.58
2	C	501	PLG	C4A-N-CA	8.69	121.65	112.58
2	A	501	PLG	C4A-N-CA	7.66	120.57	112.58
3	C	502	8AU	O1-C2-N7	4.84	113.70	110.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	8AU	O1-C2-N7	4.67	113.58	110.22
2	C	501	PLG	OP4-C5A-C5	3.97	116.92	109.35
3	B	502	8AU	O1-C2-N7	3.85	112.98	110.22
3	A	502	8AU	C6-O1-C2	3.54	122.22	118.38
2	C	501	PLG	OP3-P-OP2	3.41	120.68	107.64
3	C	502	8AU	C3-C2-N7	-3.39	124.07	127.47
3	B	502	8AU	C14-C4-C3	-3.38	104.89	110.32
3	B	502	8AU	C5-C4-C3	3.28	111.01	104.34
3	B	502	8AU	C3-C2-N7	-3.25	124.22	127.47
3	A	502	8AU	C15-C14-C4	3.22	115.85	112.40
3	A	502	8AU	C5-C4-C3	3.22	110.89	104.34
2	B	501	PLG	OP4-C5A-C5	3.17	115.39	109.35
3	C	502	8AU	C24-C14-C4	3.09	115.71	112.40
3	C	502	8AU	C5-C4-C3	3.08	110.60	104.34
2	C	501	PLG	C4-C4A-N	3.02	117.22	111.58
2	A	501	PLG	C4A-C4-C3	2.96	123.22	120.04
2	A	501	PLG	C4-C4A-N	2.95	117.09	111.58
3	A	502	8AU	C3-C2-N7	-2.92	124.54	127.47
3	C	502	8AU	C10-C5-C6	-2.92	100.93	104.01
3	A	502	8AU	C10-C5-C6	-2.89	100.96	104.01
2	B	501	PLG	OP3-P-OP1	2.85	121.84	110.68
2	A	501	PLG	OP4-C5A-C5	2.81	114.71	109.35
2	B	501	PLG	OP3-P-OP4	-2.76	99.39	106.73
2	C	501	PLG	OXT-C-CA	2.72	122.19	112.74
3	A	502	8AU	C14-C4-C3	-2.68	106.02	110.32
2	B	501	PLG	C4A-C4-C3	2.65	122.88	120.04
3	B	502	8AU	C19-C20-C22	-2.63	116.25	120.61
3	B	502	8AU	C13-C10-N11	2.58	125.32	119.65
3	A	502	8AU	C13-C10-N11	2.48	125.10	119.65
3	B	502	8AU	C10-C5-C6	-2.44	101.43	104.01
3	B	502	8AU	C6-O1-C2	2.42	121.00	118.38
2	A	501	PLG	OP3-P-OP2	2.42	116.87	107.64
3	C	502	8AU	C14-C4-C3	-2.41	106.45	110.32
2	C	501	PLG	C4A-C4-C3	2.41	122.63	120.04
2	A	501	PLG	C3-C4-C5	-2.38	116.43	118.72
2	A	501	PLG	OXT-C-CA	2.35	120.89	112.74
3	A	502	8AU	C17-C18-C23	2.31	122.70	119.58
2	A	501	PLG	C6-N1-C2	2.31	123.44	119.17
2	C	501	PLG	C6-N1-C2	2.30	123.44	119.17
2	B	501	PLG	C4-C4A-N	2.24	115.76	111.58
2	B	501	PLG	C6-N1-C2	2.22	123.27	119.17
3	C	502	8AU	C13-C10-N11	2.18	124.44	119.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PLG	C6-C5-C4	2.12	119.62	118.12
3	C	502	8AU	C6-O1-C2	2.07	120.62	118.38
2	B	501	PLG	C6-C5-C4	2.04	119.56	118.12
2	B	501	PLG	C3-C4-C5	-2.03	116.77	118.72

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PLG	C3-C4-C4A-N
2	A	501	PLG	C5-C4-C4A-N
2	A	501	PLG	C5A-OP4-P-OP1
2	B	501	PLG	C5-C4-C4A-N
2	C	501	PLG	C5-C4-C4A-N
2	C	501	PLG	C5A-OP4-P-OP1
2	C	501	PLG	OXT-C-CA-N
2	B	501	PLG	O-C-CA-N
2	B	501	PLG	C3-C4-C4A-N
2	C	501	PLG	C3-C4-C4A-N
3	A	502	8AU	C21-C16-C4-C5
3	B	502	8AU	C21-C16-C4-C5
2	C	501	PLG	O-C-CA-N
2	A	501	PLG	C5A-OP4-P-OP3
2	B	501	PLG	C5A-OP4-P-OP3
2	C	501	PLG	C5A-OP4-P-OP2
2	B	501	PLG	C4-C4A-N-CA


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	PLG	1	0
2	B	501	PLG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the


average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/442 (100%)	0.61	19 (4%) 35 45	12, 28, 51, 83	0
1	B	442/442 (100%)	0.64	23 (5%) 27 35	13, 28, 51, 70	0
1	C	442/442 (100%)	0.65	21 (4%) 30 39	14, 29, 50, 85	0
All	All	1326/1326 (100%)	0.63	63 (4%) 30 39	12, 28, 51, 85	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	LYS	4.9
1	C	419	LEU	4.2
1	B	363	ASP	4.2
1	A	363	ASP	4.0
1	C	365	VAL	3.9
1	C	363	ASP	3.6
1	C	124	LEU	3.6
1	C	364	CYS	3.5
1	A	366	SER	3.5
1	C	8	GLU	3.3
1	B	365	VAL	3.3
1	A	419	LEU	3.0
1	A	64	TYR	3.0
1	C	413	VAL	2.9
1	C	408	TYR	2.9
1	A	364	CYS	2.9
1	C	193	GLN	2.9
1	B	2	PHE	2.7
1	B	64	TYR	2.7
1	A	413	VAL	2.7
1	B	413	VAL	2.7
1	B	10	ILE	2.7
1	B	364	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	GLY	2.6
1	B	202	ASN	2.5
1	C	202	ASN	2.5
1	B	291	PRO	2.5
1	C	201	VAL	2.5
1	C	417	LYS	2.5
1	C	404	LEU	2.4
1	A	417	LYS	2.4
1	B	9	GLN	2.4
1	A	360	SER	2.4
1	C	15	HIS	2.4
1	B	366	SER	2.4
1	C	88	SER	2.4
1	A	10	ILE	2.4
1	B	138	LYS	2.3
1	B	255	PRO	2.3
1	A	409	GLY	2.3
1	B	198	CYS	2.3
1	B	219	ASN	2.3
1	B	436	ALA	2.3
1	C	412	LEU	2.2
1	C	96	VAL	2.2
1	A	203	ALA	2.2
1	C	29	ILE	2.2
1	C	434	THR	2.2
1	B	68	ASP	2.2
1	A	193	GLN	2.1
1	B	73	ILE	2.1
1	C	221	LEU	2.1
1	A	317	ASP	2.1
1	B	193	GLN	2.1
1	C	438	ALA	2.1
1	B	332	LEU	2.1
1	B	203	ALA	2.1
1	B	408	TYR	2.1
1	A	167	ARG	2.0
1	A	201	VAL	2.0
1	B	412	LEU	2.0
1	A	408	TYR	2.0
1	A	155	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	8AU	B	502	33/33	0.85	0.15	27,30,34,40	0
3	8AU	A	502	33/33	0.86	0.16	25,28,34,41	0
3	8AU	C	502	33/33	0.86	0.14	25,28,31,37	0
4	CL	A	503	1/1	0.87	0.08	31,31,31,31	0
4	CL	C	503	1/1	0.89	0.15	18,18,18,18	1
2	PLG	B	501	20/20	0.91	0.15	14,18,23,23	0
2	PLG	C	501	20/20	0.93	0.14	15,17,21,22	0
2	PLG	A	501	20/20	0.95	0.14	16,18,22,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 8AU B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8AU A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 8AU C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLG B 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PLG C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.