



# wwPDB EM Validation Summary Report ⓘ

Mar 11, 2024 – 02:28 PM JST

PDB ID : 7XOG  
EMDB ID : EMD-33347  
Title : Cryo-EM structure of S glycoprotein encoded by the Covid-19 mRNA vaccine candidate RQ3013 (Postfusion state)  
Authors : Wu, Z.; Yu, Z.; Tan, S.; Lu, J.; Lu, G.; Lin, J.  
Deposited on : 2022-05-01  
Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1293	22% 73%
1	B	1293	23% 73%
1	C	1293	22% 5% 73%
2	D	2	100%
2	H	2	100%
2	I	2	50% 50%
2	M	2	100%
2	N	2	100%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	2	 100%
3	E	4	 25% 75%
3	F	4	 50% 75% 25%
3	J	4	 50% 25% 25%
3	K	4	 25% 100%
3	O	4	 25% 50% 25% 25%
3	P	4	 50% 100%
4	G	3	 33% 67%
4	L	3	 67% 33%
4	Q	3	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	I	1	-	-	X	-
3	MAN	F	4	-	-	X	-
3	NAG	P	1	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 8808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein,peptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	C	348	2688	1683	459	534	12	0	0
1	A	348	2688	1683	459	534	12	0	0
1	B	348	2688	1683	459	534	12	0	0

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	HIS	deletion	UNP P0DTC2
C	?	-	VAL	deletion	UNP P0DTC2
C	?	-	TYR	deletion	UNP P0DTC2
C	414	ASN	LYS	variant	UNP P0DTC2
C	481	LYS	GLU	variant	UNP P0DTC2
C	498	TYR	ASN	variant	UNP P0DTC2
C	567	ASP	ALA	variant	UNP P0DTC2
C	611	GLY	ASP	variant	UNP P0DTC2
C	678	HIS	PRO	variant	UNP P0DTC2
C	679	GLY	ARG	conflict	UNP P0DTC2
C	680	SER	ARG	conflict	UNP P0DTC2
C	682	SER	ARG	conflict	UNP P0DTC2
C	698	VAL	ALA	variant	UNP P0DTC2
A	?	-	HIS	deletion	UNP P0DTC2
A	?	-	VAL	deletion	UNP P0DTC2
A	?	-	TYR	deletion	UNP P0DTC2
A	414	ASN	LYS	variant	UNP P0DTC2
A	481	LYS	GLU	variant	UNP P0DTC2
A	498	TYR	ASN	variant	UNP P0DTC2
A	567	ASP	ALA	variant	UNP P0DTC2
A	611	GLY	ASP	variant	UNP P0DTC2
A	678	HIS	PRO	variant	UNP P0DTC2
A	679	GLY	ARG	conflict	UNP P0DTC2
A	680	SER	ARG	conflict	UNP P0DTC2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	682	SER	ARG	conflict	UNP P0DTC2
A	698	VAL	ALA	variant	UNP P0DTC2
B	?	-	HIS	deletion	UNP P0DTC2
B	?	-	VAL	deletion	UNP P0DTC2
B	?	-	TYR	deletion	UNP P0DTC2
B	414	ASN	LYS	variant	UNP P0DTC2
B	481	LYS	GLU	variant	UNP P0DTC2
B	498	TYR	ASN	variant	UNP P0DTC2
B	567	ASP	ALA	variant	UNP P0DTC2
B	611	GLY	ASP	variant	UNP P0DTC2
B	678	HIS	PRO	variant	UNP P0DTC2
B	679	GLY	ARG	conflict	UNP P0DTC2
B	680	SER	ARG	conflict	UNP P0DTC2
B	682	SER	ARG	conflict	UNP P0DTC2
B	698	VAL	ALA	variant	UNP P0DTC2

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	D	2	28	16	2	10	0	0
2	H	2	28	16	2	10	0	0
2	I	2	28	16	2	10	0	0
2	M	2	28	16	2	10	0	0
2	N	2	28	16	2	10	0	0
2	R	2	28	16	2	10	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
3	E	4	50	28	2	20	0	0
3	F	4	50	28	2	20	0	0
3	J	4	50	28	2	20	0	0
3	K	4	50	28	2	20	0	0
3	O	4	50	28	2	20	0	0
3	P	4	50	28	2	20	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	G	3	39	22	2	15	0	0
4	L	3	39	22	2	15	0	0
4	Q	3	39	22	2	15	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	C	1	Total 14	8	1	5	0
5	C	1	Total 14	8	1	5	0
5	C	1	Total 14	8	1	5	0
5	A	1	Total 14	8	1	5	0
5	A	1	Total 14	8	1	5	0
5	A	1	Total 14	8	1	5	0
5	B	1	Total 14	8	1	5	0
5	B	1	Total 14	8	1	5	0
5	B	1	Total 14	8	1	5	0


- Molecule 6 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			AltConf
6	C	1	Total	C	O	0
			11	6	5	
6	A	1	Total	C	O	0
			11	6	5	
6	B	1	Total	C	O	0
			11	6	5	








● Molecule 1: Spike glycoprotein,peptide







- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  100%

  
MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain R:  100%


  
MAG1  
MAG2

- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  25%  
25% 75%

  
MAG1  
MAG2  
MAN3  
MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50%  
75% 25%

  
MAG1  
MAG2  
MAN3  
MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 25% 25%

  
MAG1  
MAG2  
MAN3  
MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  25%  
100%

  
MAG1  
MAG2  
MAN3  
MAN4


- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	19454	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.25	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.017	Depositor
Minimum map value	-0.808	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.033	Depositor
Recommended contour level	0.3	Depositor
Map size ( $\text{\AA}$ )	563.2, 563.2, 563.2	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.1, 1.1, 1.1	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/2723	0.96	2/3686 (0.1%)
1	B	0.59	0/2723	0.96	2/3686 (0.1%)
1	C	0.59	0/2723	0.96	2/3686 (0.1%)
All	All	0.59	0/8169	0.96	6/11058 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	746	CYS	CA-CB-SG	6.84	126.31	114.00
1	A	746	CYS	CA-CB-SG	6.82	126.28	114.00
1	B	746	CYS	CA-CB-SG	6.81	126.26	114.00
1	A	735	CYS	CA-CB-SG	-5.64	103.85	114.00
1	C	735	CYS	CA-CB-SG	-5.61	103.90	114.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2688	0	2641	114	0
1	B	2688	0	2638	102	0
1	C	2688	0	2638	94	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	28	0	25	0	0
2	H	28	0	25	0	0
2	I	28	0	25	9	0
2	M	28	0	25	0	0
2	N	28	0	25	0	0
2	R	28	0	25	0	0
3	E	50	0	43	1	0
3	F	50	0	43	15	0
3	J	50	0	43	1	0
3	K	50	0	43	1	0
3	O	50	0	43	2	0
3	P	50	0	40	7	0
4	G	39	0	34	3	0
4	L	39	0	34	0	0
4	Q	39	0	34	0	0
5	A	42	0	39	12	0
5	B	42	0	39	6	0
5	C	42	0	39	5	0
6	A	11	0	10	1	0
6	B	11	0	10	2	0
6	C	11	0	10	0	0
All	All	8808	0	8571	230	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 230 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:706:ASN:HD21	5:A:1303:NAG:C1	1.09	1.55
1:A:714:ASN:HD21	5:A:1304:NAG:C1	1.17	1.51
1:B:723:ILE:HD13	3:F:4:MAN:C6	1.44	1.45
1:A:714:ASN:ND2	5:A:1304:NAG:C1	1.72	1.44
1:A:706:ASN:ND2	5:A:1303:NAG:C1	1.80	1.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/1293 (25%)	309 (94%)	19 (6%)	0	100	100
1	B	328/1293 (25%)	309 (94%)	19 (6%)	0	100	100
1	C	328/1293 (25%)	309 (94%)	19 (6%)	0	100	100
All	All	984/3879 (25%)	927 (94%)	57 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/1121 (27%)	301 (98%)	6 (2%)	55	79
1	B	307/1121 (27%)	301 (98%)	6 (2%)	55	79
1	C	307/1121 (27%)	301 (98%)	6 (2%)	55	79
All	All	921/3363 (27%)	903 (98%)	18 (2%)	57	79

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1079	CYS
1	B	1193	SER
1	B	1191	ASN
1	A	1079	CYS
1	B	1071	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	922	ASN
1	B	700	ASN
1	B	1095	ASN
1	B	922	ASN
1	A	700	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

45 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	D	1	2	14,14,15	0.18	0	17,19,21	0.38	0
2	NAG	D	2	2	14,14,15	0.22	0	17,19,21	0.50	0
3	NAG	E	1	3	14,14,15	0.21	0	17,19,21	0.63	0
3	NAG	E	2	3	14,14,15	0.29	0	17,19,21	0.68	0
3	MAN	E	3	3	11,11,12	0.72	0	15,15,17	1.02	2 (13%)
3	MAN	E	4	3	11,11,12	0.74	0	15,15,17	1.22	2 (13%)
3	NAG	F	1	3	14,14,15	0.26	0	17,19,21	0.45	0
3	NAG	F	2	3	14,14,15	0.28	0	17,19,21	0.44	0
3	MAN	F	3	3	11,11,12	0.91	0	15,15,17	1.10	2 (13%)
3	MAN	F	4	3	11,11,12	0.62	0	15,15,17	0.98	2 (13%)
4	NAG	G	1	4	14,14,15	0.27	0	17,19,21	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	2	4	14,14,15	0.25	0	17,19,21	0.37	0
4	MAN	G	3	4	11,11,12	0.93	0	15,15,17	1.07	3 (20%)
2	NAG	H	1	2	14,14,15	0.23	0	17,19,21	0.50	0
2	NAG	H	2	2	14,14,15	0.29	0	17,19,21	0.45	0
2	NAG	I	1	2	14,14,15	0.21	0	17,19,21	0.41	0
2	NAG	I	2	2	14,14,15	0.20	0	17,19,21	0.49	0
3	NAG	J	1	3	14,14,15	0.23	0	17,19,21	0.62	0
3	NAG	J	2	3	14,14,15	0.27	0	17,19,21	0.68	0
3	MAN	J	3	3	11,11,12	0.72	0	15,15,17	1.03	2 (13%)
3	MAN	J	4	3	11,11,12	0.74	0	15,15,17	1.21	2 (13%)
3	NAG	K	1	3	14,14,15	0.25	0	17,19,21	0.46	0
3	NAG	K	2	3	14,14,15	0.26	0	17,19,21	0.44	0
3	MAN	K	3	3	11,11,12	0.93	0	15,15,17	1.08	2 (13%)
3	MAN	K	4	3	11,11,12	0.62	0	15,15,17	0.98	2 (13%)
4	NAG	L	1	4	14,14,15	0.26	0	17,19,21	0.38	0
4	NAG	L	2	4	14,14,15	0.22	0	17,19,21	0.37	0
4	MAN	L	3	4	11,11,12	0.92	0	15,15,17	1.07	3 (20%)
2	NAG	M	1	2	14,14,15	0.23	0	17,19,21	0.48	0
2	NAG	M	2	2	14,14,15	0.29	0	17,19,21	0.42	0
2	NAG	N	1	2	14,14,15	0.18	0	17,19,21	0.39	0
2	NAG	N	2	2	14,14,15	0.19	0	17,19,21	0.51	0
3	NAG	O	1	3	14,14,15	0.20	0	17,19,21	0.62	0
3	NAG	O	2	3	14,14,15	0.30	0	17,19,21	0.67	0
3	MAN	O	3	3	11,11,12	0.70	0	15,15,17	1.03	2 (13%)
3	MAN	O	4	3	11,11,12	0.75	0	15,15,17	1.22	2 (13%)
3	NAG	P	1	3	14,14,15	0.26	0	17,19,21	0.45	0
3	NAG	P	2	3	14,14,15	0.26	0	17,19,21	0.45	0
3	MAN	P	3	3	11,11,12	0.91	0	15,15,17	1.09	2 (13%)
3	MAN	P	4	3	11,11,12	0.61	0	15,15,17	0.98	2 (13%)
4	NAG	Q	1	4	14,14,15	0.28	0	17,19,21	0.37	0
4	NAG	Q	2	4	14,14,15	0.22	0	17,19,21	0.37	0
4	MAN	Q	3	4	11,11,12	0.93	0	15,15,17	1.09	3 (20%)
2	NAG	R	1	2	14,14,15	0.24	0	17,19,21	0.49	0
2	NAG	R	2	2	14,14,15	0.29	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
3	NAG	E	1	3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	MAN	E	3	3	-	0/2/19/22	1/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1
3	NAG	F	1	3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	MAN	F	3	3	-	2/2/19/22	1/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
4	NAG	G	1	4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	MAN	G	3	4	-	2/2/19/22	1/1/1/1
2	NAG	H	1	2	-	2/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	NAG	I	1	2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
3	NAG	J	1	3	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	0/6/23/26	0/1/1/1
3	MAN	J	3	3	-	0/2/19/22	1/1/1/1
3	MAN	J	4	3	-	0/2/19/22	0/1/1/1
3	NAG	K	1	3	-	0/6/23/26	0/1/1/1
3	NAG	K	2	3	-	2/6/23/26	0/1/1/1
3	MAN	K	3	3	-	2/2/19/22	1/1/1/1
3	MAN	K	4	3	-	0/2/19/22	0/1/1/1
4	NAG	L	1	4	-	0/6/23/26	0/1/1/1
4	NAG	L	2	4	-	0/6/23/26	0/1/1/1
4	MAN	L	3	4	-	2/2/19/22	1/1/1/1
2	NAG	M	1	2	-	2/6/23/26	0/1/1/1
2	NAG	M	2	2	-	0/6/23/26	0/1/1/1
2	NAG	N	1	2	-	0/6/23/26	0/1/1/1
2	NAG	N	2	2	-	0/6/23/26	0/1/1/1
3	NAG	O	1	3	-	0/6/23/26	0/1/1/1
3	NAG	O	2	3	-	0/6/23/26	0/1/1/1
3	MAN	O	3	3	-	0/2/19/22	0/1/1/1
3	MAN	O	4	3	-	0/2/19/22	0/1/1/1
3	NAG	P	1	3	-	0/6/23/26	0/1/1/1
3	NAG	P	2	3	-	2/6/23/26	0/1/1/1
3	MAN	P	3	3	-	2/2/19/22	1/1/1/1
3	MAN	P	4	3	-	0/2/19/22	0/1/1/1
4	NAG	Q	1	4	-	0/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	Q	2	4	-	0/6/23/26	0/1/1/1
4	MAN	Q	3	4	-	2/2/19/22	1/1/1/1
2	NAG	R	1	2	-	2/6/23/26	0/1/1/1
2	NAG	R	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	4	MAN	C1-O5-C5	3.39	116.78	112.19
3	O	4	MAN	C1-O5-C5	3.38	116.77	112.19
3	J	4	MAN	C1-O5-C5	3.33	116.70	112.19
3	F	3	MAN	C1-O5-C5	2.60	115.71	112.19
4	L	3	MAN	C1-O5-C5	2.59	115.70	112.19

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	3	MAN	C4-C5-C6-O6
4	L	3	MAN	C4-C5-C6-O6
4	Q	3	MAN	C4-C5-C6-O6
4	G	3	MAN	O5-C5-C6-O6
4	L	3	MAN	O5-C5-C6-O6

5 of 8 ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	3	MAN	C1-C2-C3-C4-C5-O5
3	J	3	MAN	C1-C2-C3-C4-C5-O5
4	Q	3	MAN	C1-C2-C3-C4-C5-O5
4	G	3	MAN	C1-C2-C3-C4-C5-O5
4	L	3	MAN	C1-C2-C3-C4-C5-O5

12 monomers are involved in 39 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	P	1	NAG	7	0
3	F	4	MAN	14	0
3	J	4	MAN	1	0


*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	K	2	NAG	1	0
4	G	1	NAG	3	0
3	P	2	NAG	1	0
3	F	2	NAG	1	0
3	E	2	NAG	1	0
2	I	1	NAG	9	0
3	K	1	NAG	1	0
3	O	4	MAN	2	0
3	F	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

































## 5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	C	1302	1	14,14,15	0.34	0	17,19,21	0.40	0
5	NAG	A	1306	-	14,14,15	0.21	0	17,19,21	0.44	0
6	MAN	B	1306	-	11,11,12	0.80	0	15,15,17	1.46	3 (20%)
5	NAG	C	1305	-	14,14,15	0.20	0	17,19,21	0.46	0
5	NAG	B	1304	1	14,14,15	0.37	0	17,19,21	0.40	0
5	NAG	B	1305	1	14,14,15	0.25	0	17,19,21	0.52	0
6	MAN	A	1305	-	11,11,12	0.80	0	15,15,17	1.45	3 (20%)
5	NAG	C	1303	1	14,14,15	0.27	0	17,19,21	0.52	0
5	NAG	B	1307	-	14,14,15	0.21	0	17,19,21	0.44	0
5	NAG	A	1303	-	14,14,15	0.36	0	17,19,21	0.38	0
6	MAN	C	1304	-	11,11,12	0.79	0	15,15,17	1.46	3 (20%)
5	NAG	A	1304	-	14,14,15	0.27	0	17,19,21	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1302	1	-	2/6/23/26	0/1/1/1
5	NAG	A	1306	-	-	2/6/23/26	0/1/1/1
6	MAN	B	1306	-	-	0/2/19/22	0/1/1/1
5	NAG	C	1305	-	-	2/6/23/26	0/1/1/1
5	NAG	B	1304	1	-	2/6/23/26	0/1/1/1
5	NAG	B	1305	1	-	2/6/23/26	0/1/1/1
6	MAN	A	1305	-	-	0/2/19/22	0/1/1/1
5	NAG	C	1303	1	-	2/6/23/26	0/1/1/1
5	NAG	B	1307	-	-	2/6/23/26	0/1/1/1
5	NAG	A	1303	-	-	2/6/23/26	0/1/1/1

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	C	1304	-	-	0/2/19/22	0/1/1/1
5	NAG	A	1304	-	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
6	B	1306	MAN	C1-O5-C5	3.57	117.03	112.19
6	A	1305	MAN	C1-O5-C5	3.53	116.98	112.19
6	C	1304	MAN	C1-O5-C5	3.50	116.93	112.19
6	C	1304	MAN	O5-C1-C2	2.41	114.49	110.77
6	B	1306	MAN	O5-C1-C2	2.41	114.49	110.77

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1303	NAG	C4-C5-C6-O6
5	A	1304	NAG	C4-C5-C6-O6
5	B	1305	NAG	C4-C5-C6-O6
5	C	1302	NAG	O5-C5-C6-O6
5	A	1303	NAG	O5-C5-C6-O6

There are no ring outliers.

7 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1306	NAG	6	0
6	B	1306	MAN	2	0
5	C	1305	NAG	5	0
6	A	1305	MAN	1	0
5	B	1307	NAG	6	0
5	A	1303	NAG	3	0
5	A	1304	NAG	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	7
1	B	7
1	A	7

The worst 5 of 21 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	727:SER	C	728:MET	N	3.93
1	B	1134:VAL	C	1135:TYR	N	3.83
1	C	1134:VAL	C	1135:TYR	N	3.73
1	A	1134:VAL	C	1135:TYR	N	3.58
1	A	727:SER	C	728:MET	N	3.55


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33347. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)

#### 6.1.1 Primary map




#### 6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 256




Y Index: 256



Z Index: 256

### 6.2.2 Raw map



X Index: 256



Y Index: 256




Z Index: 256

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 252




Y Index: 262




Z Index: 184


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)

### 6.4.1 Primary map




### 6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_33347\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 138 nm<sup>3</sup>; this corresponds to an approximate mass of 125 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.286 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.286 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.49	4.07	3.53
Unmasked-calculated*	8.53	14.90	8.67

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.53 differs from the reported value 3.5 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33347 and PDB model 7XOG. Per-residue inclusion information can be found in section [3](#) on page [9](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).









































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8670	 0.2940
A	 0.8810	 0.3020
B	 0.8820	 0.3050
C	 0.8770	 0.2950
D	 0.8930	 0.3130
E	 0.6600	 0.1880
F	 0.4200	 0.1840
G	 0.7440	 0.2140
H	 0.6070	 0.3040
I	 0.8210	 0.2630
J	 0.7600	 0.1150
K	 0.6600	 0.1970
L	 0.7690	 0.2110
M	 0.6430	 0.2760
N	 0.9290	 0.3060
O	 0.7400	 0.1780
P	 0.4400	 0.1300
Q	 0.7440	 0.2690
R	 0.6070	 0.2020

