

# wwPDB X-ray Structure Validation Summary Report (i)

#### Nov 13, 2023 – 05:00 PM JST

PDB ID	:	5XPY
Title	:	Structural basis of kindlin-mediated integrin recognition and activation
Authors	:	Li, H.; Yang, H.; Sun, K.; Zhang, Z.; Yu, C.; Wei, Z.
Deposited on		
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{l} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	5197(2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	474	90%	6%	·



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Fermitin family homolog 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	454	Total 3730	C 2391	N 626	O 692	S 21	0	8	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	11	HIS	-	expression tag	UNP Q8CIB5
А	12	MET	_	expression tag	UNP Q8CIB5
А	13	GLY	_	expression tag	UNP Q8CIB5
А	14	SER	_	expression tag	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	MET	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	MET	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5

There are 200 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	HIS	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	TRP	deletion	UNP Q8CIB5
А	?	-	PHE	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	_	GLU	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	VAL	deletion	UNP Q8CIB5



Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	PHE	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
A	?	_	PRO	deletion	UNP Q8CIB5
A	?	_	LYS	deletion	UNP Q8CIB5
A	?	-	LYS	deletion	UNP Q8CIB5
A	?	_	LEU	deletion	UNP Q8CIB5
A	?	-	THR	deletion	UNP Q8CIB5
A	?	-	LEU	deletion	UNP Q8CIB5
A	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	GLN	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	TRP	deletion	UNP Q8CIB5
А	?	-	CYS	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	PHE	deletion	UNP Q8CIB5
A	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	CYS	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
A	?	-	SER	deletion	UNP Q8CIB5
A	?	-	ARG	deletion	UNP Q8CIB5
A	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	HIS	deletion	UNP Q8CIB5
А	?	-	GLN	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5



Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	ARG	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	CYS	deletion	UNP Q8CIB5
А	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	VAL	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	VAL	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	GLN	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	PHE	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	VAL	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	MET	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	GLU	deletion	UNP Q8CIB5
A	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	TRP	deletion	UNP Q8CIB5
A	?	-	LEU	deletion	UNP Q8CIB5
A	?	-	ARG	deletion	UNP Q8CIB5
A	?	-	CYS	deletion	UNP Q8CIB5
A	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
A	?	-	GLU	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
A	?	-	GLN	deletion	UNP Q8CIB5
А	?		TYR	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5




5XPY	
------	--

Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	HIS	deletion	UNP Q8CIB5
А	?	-	TRP	deletion	UNP Q8CIB5
А	?	-	MET	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	CYS	deletion	UNP Q8CIB5
А	?	-	ARG	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	GLY	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	-	THR	deletion	UNP Q8CIB5
А	?	-	MET	deletion	UNP Q8CIB5
А	?	-	ALA	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	TYR	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	_	GLU	deletion	UNP Q8CIB5
А	?	_	VAL	deletion	UNP Q8CIB5
А	?	_	GLN	deletion	UNP Q8CIB5
А	?	_	ASN	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	_	LEU	deletion	UNP Q8CIB5
А	?	-	SER	deletion	UNP Q8CIB5
А	?	-	PHE	deletion	UNP Q8CIB5
А	?	_	LEU	deletion	UNP Q8CIB5
А	?	-	LYS	deletion	UNP Q8CIB5
А	?	_	MET	deletion	UNP Q8CIB5
А	?	_	GLN	deletion	UNP Q8CIB5
А	?	_	HIS	deletion	UNP Q8CIB5
А	?	-	LEU	deletion	UNP Q8CIB5
А	?	-	ASN	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
	?	_	ASP	deletion	UNP Q8CIB5
А	•				
A A	?	_	PRO	deletion	UNP Q8CIB5
		-	PRO GLN	deletion deletion	UNP Q8CIB5 UNP Q8CIB5



Chain	Residue	Modelled	Actual	Comment	Reference
А	?	-	ILE	deletion	UNP Q8CIB5
А	?	-	PRO	deletion	UNP Q8CIB5
А	?	-	ASP	deletion	UNP Q8CIB5
А	?	-	GLN	deletion	UNP Q8CIB5
А	?	-	ILE	deletion	UNP Q8CIB5


• Molecule 2 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).





Mol	Chain	Residues	Atom	s	ZeroOcc	AltConf
3	А	1	Total C 6 3	O 3	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	156	Total O 156 156	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fermitin family homolog 2





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants	145.07Å $145.07$ Å $59.35$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	31.41 - 2.10	Depositor
Resolution (A)	31.41 - 2.10	EDS
% Data completeness	93.4 (31.41-2.10)	Depositor
(in resolution range)	84.4 (31.41-2.10)	EDS
R <sub>merge</sub>	0.15	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	7.63 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
D D.	0.164 , $0.196$	Depositor
$R, R_{free}$	0.169 , $0.194$	DCC
$R_{free}$ test set	1756 reflections $(4.77\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	20.1	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36 , $45.4$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3912	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		Bond lengths		angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.40	0/3830	0.50	0/5173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3730	0	3758	16	0
2	А	20	0	15	1	0
3	А	6	0	8	0	0
4	А	156	0	0	1	0
All	All	3912	0	3781	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 16 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:525:ARG:HH11	1:A:529:LYS:NZ	1.82	0.78
1:A:525:ARG:HH11	1:A:529:LYS:HZ2	1.34	0.75
1:A:232:ILE:HG22	1:A:236:MET:HE2	1.70	0.71
1:A:630:ASP:HB2	1:A:632:VAL:HG23	1.73	0.69
1:A:525:ARG:HA	1:A:528:LYS:HD3	1.81	0.62

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	iles
1	А	458/474~(97%)	447 (98%)	11 (2%)	0	100 1	.00

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	413/428~(96%)	407~(98%)	6~(2%)	65 71

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	352[A]	ASP
1	А	352[B]	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	А	630	ASP
1	А	103	LEU
1	А	54	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	$\operatorname{gths}$	Bond angles		
	туре	Ullaill	ries		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	ACT	А	701	-	3,3,3	0.74	0	3,3,3	1.22	0
2	ACT	А	703	-	3,3,3	0.78	0	$3,\!3,\!3$	1.55	0
3	GOL	А	706	-	$5,\!5,\!5$	0.43	0	$5,\!5,\!5$	0.47	0
2	ACT	А	702	-	3,3,3	0.77	0	$3,\!3,\!3$	1.43	0
2	ACT	А	705	-	$3,\!3,\!3$	0.69	0	$3,\!3,\!3$	1.43	0
2	ACT	А	704	-	3,3,3	0.87	0	3,3,3	1.36	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	А	706	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	706	GOL	O1-C1-C2-C3
3	А	706	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Μ	ol	Chain	Res	Type	Clashes	Symm-Clashes
4	2	А	704	ACT	1	0

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	454/474~(95%)	-0.09	19 (4%) 36 42	13, 28, 64, 98	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	333	GLU	4.6
1	А	334	ASN	4.0
1	А	228	THR	3.7
1	А	335	HIS	3.4
1	А	336	LEU	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
3	GOL	А	706	6/6	0.95	0.08	24,29,33,35	0
2	ACT	А	704	4/4	0.96	0.09	32,37,41,45	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	ACT	А	705	4/4	0.98	0.08	$28,\!33,\!35,\!39$	0
2	ACT	А	702	4/4	0.98	0.07	27,33,35,36	0
2	ACT	А	703	4/4	0.99	0.08	19,24,25,32	0
2	ACT	А	701	4/4	0.99	0.08	29,39,40,48	0

Continued from previous page...

## 6.5 Other polymers (i)

There are no such residues in this entry.

