



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2023 – 08:12 PM JST

PDB ID : 7XPN
Title : Structure of the Spring Viraemia of Carp Virus Nucleoprotein
Authors : Wang, Z.X.; Liu, B.; Zhang, Y.A.; Ouyang, S.Y.
Deposited on : 2022-05-04
Resolution : 3.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1039 (4.26-3.70)
Clashscore	141614	1099 (4.26-3.70)
Ramachandran outliers	138981	1061 (4.26-3.70)
Sidechain outliers	138945	1053 (4.26-3.70)
RSRZ outliers	127900	1021 (4.30-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	
1	B	418	
1	C	418	
1	D	418	
1	E	418	
1	F	418	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	418	 4% 68% 32%
1	H	418	 6% 71% 28%
1	I	418	 2% 68% 31%
1	J	418	 6% 67% 32%
1	K	418	 3% 72% 27%
1	L	418	 8% 71% 28%

2 Entry composition [i](#)


There is only 1 type of molecule in this entry. The entry contains 39588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	417	3299	2108	556	618	17	0	0	0
1	E	417	3299	2108	556	618	17	0	0	0
1	F	417	3299	2108	556	618	17	0	0	0
1	I	417	3299	2108	556	618	17	0	0	0
1	K	417	3299	2108	556	618	17	0	0	0
1	B	417	3299	2108	556	618	17	0	0	0
1	C	417	3299	2108	556	618	17	0	0	0
1	G	417	3299	2108	556	618	17	0	0	0
1	H	417	3299	2108	556	618	17	0	0	0
1	D	417	3299	2108	556	618	17	0	0	0
1	J	417	3299	2108	556	618	17	0	0	0
1	L	417	3299	2108	556	618	17	0	0	0


- Molecule 1: Nucleoprotein



- Molecule 1: Nucleoprotein




- Molecule 1: Nucleoprotein








- Molecule 1: Nucleoprotein




- Molecule 1: Nucleoprotein




- Molecule 1: Nucleoprotein






• Molecule 1: Nucleoprotein




• Molecule 1: Nucleoprotein




• Molecule 1: Nucleoprotein





● Molecule 1: Nucleoprotein



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	174.25Å 298.70Å 332.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.86 – 3.98 75.26 – 3.98	Depositor EDS
% Data completeness (in resolution range)	99.9 (72.86-3.98) 91.4 (75.26-3.98)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 4.01Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.255 , 0.284 0.252 , 0.281	Depositor DCC
R_{free} test set	1986 reflections (2.67%)	wwPDB-VP
Wilson B-factor (Å ²)	135.7	Xtrriage
Anisotropy	0.509	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 100.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.016 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.026 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	39588	wwPDB-VP
Average B, all atoms (Å ²)	158.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3375	0.53	0/4575
1	B	0.25	0/3375	0.53	0/4575
1	C	0.25	0/3375	0.52	0/4575
1	D	0.25	0/3375	0.54	0/4575
1	E	0.25	0/3375	0.53	1/4575 (0.0%)
1	F	0.26	0/3375	0.51	0/4575
1	G	0.25	0/3375	0.53	0/4575
1	H	0.25	0/3375	0.51	0/4575
1	I	0.24	0/3375	0.52	0/4575
1	J	0.25	0/3375	0.52	0/4575
1	K	0.25	0/3375	0.52	0/4575
1	L	0.25	0/3375	0.53	1/4575 (0.0%)
All	All	0.25	0/40500	0.52	2/54900 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	J	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	120	LEU	CA-CB-CG	5.81	128.66	115.30
1	L	276	LEU	CA-CB-CG	5.72	128.46	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	125	SER	Peptide
1	J	112	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3299	0	3279	97	0
1	B	3299	0	3279	87	0
1	C	3299	0	3279	83	0
1	D	3299	0	3279	100	0
1	E	3299	0	3279	112	0
1	F	3299	0	3279	90	0
1	G	3299	0	3279	98	0
1	H	3299	0	3279	84	0
1	I	3299	0	3279	88	0
1	J	3299	0	3279	104	0
1	K	3299	0	3279	74	0
1	L	3299	0	3279	83	0
All	All	39588	0	39348	1017	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 1017 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:141:ARG:HB2	1:E:214:GLY:HA2	1.49	0.93
1:J:305:LEU:HA	1:J:397:VAL:HG11	1.56	0.86
1:E:61:ASN:HD21	1:B:166:LYS:HE2	1.39	0.85
1:B:138:GLY:HA2	1:B:214:GLY:HA3	1.58	0.84
1:F:35:GLU:HG2	1:F:107:ARG:HG2	1.60	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/418 (99%)	346 (83%)	64 (15%)	5 (1%)	13	49
1	B	415/418 (99%)	345 (83%)	67 (16%)	3 (1%)	22	60
1	C	415/418 (99%)	352 (85%)	59 (14%)	4 (1%)	15	52
1	D	415/418 (99%)	342 (82%)	65 (16%)	8 (2%)	8	40
1	E	415/418 (99%)	349 (84%)	63 (15%)	3 (1%)	22	60
1	F	415/418 (99%)	350 (84%)	59 (14%)	6 (1%)	11	45
1	G	415/418 (99%)	350 (84%)	64 (15%)	1 (0%)	47	79
1	H	415/418 (99%)	345 (83%)	66 (16%)	4 (1%)	15	52
1	I	415/418 (99%)	355 (86%)	56 (14%)	4 (1%)	15	52
1	J	415/418 (99%)	357 (86%)	56 (14%)	2 (0%)	29	66
1	K	415/418 (99%)	347 (84%)	61 (15%)	7 (2%)	9	42
1	L	415/418 (99%)	346 (83%)	61 (15%)	8 (2%)	8	40
All	All	4980/5016 (99%)	4184 (84%)	741 (15%)	55 (1%)	14	50

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	268	LYS
1	I	365	GLU
1	B	79	ARG
1	D	79	ARG
1	D	126	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	B	356/357 (100%)	356 (100%)	0	100	100
1	C	356/357 (100%)	356 (100%)	0	100	100
1	D	356/357 (100%)	353 (99%)	3 (1%)	81	88
1	E	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	F	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	G	356/357 (100%)	356 (100%)	0	100	100
1	H	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	I	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	J	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	K	356/357 (100%)	355 (100%)	1 (0%)	92	95
1	L	356/357 (100%)	354 (99%)	2 (1%)	86	91
All	All	4272/4284 (100%)	4260 (100%)	12 (0%)	92	95

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	42	ARG
1	D	86	TRP
1	L	201	ASN
1	J	37	ARG
1	I	327	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	104	ASN
1	L	161	GLN
1	I	161	GLN
1	C	161	GLN
1	G	51	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	417/418 (99%)	0.26	11 (2%) 56 46	102, 138, 187, 246	0
1	B	417/418 (99%)	0.39	30 (7%) 15 12	109, 162, 217, 242	0
1	C	417/418 (99%)	0.37	18 (4%) 35 29	111, 146, 199, 249	0
1	D	417/418 (99%)	0.27	21 (5%) 28 25	114, 148, 198, 244	0
1	E	417/418 (99%)	0.28	12 (2%) 51 41	108, 144, 199, 236	0
1	F	417/418 (99%)	0.21	9 (2%) 62 52	102, 145, 204, 231	0
1	G	417/418 (99%)	0.37	18 (4%) 35 29	107, 161, 212, 246	0
1	H	417/418 (99%)	0.43	24 (5%) 23 19	110, 171, 220, 235	0
1	I	417/418 (99%)	0.19	8 (1%) 66 58	106, 139, 191, 236	0
1	J	417/418 (99%)	0.35	25 (5%) 21 17	116, 150, 208, 259	0
1	K	417/418 (99%)	0.28	12 (2%) 51 41	109, 166, 214, 240	0
1	L	417/418 (99%)	0.51	35 (8%) 11 10	112, 177, 228, 245	0
All	All	5004/5016 (99%)	0.33	223 (4%) 33 28	102, 151, 214, 259	0

The worst 5 of 223 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	2	SER	7.7
1	I	362	LEU	7.7
1	L	353	ARG	7.4
1	L	361	TYR	6.6
1	I	363	VAL	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.