



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 9, 2023 – 09:26 PM EDT

PDB ID : 6XQ0  
Title : Human antibody S8V2-18 in complex with the influenza hemagglutinin head domain of A/California/7/2009(NYMC-X181)(H1N1)  
Authors : McCarthy, K.R.; Harrison, S.C.  
Deposited on : 2020-07-09  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	 4% 79% 18%
1	D	225	 2% 80% 16%
2	C	215	 7% 80% 19%
2	F	215	 23% 78% 21%
3	B	240	 7% 78% 16%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	240	
4	G	5	
4	H	5	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10954 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	216	Total 1711	C 1090	N 289	O 326	S 6	0	0	0
1	A	219	Total 1729	C 1101	N 292	O 330	S 6	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	54	ALA	-	expression tag	UNP I6T519
D	55	SER	-	expression tag	UNP I6T519
D	268	LEU	-	expression tag	UNP I6T519
D	269	GLU	-	expression tag	UNP I6T519
D	270	VAL	-	expression tag	UNP I6T519
D	271	LEU	-	expression tag	UNP I6T519
D	272	PHE	-	expression tag	UNP I6T519
D	273	GLN	-	expression tag	UNP I6T519
A	54	ALA	-	expression tag	UNP I6T519
A	55	SER	-	expression tag	UNP I6T519
A	268	LEU	-	expression tag	UNP I6T519
A	269	GLU	-	expression tag	UNP I6T519
A	270	VAL	-	expression tag	UNP I6T519
A	271	LEU	-	expression tag	UNP I6T519
A	272	PHE	-	expression tag	UNP I6T519
A	273	GLN	-	expression tag	UNP I6T519


- Molecule 2 is a protein called antibody S8V2-18 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	213	Total 1645	C 1032	N 281	O 328	S 4	0	0	0
2	F	213	Total 1645	C 1032	N 281	O 328	S 4	0	0	0

- Molecule 3 is a protein called antibody S8V2-18 heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	B	230	Total	C	N	O	S	0	0	0
			1729	1090	302	328	9			
3	E	230	Total	C	N	O	S	0	0	0
			1729	1090	302	328	9			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	5	Total	C	N	O	0	0	0
			59	34	2	23			
4	H	5	Total	C	N	O	0	0	0
			59	34	2	23			

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	132	Total O 132 132	0	0
6	A	127	Total O 127 127	0	0
6	C	86	Total O 86 86	0	0
6	B	114	Total O 114 114	0	0
6	E	114	Total O 114 114	0	0
6	F	69	Total O 69 69	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin




- Molecule 1: Hemagglutinin




- Molecule 2: antibody S8V2-18 light chain





- Molecule 2: antibody S8V2-18 light chain








- Molecule 3: antibody S8V2-18 heavy chain



- Molecule 3: antibody S8V2-18 heavy chain



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	160.20Å 55.26Å 168.00Å 90.00° 95.02° 90.00°	Depositor
Resolution (Å)	44.88 – 2.30 44.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.3 (44.88-2.30) 95.3 (44.88-2.30)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.222 , 0.268 0.221 , 0.264	Depositor DCC
$R_{free}$ test set	3042 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.9	Xtrriage
Anisotropy	0.396	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 32.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10954	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6264e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, GOL, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/1779	0.51	0/2415
1	D	0.31	0/1761	0.53	0/2391
2	C	0.30	0/1679	0.59	0/2278
2	F	0.30	0/1679	0.57	0/2278
3	B	0.31	0/1772	0.62	1/2405 (0.0%)
3	E	0.30	0/1772	0.62	2/2405 (0.1%)
All	All	0.30	0/10442	0.58	3/14172 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	161	TYR	CA-CB-CG	5.20	123.28	113.40
3	E	27	PHE	CB-CG-CD2	-5.17	117.18	120.80
3	B	161	TYR	CA-CB-CG	5.03	122.95	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	94	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1729	0	1658	38	1
1	D	1711	0	1639	28	1
2	C	1645	0	1614	40	0
2	F	1645	0	1614	41	0
3	B	1729	0	1690	37	0
3	E	1729	0	1690	49	0
4	G	59	0	52	3	0
4	H	59	0	52	3	0
5	A	6	0	8	3	0
6	A	127	0	0	9	1
6	B	114	0	0	5	1
6	C	86	0	0	7	0
6	D	132	0	0	5	0
6	E	114	0	0	9	1
6	F	69	0	0	3	0
All	All	10954	0	10017	221	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:170:ASN:O	6:D:301:HOH:O	1.80	0.98
2:C:22:SER:O	6:C:301:HOH:O	1.80	0.97
3:E:57:THR:OG1	6:E:301:HOH:O	1.83	0.95
1:A:102:PHE:HD1	1:A:232:TYR:HB2	1.34	0.92
3:E:204:SER:OG	6:E:303:HOH:O	1.88	0.92

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:503:HOH:O	6:A:515:HOH:O[4_557]	2.06	0.14

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:79:SER:OG	1:D:104:ASP:OD1[4_546]	2.14	0.06
6:B:406:HOH:O	6:E:405:HOH:O[3_455]	2.16	0.04
1:A:79:SER:OG	1:A:104:ASP:OD1[4_547]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/225 (96%)	211 (97%)	6 (3%)	0	100	100
1	D	214/225 (95%)	206 (96%)	8 (4%)	0	100	100
2	C	211/215 (98%)	205 (97%)	6 (3%)	0	100	100
2	F	211/215 (98%)	203 (96%)	8 (4%)	0	100	100
3	B	228/240 (95%)	224 (98%)	4 (2%)	0	100	100
3	E	228/240 (95%)	224 (98%)	3 (1%)	1 (0%)	34	42
All	All	1309/1360 (96%)	1273 (97%)	35 (3%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	146	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/193 (97%)	188 (100%)	0	100	100
1	D	186/193 (96%)	186 (100%)	0	100	100
2	C	184/186 (99%)	182 (99%)	2 (1%)	73	86
2	F	184/186 (99%)	183 (100%)	1 (0%)	88	95
3	B	185/195 (95%)	179 (97%)	6 (3%)	39	54
3	E	185/195 (95%)	182 (98%)	3 (2%)	62	78
All	All	1112/1148 (97%)	1100 (99%)	12 (1%)	73	86

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	161	TYR
3	E	27	PHE
2	F	49	TYR
3	E	97	ARG
3	B	32	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	147	GLN
2	F	53	ASN
2	F	160	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	1	4,1	14,14,15	0.36	0	17,19,21	0.55	0
4	NAG	G	2	4	14,14,15	0.45	0	17,19,21	0.38	0
4	BMA	G	3	4	11,11,12	0.62	0	15,15,17	0.75	0
4	FUC	G	4	4	10,10,11	0.69	0	14,14,16	0.85	0
4	FUC	G	5	4	10,10,11	0.67	1 (10%)	14,14,16	1.53	3 (21%)
4	NAG	H	1	4,1	14,14,15	0.35	0	17,19,21	0.69	1 (5%)
4	NAG	H	2	4	14,14,15	0.28	0	17,19,21	0.41	0
4	BMA	H	3	4	11,11,12	0.55	0	15,15,17	0.72	0
4	FUC	H	4	4	10,10,11	0.65	0	14,14,16	0.86	0
4	FUC	H	5	4	10,10,11	0.86	0	14,14,16	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	FUC	G	4	4	-	-	0/1/1/1
4	FUC	G	5	4	-	-	0/1/1/1
4	NAG	H	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
4	FUC	H	4	4	-	-	0/1/1/1
4	FUC	H	5	4	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	5	FUC	C1-C2	2.01	1.56	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	5	FUC	C1-O5-C5	3.22	120.08	112.78
4	G	5	FUC	O5-C5-C6	-2.20	102.59	107.33
4	H	1	NAG	C1-O5-C5	2.12	115.07	112.19
4	G	5	FUC	O5-C5-C4	2.01	113.12	109.52

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	1	NAG	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	H	1	NAG	O5-C5-C6-O6
4	H	1	NAG	C4-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6

There are no ring outliers.


5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	2	NAG	1	0
4	H	3	BMA	1	0
4	G	4	FUC	1	0
4	H	5	FUC	1	0
4	G	5	FUC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	300	-	5,5,5	0.91	0	5,5,5	1.24	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	300	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	300	GOL	C3-C2-C1	-2.13	103.44	111.70

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	300	GOL	O1-C1-C2-C3
5	A	300	GOL	C1-C2-C3-O3
5	A	300	GOL	O2-C2-C3-O3
5	A	300	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	300	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	219/225 (97%)	0.47	9 (4%) 37 44	24, 34, 55, 76	0
1	D	216/225 (96%)	0.50	5 (2%) 60 67	19, 32, 48, 71	0
2	C	213/215 (99%)	0.62	15 (7%) 16 21	29, 41, 66, 88	0
2	F	213/215 (99%)	1.15	49 (23%) 0 1	31, 56, 89, 108	0
3	B	230/240 (95%)	0.75	16 (6%) 16 21	24, 38, 70, 186	0
3	E	230/240 (95%)	0.92	20 (8%) 10 14	26, 42, 70, 195	0
All	All	1321/1360 (97%)	0.73	114 (8%) 10 14	19, 39, 74, 195	0

The worst 5 of 114 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	149	GLY	24.7
3	B	146	SER	13.9
3	B	147	THR	13.4
3	E	148	SER	10.9
3	E	146	SER	7.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FUC	H	5	10/11	0.69	0.26	53,57,61,64	0
4	BMA	G	3	11/12	0.72	0.23	66,69,73,76	0
4	FUC	G	5	10/11	0.74	0.26	43,48,55,62	0
4	BMA	H	3	11/12	0.78	0.20	61,66,68,69	0
4	FUC	G	4	10/11	0.79	0.17	40,48,50,52	0
4	NAG	G	2	14/15	0.81	0.18	40,49,58,62	0
4	NAG	H	1	14/15	0.85	0.18	24,33,44,53	0
4	FUC	H	4	10/11	0.85	0.22	44,49,52,52	0
4	NAG	H	2	14/15	0.85	0.17	43,50,55,56	0
4	NAG	G	1	14/15	0.88	0.19	27,36,45,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	300	6/6	0.88	0.29	40,53,54,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.