



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 07:21 AM EDT

PDB ID : 1XRN  
Title : Crystal structure of active site F1-mutant E213Q soaked with peptide Phe-Ala  
Authors : Goettig, P.; Brandstetter, H.; Groll, M.; Goehring, W.; Konarev, P.V.; Svergun, D.I.; Huber, R.; Kim, J.-S.  
Deposited on : 2004-10-15  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ALA	A	300	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline iminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	290	2328	1488	386	442	12	17	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	GLN	GLU	engineered mutation	UNP P96084

- Molecule 2 is ALANINE (three-letter code: ALA) (formula:  $C_3H_7NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	6	3	1	2	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	A	12	Total O 12 12	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proline iminopeptidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.33Å 61.51Å 80.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.94 – 2.80 16.85 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (10.94-2.80) 99.0 (16.85-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.50 (at 2.78Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.256 , 0.329 0.254 , 0.325	Depositor DCC
$R_{free}$ test set	408 reflections (5.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.0	Xtrriage
Anisotropy	0.761	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.55 , 109.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	2346	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/2382	0.54	1/3216 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	GLU	N-CA-C	-5.54	96.04	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	2285	93	0
2	A	6	0	4	1	0
3	A	12	0	0	0	0
All	All	2346	0	2289	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ASN:HD21	1:A:169:GLU:HB2	1.40	0.84
1:A:38:PRO:HD2	1:A:209:ASN:HB2	1.64	0.79
1:A:38:PRO:HG3	1:A:106:TYR:CD2	2.23	0.74
1:A:167:ASN:ND2	1:A:169:GLU:HB2	2.06	0.71
1:A:136:LEU:HG	1:A:250:VAL:HG22	1.75	0.68
1:A:30:LYS:HG2	1:A:57:THR:HB	1.77	0.67
1:A:23:LYS:HA	1:A:57:THR:HG23	1.79	0.65
1:A:249:ASN:ND2	1:A:252:ARG:HE	1.95	0.65
1:A:141:MET:HE1	2:A:300:ALA:O	1.97	0.64
1:A:115:ALA:O	1:A:119:GLN:HB3	1.97	0.63
1:A:184:ARG:HG3	1:A:270:SER:HB3	1.79	0.63
1:A:32:MET:HE3	1:A:90:ARG:HG3	1.81	0.62
1:A:126:ILE:HD13	1:A:285:LEU:HG	1.80	0.62
1:A:37:GLY:HA2	1:A:40:MET:HG3	1.82	0.62
1:A:109:ALA:HB1	1:A:223:TRP:CZ3	2.35	0.61
1:A:105:SER:HB2	1:A:271:HIS:NE2	2.16	0.60
1:A:167:ASN:OD1	1:A:168:PRO:HD2	2.01	0.59
1:A:109:ALA:HB2	1:A:131:LEU:HG	1.85	0.58
1:A:38:PRO:HG3	1:A:106:TYR:CG	2.40	0.57
1:A:134:VAL:HG11	1:A:221:LYS:HA	1.86	0.56
1:A:32:MET:HG3	1:A:59:LEU:HD23	1.88	0.56
1:A:35:HIS:HE1	1:A:63:GLN:H	1.52	0.56
1:A:67:GLY:HA2	1:A:198:TYR:CD1	2.41	0.55
1:A:79:ILE:O	1:A:83:VAL:HG23	2.07	0.55
1:A:241:GLY:HA3	1:A:270:SER:O	2.08	0.54
1:A:145:ILE:O	1:A:148:LEU:HB2	2.08	0.54
1:A:144:LEU:O	1:A:148:LEU:HD13	2.09	0.52
1:A:15:ILE:HD13	1:A:72:PRO:HB3	1.92	0.52
1:A:35:HIS:CE1	1:A:63:GLN:H	2.27	0.52
1:A:286:SER:O	1:A:290:LEU:HG	2.09	0.52
1:A:79:ILE:HG23	1:A:110:LEU:HD12	1.91	0.51
1:A:100:PHE:CE2	1:A:289:ILE:HG23	2.46	0.51
1:A:30:LYS:O	1:A:99:VAL:HB	2.10	0.51
1:A:47:SER:HB3	1:A:275:TRP:CH2	2.46	0.51
1:A:15:ILE:HB	1:A:70:GLU:CG	2.41	0.51
1:A:37:GLY:CA	1:A:40:MET:HG3	2.40	0.50
1:A:48:LEU:HA	1:A:282:ASN:HD21	1.76	0.50
1:A:145:ILE:HG23	1:A:156:ILE:HD12	1.94	0.49
1:A:227:ASP:OD1	1:A:228:LYS:HG3	2.12	0.49
1:A:267:ARG:HH11	1:A:267:ARG:HG2	1.77	0.49
1:A:68:ARG:HH21	1:A:68:ARG:HG3	1.76	0.49
1:A:20:LYS:HB3	1:A:60:PHE:HB2	1.94	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:TYR:HA	1:A:162:SER:OG	2.13	0.49
1:A:43:ASP:O	1:A:191:GLU:HB3	2.13	0.48
1:A:137:THR:HG23	1:A:246:VAL:HG13	1.95	0.48
1:A:46:LEU:HD12	1:A:191:GLU:HG2	1.95	0.48
1:A:12:VAL:HG21	1:A:89:LEU:CD1	2.43	0.48
1:A:178:TYR:OH	1:A:212:ASN:HB2	2.14	0.48
1:A:183:LEU:HD12	1:A:272:LEU:HD12	1.96	0.48
1:A:243:TYR:HE2	1:A:268:ASP:HA	1.79	0.47
1:A:63:GLN:HE22	1:A:106:TYR:HE1	1.60	0.47
1:A:76:LYS:HD3	1:A:81:TYR:CE1	2.50	0.47
1:A:27:GLU:HG3	1:A:57:THR:OG1	2.14	0.47
1:A:34:MET:O	1:A:107:GLY:HA3	2.15	0.46
1:A:43:ASP:OD1	1:A:194:LYS:HG2	2.15	0.46
1:A:189:PRO:HG2	1:A:192:VAL:HG23	1.97	0.46
1:A:17:ILE:HD13	1:A:69:SER:HB3	1.97	0.46
1:A:131:LEU:HA	1:A:254:ILE:HD11	1.98	0.46
1:A:207:ILE:O	1:A:219:THR:HG23	2.15	0.46
1:A:119:GLN:HE22	1:A:233:LYS:HB2	1.81	0.46
1:A:38:PRO:O	1:A:205:TYR:HD1	1.99	0.45
1:A:63:GLN:NE2	1:A:106:TYR:HE1	2.13	0.45
1:A:38:PRO:O	1:A:205:TYR:HA	2.17	0.45
1:A:45:LEU:O	1:A:48:LEU:HD13	2.17	0.45
1:A:152:TYR:HE2	1:A:172:GLU:HG2	1.82	0.45
1:A:134:VAL:N	1:A:135:PRO:CD	2.79	0.44
1:A:140:GLU:HB3	1:A:247:THR:CG2	2.47	0.44
1:A:119:GLN:HE22	1:A:233:LYS:H	1.65	0.44
1:A:87:GLU:HA	1:A:87:GLU:OE2	2.17	0.44
1:A:43:ASP:HB2	1:A:195:SER:OG	2.18	0.43
1:A:197:GLU:O	1:A:201:ARG:HG3	2.18	0.43
1:A:28:LYS:HB3	1:A:28:LYS:NZ	2.33	0.43
1:A:31:LEU:HD12	1:A:100:PHE:O	2.18	0.43
1:A:101:LEU:HB2	1:A:122:LEU:CD2	2.47	0.43
1:A:8:ASN:HB3	1:A:19:TYR:CE1	2.53	0.43
1:A:149:PRO:O	1:A:150:ALA:C	2.57	0.43
1:A:15:ILE:HB	1:A:70:GLU:HG3	2.00	0.42
1:A:68:ARG:HG3	1:A:68:ARG:NH2	2.33	0.42
1:A:149:PRO:HG2	1:A:152:TYR:HD1	1.83	0.42
1:A:35:HIS:HE1	1:A:62:ASP:HA	1.84	0.42
1:A:35:HIS:HD2	1:A:36:GLY:O	2.03	0.42
1:A:119:GLN:NE2	1:A:233:LYS:H	2.17	0.42
1:A:36:GLY:HA2	1:A:107:GLY:H	1.84	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:TYR:CZ	1:A:213:GLN:HG2	2.55	0.42
1:A:152:TYR:O	1:A:156:ILE:HG13	2.21	0.41
1:A:100:PHE:CD2	1:A:289:ILE:HG23	2.56	0.41
1:A:189:PRO:HG2	1:A:192:VAL:CG2	2.51	0.41
1:A:267:ARG:O	1:A:268:ASP:HB2	2.21	0.41
1:A:208:MET:O	1:A:220:ILE:HG12	2.19	0.41
1:A:140:GLU:HB3	1:A:247:THR:HG23	2.03	0.41
1:A:174:VAL:HG22	1:A:214:PHE:HZ	1.85	0.41
1:A:22:CYS:O	1:A:57:THR:HG23	2.21	0.41
1:A:179:HIS:CE1	1:A:188:TRP:HE1	2.39	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	288/293 (98%)	247 (86%)	36 (12%)	5 (2%)	<b>9</b> <b>29</b>

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	150	ALA
1	A	211	PRO
1	A	247	THR
1	A	67	GLY
1	A	258	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/255 (98%)	247 (98%)	4 (2%)	62 88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	131	LEU
1	A	185	SER
1	A	230	SER
1	A	293	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	HIS
1	A	63	GLN
1	A	119	GLN
1	A	142	ASN
1	A	167	ASN
1	A	213	GLN
1	A	249	ASN
1	A	255	HIS
1	A	282	ASN
1	A	292	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ALA	A	300	-	2,5,5	0.28	0	2,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALA	A	300	-	-	0/0/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	ALA	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/293 (98%)	0.99	49 (16%) <b>1</b> <b>1</b>	20, 46, 69, 78	5 (1%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	161	SER	8.0
1	A	154	ASP	5.7
1	A	291	LYS	5.4
1	A	175	ASN	5.2
1	A	14	GLY	5.1
1	A	75	SER	4.7
1	A	146	ASP	4.7
1	A	227	ASP	4.5
1	A	162	SER	4.1
1	A	156	ILE	4.0
1	A	212	ASN	3.8
1	A	167	ASN	3.6
1	A	187	ASP	3.3
1	A	268	ASP	3.3
1	A	287	ASP	3.3
1	A	211	PRO	3.2
1	A	220	ILE	3.2
1	A	197	GLU	3.2
1	A	280	GLY	3.1
1	A	150	ALA	3.0
1	A	5	CYS	2.9
1	A	27	GLU	2.9
1	A	80	ASP	2.9
1	A	120	ASP	2.9
1	A	73	ASP	2.8
1	A	221	LYS	2.8
1	A	32	MET	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	293	LEU	2.8
1	A	203	ASN	2.8
1	A	166	GLU	2.7
1	A	186	GLU	2.6
1	A	93	LEU	2.6
1	A	155	ALA	2.6
1	A	96	ASN	2.5
1	A	89	LEU	2.4
1	A	215	THR	2.4
1	A	147	GLU	2.4
1	A	174	VAL	2.4
1	A	111	ALA	2.4
1	A	95	GLY	2.3
1	A	122	LEU	2.3
1	A	4	GLU	2.3
1	A	139	LYS	2.2
1	A	185	SER	2.2
1	A	289	ILE	2.1
1	A	272	LEU	2.1
1	A	216	ILE	2.1
1	A	160	GLY	2.1
1	A	230	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ALA	A	300	6/6	0.65	0.41	47,55,58,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.