



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 6, 2023 – 06:33 PM JST

PDB ID : 7XRE  
Title : Crystal structure of DgpA  
Authors : Ma, W.; He, P.  
Deposited on : 2022-05-10  
Resolution : 2.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	 20% 57% 32% 6% ..
1	B	367	 6% 72% 25% .
1	C	367	 13% 63% 32% ..
1	D	367	 11% 63% 34% ..
1	E	367	 7% 74% 24% ..
1	F	367	 14% 63% 30% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DgpA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	355	2662	1666	463	518	6	9	0	0	0
1	B	357	2702	1695	466	524	7	10	0	0	0
1	C	363	2742	1723	472	530	7	10	0	0	0
1	D	361	2734	1718	469	530	7	10	0	0	0
1	E	360	2717	1704	469	527	7	10	0	0	0
1	F	355	2691	1691	463	520	7	10	0	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A0A3Q9WWX8
A	?	-	GLU	deletion	UNP A0A3Q9WWX8
B	0	GLY	-	expression tag	UNP A0A3Q9WWX8
B	?	-	GLU	deletion	UNP A0A3Q9WWX8
C	0	GLY	-	expression tag	UNP A0A3Q9WWX8
C	?	-	GLU	deletion	UNP A0A3Q9WWX8
D	0	GLY	-	expression tag	UNP A0A3Q9WWX8
D	?	-	GLU	deletion	UNP A0A3Q9WWX8
E	0	GLY	-	expression tag	UNP A0A3Q9WWX8
E	?	-	GLU	deletion	UNP A0A3Q9WWX8
F	0	GLY	-	expression tag	UNP A0A3Q9WWX8
F	?	-	GLU	deletion	UNP A0A3Q9WWX8


- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).








- Molecule 1: DgpA




- Molecule 1: DgpA





- Molecule 1: DgpA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.79Å 130.76Å 206.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.08 – 2.76 26.04 – 2.76	Depositor EDS
% Data completeness (in resolution range)	98.5 (17.08-2.76) 93.9 (26.04-2.76)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.46 (at 2.76Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.235 , 0.294 0.253 , 0.305	Depositor DCC
$R_{free}$ test set	3318 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.3	Xtrriage
Anisotropy	0.174	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	16550	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/2701	0.68	3/3638 (0.1%)
1	B	0.33	1/2741 (0.0%)	0.51	3/3687 (0.1%)
1	C	0.30	0/2784	0.55	0/3750
1	D	0.31	0/2775	0.53	0/3735
1	E	0.28	0/2757	0.48	0/3711
1	F	0.34	1/2731 (0.0%)	0.59	1/3676 (0.0%)
All	All	0.33	2/16489 (0.0%)	0.56	7/22197 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	100	LYS	C-N	8.38	1.50	1.34
1	F	67	ASN	C-N	6.87	1.47	1.34

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	67	ASN	CB-CA-C	-12.64	85.12	110.40
1	B	99	GLU	O-C-N	6.38	132.91	122.70
1	A	57	VAL	C-N-CA	5.64	135.81	121.70
1	A	102	MSE	CG-SE-CE	5.60	111.21	98.90
1	B	99	GLU	CA-C-N	-5.50	105.11	117.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2662	0	2533	170	0
1	B	2702	0	2603	65	0
1	C	2742	0	2650	135	0
1	D	2734	0	2638	93	0
1	E	2717	0	2613	58	0
1	F	2691	0	2599	108	0
2	A	44	0	26	12	0
2	B	44	0	25	6	0
2	C	44	0	25	4	0
2	D	44	0	25	4	0
2	E	44	0	24	4	0
2	F	44	0	26	4	0
3	A	10	0	0	1	0
3	B	11	0	0	1	0
3	C	4	0	0	0	0
3	D	3	0	0	0	0
3	E	4	0	0	1	0
3	F	6	0	0	2	0
All	All	16550	0	15787	605	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 605 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:GLU:HA	1:A:113:VAL:CG1	1.50	1.38
1:C:33:VAL:CG2	1:C:56:GLN:H	1.46	1.29
1:C:100:LYS:HB3	1:C:101:PRO:CD	1.62	1.27
1:A:110:GLU:CA	1:A:113:VAL:CG1	2.18	1.20
1:A:110:GLU:CA	1:A:113:VAL:HG12	1.70	1.18

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/367 (95%)	301 (86%)	30 (9%)	18 (5%)	2	2
1	B	349/367 (95%)	320 (92%)	23 (7%)	6 (2%)	9	16
1	C	359/367 (98%)	311 (87%)	36 (10%)	12 (3%)	4	5
1	D	355/367 (97%)	305 (86%)	38 (11%)	12 (3%)	3	5
1	E	354/367 (96%)	326 (92%)	24 (7%)	4 (1%)	14	25
1	F	349/367 (95%)	311 (89%)	33 (10%)	5 (1%)	11	19
All	All	2115/2202 (96%)	1874 (89%)	184 (9%)	57 (3%)	5	7

5 of 57 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	ILE
1	A	44	GLU
1	A	45	LYS
1	A	58	THR
1	A	100	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/284 (94%)	243 (91%)	23 (9%)	10	18
1	B	276/284 (97%)	272 (99%)	4 (1%)	67	79
1	C	279/284 (98%)	268 (96%)	11 (4%)	32	52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	280/284 (99%)	267 (95%)	13 (5%)	27	46
1	E	276/284 (97%)	269 (98%)	7 (2%)	47	67
1	F	275/284 (97%)	267 (97%)	8 (3%)	42	62
All	All	1652/1704 (97%)	1586 (96%)	66 (4%)	31	51

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	327	GLN
1	F	19	HIS
1	F	365	LYS
1	B	100	LYS
1	B	2	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	333	GLN
1	E	129	ASN
1	E	333	GLN
1	E	196	ASN
1	A	327	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	A	401	-	42,48,48	3.84	19 (45%)	50,73,73	2.15	8 (16%)
2	NAD	B	401	-	42,48,48	4.14	18 (42%)	50,73,73	2.46	12 (24%)
2	NAD	C	401	1	42,48,48	3.89	19 (45%)	50,73,73	2.37	11 (22%)
2	NAD	F	401	-	42,48,48	3.82	19 (45%)	50,73,73	2.30	8 (16%)
2	NAD	D	401	-	42,48,48	3.83	19 (45%)	50,73,73	2.28	9 (18%)
2	NAD	E	401	1	42,48,48	3.87	19 (45%)	50,73,73	2.31	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	401	-	-	14/26/62/62	0/5/5/5
2	NAD	B	401	-	-	16/26/62/62	0/5/5/5
2	NAD	C	401	1	-	10/26/62/62	0/5/5/5
2	NAD	F	401	-	-	11/26/62/62	0/5/5/5
2	NAD	D	401	-	-	11/26/62/62	0/5/5/5
2	NAD	E	401	1	-	9/26/62/62	0/5/5/5

The worst 5 of 113 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAD	C7N-N7N	12.01	1.55	1.33
2	B	401	NAD	O4D-C1D	-10.79	1.26	1.41
2	E	401	NAD	O4D-C1D	-9.97	1.27	1.41
2	A	401	NAD	O4D-C1D	-9.84	1.27	1.41
2	D	401	NAD	O4D-C1D	-9.78	1.27	1.41

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	401	NAD	C1B-N9A-C4A	-9.29	110.32	126.64
2	B	401	NAD	C1B-N9A-C4A	-9.04	110.76	126.64
2	C	401	NAD	C1B-N9A-C4A	-8.57	111.58	126.64
2	E	401	NAD	C1B-N9A-C4A	-8.54	111.64	126.64
2	D	401	NAD	C1B-N9A-C4A	-8.35	111.97	126.64

There are no chirality outliers.

5 of 71 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5B-O5B-PA-O1A
2	A	401	NAD	C5B-O5B-PA-O2A
2	A	401	NAD	PN-O3-PA-O5B
2	A	401	NAD	O4D-C1D-N1N-C2N
2	A	401	NAD	O4D-C1D-N1N-C6N


There are no ring outliers.

6 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAD	12	0
2	B	401	NAD	6	0
2	C	401	NAD	4	0
2	F	401	NAD	4	0
2	D	401	NAD	4	0
2	E	401	NAD	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/367 (94%)	0.97	74 (21%) 0 0	30, 69, 127, 134	0
1	B	347/367 (94%)	0.31	22 (6%) 20 24	39, 62, 86, 118	0
1	C	353/367 (96%)	0.80	48 (13%) 3 3	49, 73, 110, 154	0
1	D	351/367 (95%)	0.65	42 (11%) 4 4	30, 71, 104, 125	0
1	E	350/367 (95%)	0.51	26 (7%) 14 17	48, 67, 92, 126	0
1	F	345/367 (94%)	0.75	51 (14%) 2 2	42, 72, 115, 142	0
All	All	2091/2202 (94%)	0.67	263 (12%) 3 4	30, 69, 115, 154	0

The worst 5 of 263 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	0	GLY	16.4
1	E	88	ALA	9.3
1	A	88	ALA	8.1
1	C	40	ILE	7.8
1	D	2	SER	7.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAD	A	401	44/44	0.74	0.23	73,111,120,122	0
2	NAD	F	401	44/44	0.81	0.23	67,98,108,111	0
2	NAD	C	401	44/44	0.88	0.19	71,90,103,106	0
2	NAD	D	401	44/44	0.92	0.20	64,79,88,90	0
2	NAD	E	401	44/44	0.93	0.17	57,68,74,78	0
2	NAD	B	401	44/44	0.93	0.21	63,71,76,78	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around NAD D 401:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD E 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.