



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 12:29 am BST

PDB ID : 2XSM
Title : Crystal structure of the mammalian cytosolic chaperonin CCT in complex with tubulin
Authors : Munoz, I.G.; Yebenes, H.; Zhou, M.; Mesa, P.; Serna, M.; Bragado-Nilsson, E.; Beloso, A.; Robinson, C.V.; Valpuesta, J.M.; Montoya, G.
Deposited on : 2010-10-29
Resolution : 5.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at


<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

Continued from previous page...

Mol	Chain	Length	Quality of chain
12	L	297	 99%
12	O	297	 99%
13	M	298	 98%
14	N	289	 99%
15	P	481	 99%

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 6438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	489	Total C 489 489	0	0	489

- Molecule 2 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	478	Total C 478 478	0	0	478

- Molecule 3 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	455	Total C 455 455	0	0	455

- Molecule 4 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	D	471	Total C 471 471	0	0	471

- Molecule 5 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	E	472	Total C 472 472	0	0	472

- Molecule 6 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
6	F	466	Total C 466 466	0	0	466

- Molecule 7 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
7	G	485	Total C 485 485	0	0	485

- Molecule 8 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
8	H	474	Total C 474 474	0	0	474

- Molecule 9 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
9	I	293	Total C 293 293	0	0	293

- Molecule 10 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
10	J	299	Total C 299 299	0	0	299

- Molecule 11 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
11	K	394	Total C 394 394	0	0	394

- Molecule 12 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
12	L	297	Total C 297 297	0	0	297
12	O	297	Total C 297 297	0	0	297

- Molecule 13 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
13	M	298	Total C 298 298	0	0	298

- Molecule 14 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
14	N	289	Total C 289 289	0	0	289


- Molecule 15 is a protein called CCT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
15	P	481	Total C 481 481	0	0	481


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCT




- Molecule 2: CCT




- Molecule 3: CCT



- Molecule 4: CCT



- Molecule 5: CCT




- Molecule 6: CCT





- Molecule 7: CCT




- Molecule 8: CCT



- Molecule 9: CCT




- Molecule 10: CCT



- Molecule 11: CCT




- Molecule 12: CCT



- Molecule 12: CCT






- Molecule 13: CCT

Chain M:  98%



- Molecule 14: CCT

Chain N:  99%



- Molecule 15: CCT

Chain P:  99%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	272.70Å 313.50Å 158.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 5.50 97.58 – 5.44	Depositor EDS
% Data completeness (in resolution range)	(Not available) (100.00-5.50) 98.4 (97.58-5.44)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 5.41Å)	Xtriage
Refinement program	REFMAC 5.5.0099	Depositor
R, R_{free}	(Not available) , (Not available) 0.433 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	257.8	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.52 , 11.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.32$, $\langle L^2 \rangle = 0.16$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.76	EDS
Total number of atoms	6438	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	489	0	0	2	0
2	B	478	0	0	5	0
3	C	455	0	0	5	0
4	D	471	0	0	4	0
5	E	472	0	0	11	0
6	F	466	0	0	5	0
7	G	485	0	0	5	0
8	H	474	0	0	2	0
9	I	293	0	0	6	0
10	J	299	0	0	4	0
11	K	394	0	0	8	0
12	L	297	0	0	2	0
12	O	297	0	0	1	0
13	M	298	0	0	3	0
14	N	289	0	0	1	0
15	P	481	0	0	4	0
All	All	6438	0	0	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:K:171:UNK:C	11:K:172:UNK:CA	2.24	1.15
7:G:283:UNK:CA	7:G:284:UNK:CA	2.25	1.13
4:D:448:UNK:CA	4:D:449:UNK:CA	2.27	1.13
3:C:239:UNK:CA	3:C:290:UNK:CA	2.30	1.10
2:B:232:UNK:CA	2:B:346:UNK:CA	2.34	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.