



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 17, 2023 – 01:13 PM EDT

PDB ID : 4XTR  
Title : Structure of Get3 bound to the transmembrane domain of Pep12  
Authors : Mateja, A.; Paduch, M.; Chang, H.-Y.; Szydłowska, A.; Kossiakoff, A.A.; Hegde, R.S.; Keenan, R.J.  
Deposited on : 2015-01-23  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	 6% 82% 5% 14%
1	B	354	 4% 83% 5% 12%
2	C	230	 7% 92% 6% 6%
2	E	230	 96% 2% 2%
3	D	217	 10% 95% 5%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	217	 <p>2% 96%</p>
4	G	37	 <p>46% 46% 14% 41%</p>

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 24293 atoms, of which 11564 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	305	Total	C	H	N	O	S	0	2	0
			4815	1530	2400	401	466	18			
1	B	311	Total	C	H	N	O	S	0	1	0
			4873	1543	2432	404	476	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ASN	ASP	engineered mutation	UNP Q12154
B	57	ASN	ASP	engineered mutation	UNP Q12154

- Molecule 2 is a protein called Antibody Heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	C	217	Total	C	H	N	O	S	0	1	0
			3241	1037	1600	278	320	6			
2	E	222	Total	C	H	N	O	S	0	1	0
			3306	1055	1633	284	328	6			

- Molecule 3 is a protein called Antibody Light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	D	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			
3	F	216	Total	C	H	N	O	S	0	0	0
			3269	1038	1611	276	338	6			


- Molecule 4 is a protein called Pep12p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
4	G	22	Total	C	H	N	O	S	0	0	0
			427	141	231	30	23	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	252	MET	-	initiating methionine	UNP E7M086
G	253	GLY	-	expression tag	UNP E7M086
G	254	SER	-	expression tag	UNP E7M086
G	255	HIS	-	expression tag	UNP E7M086
G	256	HIS	-	expression tag	UNP E7M086
G	257	HIS	-	expression tag	UNP E7M086
G	258	HIS	-	expression tag	UNP E7M086
G	259	HIS	-	expression tag	UNP E7M086
G	260	HIS	-	expression tag	UNP E7M086
G	261	SER	-	expression tag	UNP E7M086

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
5	A	1	38	10	11	5	10	2	0	1
5	B	1	39	10	12	5	10	2	0	1

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		

- Molecule 8 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
8	A	1	42	10	11	5	13	3	0	1
8	B	1	43	10	12	5	13	3	0	1

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	104	Total	O	0	0
			104	104		
9	B	149	Total	O	0	0
			149	149		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	162	Total 162	O 162	0	0
9	D	164	Total 164	O 164	0	0
9	E	211	Total 211	O 211	0	0
9	F	137	Total 137	O 137	0	0
9	G	1	Total 1	O 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATPase GET3




- Molecule 1: ATPase GET3



- Molecule 2: Antibody Heavy chain



- Molecule 2: Antibody Heavy chain




- Molecule 3: Antibody Light chain






- Molecule 3: Antibody Light chain



- Molecule 4: Pep12p



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.87Å 112.03Å 153.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.95 – 2.05 61.56 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.8 (53.95-2.05) 92.1 (61.56-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 2.05Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, $R_{free}$	0.187 , 0.216 0.188 , 0.217	Depositor DCC
$R_{free}$ test set	5469 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtrriage
Anisotropy	0.474	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	24293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, ATP, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/2457	0.40	0/3310
1	B	0.24	0/2482	0.41	0/3345
2	C	0.23	0/1682	0.43	0/2293
2	E	0.23	0/1715	0.44	0/2338
3	D	0.23	0/1694	0.41	0/2299
3	F	0.23	0/1694	0.41	0/2299
4	G	0.24	0/200	0.47	0/269
All	All	0.23	0/11924	0.42	0/16153

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	2400	2397	6	0
1	B	2441	2432	2431	6	0
2	C	1641	1600	1598	4	0
2	E	1673	1633	1632	0	0
3	D	1658	1611	1611	4	0
3	F	1658	1611	1611	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	196	231	230	3	0
5	A	27	11	12	0	0
5	B	27	12	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	1	0	0	0	0
8	A	31	11	12	0	0
8	B	31	12	12	0	0
9	A	104	0	0	0	0
9	B	149	0	0	2	0
9	C	162	0	0	2	0
9	D	164	0	0	0	0
9	E	211	0	0	0	0
9	F	137	0	0	2	0
9	G	1	0	0	0	0
All	All	12729	11564	11558	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:320:GLU:OE1	1:B:322:ARG:NH1	2.01	0.93
3:D:111:ARG:NH1	3:D:112:THR:O	2.10	0.83
1:A:277:ALA:O	1:A:279:ASN:N	2.17	0.77
3:F:192:HIS:O	3:F:214:ARG:NH1	2.24	0.70
1:B:310:HIS:NE2	9:B:577:HOH:O	2.30	0.63
1:B:307:GLU:OE2	9:B:599:HOH:O	2.15	0.63
3:F:141:ASN:ND2	9:F:412:HOH:O	2.31	0.63
3:D:108:GLU:OE2	3:D:176:TYR:OH	2.17	0.56
2:C:107:ARG:NH1	9:C:439:HOH:O	2.37	0.56
1:B:220:LYS:NZ	1:B:224:GLU:OE2	2.44	0.51
1:A:275:LEU:O	1:A:277:ALA:N	2.42	0.50
3:F:25:ARG:NE	3:F:71:ASP:OD2	2.37	0.47
4:G:281:PHE:CZ	4:G:285:ILE:HD11	2.50	0.47
4:G:277:VAL:O	4:G:281:PHE:N	2.49	0.46
3:F:145:ARG:NH1	9:F:413:HOH:O	2.49	0.45
4:G:279:LEU:HA	4:G:282:ILE:HD12	1.99	0.45
1:A:64:ASP:OD1	1:B:254:ARG:NH1	2.48	0.45
3:F:15:SER:N	3:F:18:ASP:OD2	2.39	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246[B]:PHE:CG	1:A:247:LEU:N	2.83	0.43
2:C:90:ARG:NH1	9:C:357:HOH:O	2.43	0.43
2:C:210:LYS:N	2:C:211:PRO:HD2	2.33	0.43
1:A:277:ALA:O	1:A:279:ASN:ND2	2.52	0.42
3:D:135:VAL:HG13	3:D:182:LEU:HB3	2.01	0.42
2:C:107:ARG:NH2	3:D:96:SER:O	2.53	0.41
1:A:246[B]:PHE:CZ	1:B:322:ARG:HB3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/354 (84%)	285 (96%)	10 (3%)	2 (1%)	22	12
1	B	304/354 (86%)	297 (98%)	5 (2%)	2 (1%)	22	12
2	C	214/230 (93%)	208 (97%)	6 (3%)	0	100	100
2	E	221/230 (96%)	218 (99%)	3 (1%)	0	100	100
3	D	214/217 (99%)	209 (98%)	5 (2%)	0	100	100
3	F	214/217 (99%)	210 (98%)	4 (2%)	0	100	100
4	G	20/37 (54%)	19 (95%)	1 (5%)	0	100	100
All	All	1484/1639 (90%)	1446 (97%)	34 (2%)	4 (0%)	41	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	170	THR
1	A	170	THR
1	A	278	GLU
1	B	285	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/309 (88%)	268 (98%)	5 (2%)	59	55
1	B	276/309 (89%)	268 (97%)	8 (3%)	42	35
2	C	182/193 (94%)	180 (99%)	2 (1%)	73	73
2	E	186/193 (96%)	183 (98%)	3 (2%)	62	59
3	D	191/192 (100%)	188 (98%)	3 (2%)	62	59
3	F	191/192 (100%)	191 (100%)	0	100	100
4	G	22/36 (61%)	22 (100%)	0	100	100
All	All	1321/1424 (93%)	1300 (98%)	21 (2%)	65	59

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	LEU
1	A	152	GLN
1	A	216	LEU
1	A	283	HIS
1	A	332	GLN
1	B	98	ASN
1	B	100	MET
1	B	132	SER
1	B	149	ILE
1	B	193	ILE
1	B	194	THR
1	B	228	GLN
1	B	353	LYS
2	C	32	LEU
2	C	206	ASN
3	D	108	GLU
3	D	130	SER
3	D	202	GLN
2	E	32	LEU
2	E	114[A]	GLN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
2	E	114[B]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	ADP	B	401[B]	6	24,29,29	4.72	10 (41%)	29,45,45	2.64	4 (13%)
5	ADP	A	401[B]	6	24,29,29	4.71	10 (41%)	29,45,45	2.65	5 (17%)
8	ATP	A	404[A]	6	26,33,33	2.30	10 (38%)	31,52,52	1.86	9 (29%)
8	ATP	B	403[A]	6	26,33,33	2.28	9 (34%)	31,52,52	1.87	9 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	B	401[B]	6	-	0/12/32/32	0/3/3/3
5	ADP	A	401[B]	6	-	0/12/32/32	0/3/3/3
8	ATP	A	404[A]	6	-	0/18/38/38	0/3/3/3
8	ATP	B	403[A]	6	-	0/18/38/38	0/3/3/3

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	401[B]	ADP	O4'-C1'	14.97	1.62	1.41
5	A	401[B]	ADP	O4'-C1'	14.92	1.61	1.41
5	A	401[B]	ADP	C2'-C1'	-14.56	1.31	1.53
5	B	401[B]	ADP	C2'-C1'	-14.49	1.31	1.53
5	B	401[B]	ADP	O4'-C4'	-5.70	1.32	1.45
5	A	401[B]	ADP	O4'-C4'	-5.68	1.32	1.45
8	B	403[A]	ATP	C2-N1	-5.22	1.24	1.33
8	A	404[A]	ATP	C2-N1	-5.20	1.24	1.33
5	B	401[B]	ADP	C6-N6	4.50	1.50	1.34
5	A	401[B]	ADP	C6-N6	4.50	1.50	1.34
8	B	403[A]	ATP	C4-N3	4.17	1.41	1.35
8	A	404[A]	ATP	C4-N3	4.16	1.41	1.35
8	A	404[A]	ATP	O4'-C1'	4.08	1.46	1.41
8	B	403[A]	ATP	O4'-C1'	3.96	1.46	1.41
8	A	404[A]	ATP	C8-N7	3.95	1.41	1.34
8	B	403[A]	ATP	C8-N7	3.94	1.41	1.34
8	A	404[A]	ATP	C2'-C1'	-3.72	1.48	1.53
8	B	403[A]	ATP	C2'-C1'	-3.71	1.48	1.53
5	A	401[B]	ADP	O2'-C2'	3.28	1.50	1.43
5	B	401[B]	ADP	O2'-C2'	3.27	1.50	1.43
8	A	404[A]	ATP	C5-C4	2.81	1.48	1.40
8	B	403[A]	ATP	C5-C4	2.81	1.48	1.40
8	B	403[A]	ATP	C6-N6	2.67	1.43	1.34
8	A	404[A]	ATP	C6-N6	2.67	1.43	1.34
8	A	404[A]	ATP	C6-N1	-2.64	1.25	1.37
8	B	403[A]	ATP	C6-N1	-2.63	1.25	1.37
5	B	401[B]	ADP	C5-C4	-2.21	1.35	1.40
8	B	403[A]	ATP	C2'-C3'	-2.20	1.47	1.53
5	A	401[B]	ADP	C5-C4	-2.20	1.35	1.40
5	A	401[B]	ADP	C2-N3	2.19	1.35	1.32
5	B	401[B]	ADP	C2-N3	2.17	1.35	1.32
8	A	404[A]	ATP	C2'-C3'	-2.15	1.47	1.53

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	401[B]	ADP	C2-N1	2.13	1.37	1.33
5	B	401[B]	ADP	PA-O5'	2.09	1.67	1.59
5	A	401[B]	ADP	C2-N1	2.06	1.37	1.33
5	A	401[B]	ADP	O3'-C3'	-2.06	1.38	1.43
5	A	401[B]	ADP	PA-O5'	2.06	1.67	1.59
8	A	404[A]	ATP	PG-O2G	-2.01	1.47	1.54
5	B	401[B]	ADP	O3'-C3'	-2.01	1.38	1.43

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401[B]	ADP	C5-C6-N6	10.17	135.80	120.35
5	B	401[B]	ADP	C5-C6-N6	10.16	135.79	120.35
5	A	401[B]	ADP	N6-C6-N1	-6.83	104.39	118.57
5	B	401[B]	ADP	N6-C6-N1	-6.79	104.49	118.57
5	B	401[B]	ADP	N3-C2-N1	-5.66	119.83	128.68
5	A	401[B]	ADP	N3-C2-N1	-5.60	119.92	128.68
8	B	403[A]	ATP	C2-N1-C6	5.45	128.08	118.75
8	A	404[A]	ATP	C2-N1-C6	5.36	127.92	118.75
8	B	403[A]	ATP	N3-C2-N1	-3.85	122.66	128.68
8	A	404[A]	ATP	N3-C2-N1	-3.80	122.73	128.68
8	B	403[A]	ATP	O3G-PG-O3B	3.01	114.73	104.64
8	A	404[A]	ATP	O3G-PG-O3B	2.97	114.60	104.64
8	B	403[A]	ATP	O2G-PG-O3B	2.88	114.29	104.64
8	A	404[A]	ATP	O2G-PG-O3B	2.74	113.83	104.64
8	A	404[A]	ATP	C3'-C2'-C1'	2.74	105.10	100.98
8	B	403[A]	ATP	PB-O3B-PG	-2.69	123.59	132.83
8	A	404[A]	ATP	PB-O3B-PG	-2.63	123.80	132.83
5	B	401[B]	ADP	PA-O3A-PB	-2.57	123.99	132.83
8	B	403[A]	ATP	C3'-C2'-C1'	2.56	104.83	100.98
8	B	403[A]	ATP	PA-O3A-PB	-2.49	124.29	132.83
8	A	404[A]	ATP	PA-O3A-PB	-2.48	124.33	132.83
5	A	401[B]	ADP	PA-O3A-PB	-2.46	124.39	132.83
5	A	401[B]	ADP	C3'-C2'-C1'	2.22	104.32	100.98
8	A	404[A]	ATP	O2B-PB-O1B	-2.21	101.33	112.24
8	A	404[A]	ATP	O2A-PA-O1A	-2.16	101.58	112.24
8	B	403[A]	ATP	O2B-PB-O1B	-2.13	101.73	112.24
8	B	403[A]	ATP	O2A-PA-O1A	-2.09	101.90	112.24


There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/354 (86%)	0.57	23 (7%) 14 15	32, 50, 122, 138	0
1	B	311/354 (87%)	0.44	14 (4%) 33 35	25, 40, 97, 134	0
2	C	217/230 (94%)	0.52	16 (7%) 14 16	24, 42, 97, 131	0
2	E	222/230 (96%)	0.13	0 100 100	22, 35, 58, 102	0
3	D	216/217 (99%)	0.49	21 (9%) 7 8	26, 46, 90, 135	0
3	F	216/217 (99%)	0.22	4 (1%) 66 71	28, 49, 71, 111	0
4	G	22/37 (59%)	3.96	17 (77%) 0 0	90, 112, 124, 143	0
All	All	1509/1639 (92%)	0.46	95 (6%) 20 21	22, 44, 103, 143	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	ILE	10.2
4	G	275	LEU	8.8
1	A	126	LEU	8.2
1	A	190	PHE	7.8
4	G	270	TYR	7.6
4	G	281	PHE	7.0
1	B	190	PHE	6.4
4	G	279	LEU	6.3
4	G	274	VAL	6.2
4	G	271	LEU	6.1
3	D	157	LEU	6.0
3	D	217	CYS	5.9
2	C	223	LYS	5.9
2	C	225	CYS	5.7
4	G	267	TRP	5.6
1	A	193	ILE	5.3
3	D	153	VAL	5.2

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
4	G	280	LEU	5.2
1	A	125	ALA	4.9
2	C	224	SER	4.8
1	B	193	ILE	4.8
1	A	186	LEU	4.7
1	A	100	MET	4.6
1	A	189	LYS	4.6
2	C	137	SER	4.6
3	D	212	PHE	4.4
4	G	277	VAL	4.4
2	C	168	LEU	4.4
4	G	276	LEU	4.2
4	G	278	MET	3.8
3	D	216	GLU	3.7
1	A	246[A]	PHE	3.6
1	B	129	LEU	3.6
3	D	152	LYS	3.5
1	A	129	LEU	3.5
1	A	216	LEU	3.5
1	A	219	LEU	3.5
1	B	284	ASN	3.4
2	C	198	LEU	3.3
1	A	215	LYS	3.3
1	A	124	GLY	3.2
1	A	187	LEU	3.2
4	G	285	ILE	3.2
3	D	215	GLY	3.2
1	B	128	ASP	3.1
2	C	143	GLY	3.1
1	B	100	MET	3.0
3	D	197	CYS	3.0
2	C	210	LYS	2.9
3	D	210	LYS	2.9
3	F	77	SER	2.9
2	C	222	PRO	2.8
3	D	193	LYS	2.8
4	G	284	LEU	2.7
1	B	246	PHE	2.7
1	B	126	LEU	2.7
1	B	212	ILE	2.7
1	B	192	GLU	2.7
1	A	211	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	128	ASP	2.6
3	D	117	SER	2.6
1	B	345	LYS	2.6
3	D	211	SER	2.6
4	G	287	LYS	2.5
2	C	136	SER	2.5
4	G	282	ILE	2.4
3	F	157	LEU	2.4
3	D	195	TYR	2.4
1	A	278	GLU	2.3
2	C	191	VAL	2.3
1	A	284	ASN	2.3
3	D	191	LYS	2.3
3	D	135	VAL	2.3
3	D	194	VAL	2.3
1	B	194	THR	2.3
4	G	269	VAL	2.3
1	A	188	GLU	2.2
4	G	273	ILE	2.2
1	B	101	ALA	2.2
1	B	4	THR	2.2
3	D	113	VAL	2.2
3	D	2	ASP	2.2
2	C	219	LYS	2.2
2	C	130	VAL	2.1
1	A	294	MET	2.1
3	F	129	LYS	2.1
1	A	194	THR	2.1
2	C	172	VAL	2.1
3	F	109	ILE	2.1
2	C	202	THR	2.1
3	D	120	ILE	2.1
3	D	196	ALA	2.1
1	A	153	GLU	2.0
2	C	199	GLY	2.0
3	D	122	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	A	402	1/1	0.94	0.05	41,41,41,41	0
6	MG	B	402	1/1	0.94	0.08	35,35,35,35	0
7	ZN	A	403	1/1	0.94	0.09	68,68,68,68	0
5	ADP	B	401[B]	27/27	0.95	0.15	30,36,45,50	39
8	ATP	A	404[A]	31/31	0.96	0.14	30,36,45,48	42
8	ATP	B	403[A]	31/31	0.96	0.15	25,34,45,52	43
5	ADP	A	401[B]	27/27	0.97	0.14	30,36,46,48	38

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around ADP B 401 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ATP A 404 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ATP B 403 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP A 401 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.