



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 13, 2023 – 06:52 PM JST

PDB ID : 5XT5
Title : SufS-SufU complex from *Bacillus subtilis*
Authors : Fujishiro, T.; Takahashi, Y.
Deposited on : 2017-06-17
Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	155	
1	D	155	
2	A	419	
2	B	419	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8619 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zinc-dependent sulfurtransferase SufU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	137	1050	651	178	209	12	0	0	0
1	C	135	1036	642	175	207	12	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	148	LEU	-	expression tag	UNP O32163
D	149	GLU	-	expression tag	UNP O32163
D	150	HIS	-	expression tag	UNP O32163
D	151	HIS	-	expression tag	UNP O32163
D	152	HIS	-	expression tag	UNP O32163
D	153	HIS	-	expression tag	UNP O32163
D	154	HIS	-	expression tag	UNP O32163
D	155	HIS	-	expression tag	UNP O32163
C	148	LEU	-	expression tag	UNP O32163
C	149	GLU	-	expression tag	UNP O32163
C	150	HIS	-	expression tag	UNP O32163
C	151	HIS	-	expression tag	UNP O32163
C	152	HIS	-	expression tag	UNP O32163
C	153	HIS	-	expression tag	UNP O32163
C	154	HIS	-	expression tag	UNP O32163
C	155	HIS	-	expression tag	UNP O32163

- Molecule 2 is a protein called Cysteine desulfurase SufS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	405	3160	2002	538	606	14	0	1	0
2	B	405	3152	1997	536	606	13	0	0	0


There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP O32164
A	-1	GLY	-	expression tag	UNP O32164
A	0	HIS	-	expression tag	UNP O32164
A	407	VAL	-	expression tag	UNP O32164
A	408	ASP	-	expression tag	UNP O32164
A	409	LEU	-	expression tag	UNP O32164
A	410	GLU	-	expression tag	UNP O32164
A	411	HIS	-	expression tag	UNP O32164
A	412	HIS	-	expression tag	UNP O32164
A	413	HIS	-	expression tag	UNP O32164
A	414	HIS	-	expression tag	UNP O32164
A	415	HIS	-	expression tag	UNP O32164
A	416	HIS	-	expression tag	UNP O32164
B	-2	MET	-	expression tag	UNP O32164
B	-1	GLY	-	expression tag	UNP O32164
B	0	HIS	-	expression tag	UNP O32164
B	407	VAL	-	expression tag	UNP O32164
B	408	ASP	-	expression tag	UNP O32164
B	409	LEU	-	expression tag	UNP O32164
B	410	GLU	-	expression tag	UNP O32164
B	411	HIS	-	expression tag	UNP O32164
B	412	HIS	-	expression tag	UNP O32164
B	413	HIS	-	expression tag	UNP O32164
B	414	HIS	-	expression tag	UNP O32164
B	415	HIS	-	expression tag	UNP O32164
B	416	HIS	-	expression tag	UNP O32164

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Zn 1 1	0	0
3	C	1	Total Zn 1 1	0	0

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	15	8	1	5	1	0	0
4	B	1	15	8	1	5	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	D	3	3	3	0	0
5	A	97	97	97	0	0
5	B	82	82	82	0	0
5	C	7	7	7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Zinc-dependent sulfurtransferase SufU



- Molecule 1: Zinc-dependent sulfurtransferase SufU



- Molecule 2: Cysteine desulfurase SufS



HIS
HIS
HIS
HIS
HIS
HIS

• Molecule 2: Cysteine desulfurase SufS

Chain B: 4% 82% 13%

MET GLY HIS M1 N2 I3 T4 D5 I6 Q16 D21 L22 V23 A29 R35 H53 R54 G55 V56 G69 G91 L101 K109 A123 I142 P143 L144 I150 S151 L152 E153 D154 V155 Q201 K207 S221 S222 H223 K224 V231 E248 L259

Y260 E261 K265 V269 T275 D297 E298 I299 S300 R301 H302 F314 R315 Q316 L317 D318 G319 V320 T321 V322 E327 D337 D338 V339 H340 P341 D349 A350 E351 G352 I353 R356 A357 G358 H359 H360 T372 Y380 L381 E386 E387 I388 V392 E393 A394 L395 Q396

K399 V405 PHE VAL ASP LEU GLU HIS HIS HIS HIS HIS HIS

4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	74.98Å 74.98Å 364.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.93 – 2.34 44.36 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.5 (64.93-2.34) 99.5 (44.36-2.34)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.199 , 0.243 0.198 , 0.243	Depositor DCC
R_{free} test set	2414 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	50.7	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8619	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.86	0/1047	1.10	7/1403 (0.5%)
1	D	0.79	0/1061	1.09	7/1421 (0.5%)
2	A	1.05	3/3229 (0.1%)	0.97	7/4384 (0.2%)
2	B	1.06	5/3217 (0.2%)	1.01	13/4369 (0.3%)
All	All	1.00	8/8554 (0.1%)	1.02	34/11577 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	B	0	1
All	All	0	2

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	261	GLU	CG-CD	9.02	1.65	1.51
2	B	248	GLU	CD-OE2	-6.45	1.18	1.25
2	A	39	GLU	CD-OE1	5.95	1.32	1.25
2	B	393	GLU	CD-OE2	5.88	1.32	1.25
2	A	39	GLU	CD-OE2	5.73	1.31	1.25

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	ARG	NE-CZ-NH2	10.85	125.72	120.30
2	B	315	ARG	NE-CZ-NH1	8.24	124.42	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	156	ARG	NE-CZ-NH1	8.05	124.33	120.30
2	B	356	ARG	NE-CZ-NH2	7.50	124.05	120.30
2	A	154	ASP	CB-CG-OD1	7.48	125.03	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	54	ARG	Peptide
1	C	101	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1036	0	1038	25	0
1	D	1050	0	1056	61	0
2	A	3160	0	3115	29	0
2	B	3152	0	3108	30	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	15	0	6	2	0
4	B	15	0	6	2	0
5	A	97	0	0	2	0
5	B	82	0	0	0	0
5	C	7	0	0	0	0
5	D	3	0	0	0	0
All	All	8619	0	8329	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:27:VAL:HG12	1:D:46:ARG:HH12	1.25	1.01

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:117:THR:CG2	2:A:119:MET:HG2	1.91	1.00
2:B:315:ARG:NH1	2:B:322:VAL:HG11	1.77	0.97
1:D:113:ILE:HD12	1:D:114:GLU:H	1.30	0.96
2:A:117:THR:HG21	2:A:119:MET:HG2	1.50	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	133/155 (86%)	122 (92%)	7 (5%)	4 (3%)	4 2
1	D	135/155 (87%)	114 (84%)	15 (11%)	6 (4%)	2 1
2	A	404/419 (96%)	391 (97%)	12 (3%)	1 (0%)	47 55
2	B	403/419 (96%)	393 (98%)	8 (2%)	2 (0%)	29 31
All	All	1075/1148 (94%)	1020 (95%)	42 (4%)	13 (1%)	13 11

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	114	GLU
2	B	359	HIS
1	C	29	ASN
1	D	30	ASP
1	D	85	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	116/132 (88%)	107 (92%)	9 (8%)	12	12
1	D	117/132 (89%)	105 (90%)	12 (10%)	7	6
2	A	337/349 (97%)	325 (96%)	12 (4%)	35	44
2	B	336/349 (96%)	321 (96%)	15 (4%)	27	34
All	All	906/962 (94%)	858 (95%)	48 (5%)	22	27

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	109	LYS
2	B	380	TYR
2	B	207	LYS
2	B	337	ASP
2	B	396	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	13	GLN
1	D	21	ASN
1	C	117	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PLP	A	501	2	15,15,16	2.72	4 (26%)	20,22,23	1.53	5 (25%)
4	PLP	B	501	2	15,15,16	3.09	3 (20%)	20,22,23	1.75	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PLP	A	501	2	-	1/6/6/8	0/1/1/1
4	PLP	B	501	2	-	1/6/6/8	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	PLP	C3-C2	7.97	1.48	1.40
4	B	501	PLP	C5-C4	7.63	1.48	1.40
4	B	501	PLP	C3-C2	7.43	1.48	1.40
4	A	501	PLP	C5-C4	4.41	1.45	1.40
4	B	501	PLP	C3-C4	3.74	1.47	1.40

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	PLP	C4A-C4-C5	4.73	125.81	120.94
4	A	501	PLP	C6-N1-C2	3.17	125.03	119.17
4	B	501	PLP	C3-C4-C5	-2.80	115.72	118.74
4	A	501	PLP	O3P-P-O4P	-2.48	100.12	106.73
4	B	501	PLP	O3P-P-O2P	2.44	116.95	107.64

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	PLP	C5A-O4P-P-O1P
4	B	501	PLP	C5A-O4P-P-O1P

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	PLP	2	0
4	B	501	PLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	135/155 (87%)	0.80	24 (17%) 1 2	53, 86, 156, 224	0
1	D	137/155 (88%)	1.40	38 (27%) 0 0	78, 112, 154, 182	0
2	A	405/419 (96%)	0.09	18 (4%) 34 45	33, 51, 78, 113	0
2	B	405/419 (96%)	0.19	16 (3%) 38 49	35, 58, 111, 162	0
All	All	1082/1148 (94%)	0.38	96 (8%) 9 15	33, 61, 130, 224	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	27	VAL	6.1
1	D	104	TYR	5.3
1	D	78	GLN	4.4
1	D	50	LYS	4.1
1	D	135	ALA	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	C	201	1/1	0.94	0.07	64,64,64,64	0
3	ZN	D	201	1/1	0.96	0.04	73,73,73,73	0
4	PLP	B	501	15/16	0.98	0.20	41,47,57,61	0
4	PLP	A	501	15/16	0.99	0.19	34,40,43,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.