



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2024 – 01:59 am GMT

PDB ID : 6XTJ
Title : The high resolution structure of the FERM domain of human FERMT2
Authors : Bradshaw, W.J.; Katis, V.L.; Newman, J.A.; von Delft, F.; Arrowsmith, C.H.;
Edwards, A.; Bountra, C.; Gileadi, O.
Deposited on : 2020-01-16
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	478	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fermitin family homolog 2, Fermitin family homolog 2, Fermitin family homolog 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	451	3927	2531	646	726	24	0	35	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	7	SER	-	expression tag	UNP Q96AC1

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	AAA	1	13	6	7	0	0
2	AAA	1	13	6	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			13	6	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	377	Total	O	0	5
			378	378		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fermitin family homolog 2, Fermitin family homolog 2, Fermitin family homolog 2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.17Å 145.17Å 59.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	72.70 – 1.60 72.59 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.70-1.60) 99.7 (72.59-1.60)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.181 , 0.204 0.190 , 0.212	Depositor DCC
R_{free} test set	1874 reflections (1.98%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtrriage
Anisotropy	0.654	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4344	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.64	0/4057	0.75	0/5473

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	362[B]	THR	Mainchain
1	AAA	679[B]	TRP	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3927	0	3978	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	AAA	39	0	15	0	0
3	AAA	378	0	0	5	0
All	All	4344	0	3993	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:538[B]:ARG:HG2	1:AAA:538[B]:ARG:HH21	1.08	1.07
1:AAA:361[B]:LYS:HE2	3:AAA:1110:HOH:O	1.62	0.98
1:AAA:538[B]:ARG:HH21	1:AAA:538[B]:ARG:CG	1.93	0.80
1:AAA:538[B]:ARG:HG2	1:AAA:538[B]:ARG:NH2	1.89	0.79
1:AAA:361[B]:LYS:CE	3:AAA:1110:HOH:O	2.30	0.69
1:AAA:358[B]:GLU:N	1:AAA:358[B]:GLU:OE1	2.27	0.67
1:AAA:680[B]:VAL:CG1	1:AAA:680[B]:VAL:O	2.44	0.65
1:AAA:680[B]:VAL:O	1:AAA:680[B]:VAL:HG13	2.00	0.62
1:AAA:609:PHE:HD1	1:AAA:612[B]:MET:HE3	1.71	0.55
1:AAA:609:PHE:CD1	1:AAA:612[B]:MET:HE3	2.44	0.52
1:AAA:590:TYR:HA	1:AAA:653:TYR:CD1	2.46	0.51
1:AAA:365[A]:ILE:HG23	1:AAA:365[A]:ILE:O	2.12	0.49
1:AAA:142:LYS:HE3	1:AAA:253:LEU:CD2	2.45	0.47
1:AAA:330:MET:HG2	1:AAA:341[A]:LYS:HE3	1.99	0.44
1:AAA:638[B]:CYS:HB2	1:AAA:642:ASP:HB2	1.98	0.44
1:AAA:277:TYR:CG	1:AAA:569:GLY:HA2	2.53	0.43
1:AAA:538[B]:ARG:CG	1:AAA:538[B]:ARG:NH2	2.63	0.43
1:AAA:43:GLY:O	1:AAA:47:LYS:HG3	2.19	0.43
1:AAA:665[B]:GLU:HG3	3:AAA:1097:HOH:O	2.18	0.43
1:AAA:356:THR:HG21	1:AAA:360[B]:GLY:HA3	2.01	0.42
1:AAA:36:THR:HG22	1:AAA:37:GLY:N	2.35	0.41
1:AAA:326:LYS:HG2	3:AAA:1080:HOH:O	2.20	0.41
1:AAA:623[B]:MET:HE3	1:AAA:637:ILE:HD11	2.03	0.41
1:AAA:516:ILE:HD12	1:AAA:540:LEU:HD23	2.03	0.41
1:AAA:627[A]:GLU:HG3	3:AAA:1068:HOH:O	2.20	0.40
1:AAA:365[A]:ILE:O	1:AAA:365[A]:ILE:CG2	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	481/478 (101%)	466 (97%)	15 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	439/431 (102%)	439 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	AAA	701	-	12,12,12	1.54	3 (25%)	17,17,17	1.12	3 (17%)
2	CIT	AAA	702	-	12,12,12	1.16	1 (8%)	17,17,17	1.49	2 (11%)
2	CIT	AAA	703	-	12,12,12	1.20	1 (8%)	17,17,17	1.31	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	AAA	701	-	-	0/16/16/16	-
2	CIT	AAA	702	-	-	1/16/16/16	-
2	CIT	AAA	703	-	-	7/16/16/16	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	701	CIT	C3-C6	3.54	1.57	1.53
2	AAA	703	CIT	C3-C6	2.44	1.55	1.53
2	AAA	702	CIT	C3-C6	2.20	1.55	1.53
2	AAA	701	CIT	O3-C5	2.03	1.28	1.22
2	AAA	701	CIT	O2-C1	-2.02	1.24	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	702	CIT	O5-C6-C3	-4.13	116.40	122.25
2	AAA	703	CIT	O5-C6-C3	-3.48	117.33	122.25
2	AAA	702	CIT	O6-C6-C3	3.17	118.56	113.05
2	AAA	703	CIT	O6-C6-C3	2.54	117.47	113.05
2	AAA	701	CIT	O5-C6-C3	-2.37	118.89	122.25
2	AAA	701	CIT	O7-C3-C6	2.12	111.83	108.86
2	AAA	701	CIT	O6-C6-C3	2.11	116.71	113.05

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	703	CIT	C1-C2-C3-O7
2	AAA	703	CIT	C1-C2-C3-C4
2	AAA	703	CIT	C1-C2-C3-C6
2	AAA	703	CIT	C2-C3-C6-O5
2	AAA	703	CIT	C2-C3-C6-O6
2	AAA	703	CIT	C4-C3-C6-O5
2	AAA	703	CIT	C4-C3-C6-O6
2	AAA	702	CIT	C1-C2-C3-C6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	451/478 (94%)	-0.09	26 (5%) 23 20	14, 25, 66, 91	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	679[A]	TRP	7.1
1	AAA	680[A]	VAL	5.3
1	AAA	238	LYS	5.3
1	AAA	239	PRO	4.9
1	AAA	162	GLU	4.9
1	AAA	228	THR	4.8
1	AAA	144	PRO	3.7
1	AAA	331	THR	3.7
1	AAA	241	ALA	3.5
1	AAA	87	ASP	3.3
1	AAA	18[A]	TRP	3.2
1	AAA	361[A]	LYS	3.2
1	AAA	332	SER	3.1
1	AAA	17	THR	3.0
1	AAA	237	PHE	2.7
1	AAA	240	GLN	2.7
1	AAA	53	ASP	2.7
1	AAA	599	SER	2.7
1	AAA	333	GLU	2.5
1	AAA	678[A]	GLY	2.4
1	AAA	225	GLN	2.4
1	AAA	55	LYS	2.3
1	AAA	86	ALA	2.2
1	AAA	365[A]	ILE	2.2
1	AAA	359[A]	GLY	2.0
1	AAA	52	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIT	AAA	701	13/13	0.77	0.22	32,42,48,53	0
2	CIT	AAA	703	13/13	0.77	0.28	51,75,88,93	0
2	CIT	AAA	702	13/13	0.87	0.25	32,49,54,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.