



Full wwPDB EM Validation Report ⓘ

Oct 12, 2022 – 12:40 AM JST

PDB ID : 7XTD
EMDB ID : EMD-33447
Title : RNA polymerase II elongation complex transcribing a nucleosome (EC58oct)
Authors : Ehara, H.; Kujirai, T.; Shirouzu, M.; Kurumizaka, H.; Sekine, S.
Deposited on : 2022-05-16
Resolution : 3.90 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1743	 80% 19%
2	B	1227	 95% 5%
3	C	304	 87% 13%
4	D	186	 20% 94% 6%
5	E	214	 99%
6	F	155	 54% 46%
7	G	171	 15% 100%
8	H	145	 92% 8%



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	I	115	
10	J	72	
11	K	118	
12	L	72	
13	M	113	
14	N	198	
15	P	19	
16	T	198	
17	V	108	
18	W	911	
19	m	1503	
20	n	417	
21	q	1084	
22	r	544	
23	u	459	
24	v	396	
25	x	395	
26	a	139	
26	e	139	
27	b	106	
27	f	106	
28	c	133	
28	g	133	
29	d	129	
29	h	129	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
30	j	1008	 46% 54%
31	k	531	 81% 79% 19%

2 Entry composition [i](#)

There are 33 unique types of molecules in this entry. The entry contains 81668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1404	11064	6975	1930	2089	70	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1164	9284	5848	1639	1739	58	0	0

- Molecule 3 is a protein called RNA polymerase II third largest subunit B44, part of central core.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	263	2098	1319	354	413	12	0	0

- Molecule 4 is a protein called RNA polymerase II subunit B32.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	174	1349	828	244	274	3	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	213	1741	1094	312	325	10	0	0

- Molecule 6 is a protein called RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	84	677	429	114	131	3	0	0

- Molecule 7 is a protein called RNA polymerase II subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	171	1325	858	214	248	5	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	133	1053	671	169	209	4	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	111	917	565	161	180	11	0	0

- Molecule 10 is a protein called RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	67	554	355	97	96	6	0	0

- Molecule 11 is a protein called RNA polymerase II subunit B12.5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	K	113	932	599	160	169	4	0	0

- Molecule 12 is a protein called RNA polymerase subunit ABC10-alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	45	359	221	72	61	5	0	0

- Molecule 13 is a protein called Transcription elongation factor 1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	M	64	505	318	82	99	6	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-2	GLY	-	expression tag	UNP C4QZ45
M	-1	PRO	-	expression tag	UNP C4QZ45
M	0	GLY	-	expression tag	UNP C4QZ45

- Molecule 14 is a DNA chain called DNA (198-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
14	N	112	2305	1099	389	705	112	0	0

- Molecule 15 is a RNA chain called RNA (5'-R(P*UP*GP*UP*AP*AP*UP*CP*CP*CP*C P*UP*UP*GP*GP*CP*GP*GP*UP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
15	P	19	399	178	64	138	19	0	0

- Molecule 16 is a DNA chain called DNA (198-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
16	T	123	2512	1189	497	703	123	0	0

- Molecule 17 is a protein called Transcription elongation factor SPT4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
17	V	106	824	512	150	155	7	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
V	7	MET	-	initiating methionine	UNP C4R0E6

- Molecule 18 is a protein called Transcription elongation factor SPT5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	W	533	4232	2666	752	812	2	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
W	-2	GLY	-	expression tag	UNP C4R370
W	-1	PRO	-	expression tag	UNP C4R370
W	0	GLY	-	expression tag	UNP C4R370

- Molecule 19 is a protein called Transcription elongation factor Spt6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	m	1187	9730	6162	1663	1877	28	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
m	-2	GLY	-	expression tag	UNP C4R7H2
m	-1	PRO	-	expression tag	UNP C4R7H2
m	0	GLY	-	expression tag	UNP C4R7H2

- Molecule 20 is a protein called Protein that interacts with Spt6p and copurifies with Spt5p and RNA polymerase II.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	n	139	1115	716	193	202	4	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
n	-2	GLY	-	expression tag	UNP C4R7L8
n	-1	PRO	-	expression tag	UNP C4R7L8
n	0	GLY	-	expression tag	UNP C4R7L8

- Molecule 21 is a protein called Component of the Paf1p complex.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	q	930	7552	4805	1283	1439	25	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
q	-39	MET	-	initiating methionine	UNP C4R6B2
q	-38	LYS	-	expression tag	UNP C4R6B2
q	-37	ASP	-	expression tag	UNP C4R6B2
q	-36	HIS	-	expression tag	UNP C4R6B2
q	-35	LEU	-	expression tag	UNP C4R6B2
q	-34	ILE	-	expression tag	UNP C4R6B2
q	-33	HIS	-	expression tag	UNP C4R6B2
q	-32	ASN	-	expression tag	UNP C4R6B2
q	-31	HIS	-	expression tag	UNP C4R6B2
q	-30	HIS	-	expression tag	UNP C4R6B2
q	-29	LYS	-	expression tag	UNP C4R6B2
q	-28	HIS	-	expression tag	UNP C4R6B2
q	-27	GLU	-	expression tag	UNP C4R6B2
q	-26	HIS	-	expression tag	UNP C4R6B2
q	-25	ALA	-	expression tag	UNP C4R6B2
q	-24	HIS	-	expression tag	UNP C4R6B2
q	-23	ALA	-	expression tag	UNP C4R6B2
q	-22	GLU	-	expression tag	UNP C4R6B2
q	-21	HIS	-	expression tag	UNP C4R6B2
q	-20	ASP	-	expression tag	UNP C4R6B2
q	-19	TYR	-	expression tag	UNP C4R6B2
q	-18	LYS	-	expression tag	UNP C4R6B2
q	-17	ASP	-	expression tag	UNP C4R6B2
q	-16	ASP	-	expression tag	UNP C4R6B2
q	-15	ASP	-	expression tag	UNP C4R6B2
q	-14	ASP	-	expression tag	UNP C4R6B2
q	-13	LYS	-	expression tag	UNP C4R6B2
q	-12	GLU	-	expression tag	UNP C4R6B2
q	-11	HIS	-	expression tag	UNP C4R6B2
q	-10	LEU	-	expression tag	UNP C4R6B2
q	-9	TYR	-	expression tag	UNP C4R6B2
q	-8	PHE	-	expression tag	UNP C4R6B2
q	-7	GLN	-	expression tag	UNP C4R6B2
q	-6	GLY	-	expression tag	UNP C4R6B2
q	-5	SER	-	expression tag	UNP C4R6B2
q	-4	SER	-	expression tag	UNP C4R6B2
q	-3	GLY	-	expression tag	UNP C4R6B2
q	-2	SER	-	expression tag	UNP C4R6B2
q	-1	SER	-	expression tag	UNP C4R6B2
q	0	GLY	-	expression tag	UNP C4R6B2

- Molecule 22 is a protein called RNAPII-associated chromatin remodeling Paf1 complex sub-

unit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	r	266	2139	1342	374	412	11	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
r	-29	MET	-	initiating methionine	UNP F2QQ42
r	-28	LYS	-	expression tag	UNP F2QQ42
r	-27	ASP	-	expression tag	UNP F2QQ42
r	-26	HIS	-	expression tag	UNP F2QQ42
r	-25	LEU	-	expression tag	UNP F2QQ42
r	-24	ILE	-	expression tag	UNP F2QQ42
r	-23	HIS	-	expression tag	UNP F2QQ42
r	-22	ASN	-	expression tag	UNP F2QQ42
r	-21	HIS	-	expression tag	UNP F2QQ42
r	-20	HIS	-	expression tag	UNP F2QQ42
r	-19	LYS	-	expression tag	UNP F2QQ42
r	-18	HIS	-	expression tag	UNP F2QQ42
r	-17	GLU	-	expression tag	UNP F2QQ42
r	-16	HIS	-	expression tag	UNP F2QQ42
r	-15	ALA	-	expression tag	UNP F2QQ42
r	-14	HIS	-	expression tag	UNP F2QQ42
r	-13	ALA	-	expression tag	UNP F2QQ42
r	-12	GLU	-	expression tag	UNP F2QQ42
r	-11	HIS	-	expression tag	UNP F2QQ42
r	-10	LEU	-	expression tag	UNP F2QQ42
r	-9	TYR	-	expression tag	UNP F2QQ42
r	-8	PHE	-	expression tag	UNP F2QQ42
r	-7	GLN	-	expression tag	UNP F2QQ42
r	-6	GLY	-	expression tag	UNP F2QQ42
r	-5	SER	-	expression tag	UNP F2QQ42
r	-4	SER	-	expression tag	UNP F2QQ42
r	-3	GLY	-	expression tag	UNP F2QQ42
r	-2	SER	-	expression tag	UNP F2QQ42
r	-1	SER	-	expression tag	UNP F2QQ42
r	0	GLY	-	expression tag	UNP F2QQ42

- Molecule 23 is a protein called Leo1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	u	208	1707	1063	304	337	3	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
u	-29	MET	-	initiating methionine	UNP C4R3K1
u	-28	LYS	-	expression tag	UNP C4R3K1
u	-27	ASP	-	expression tag	UNP C4R3K1
u	-26	HIS	-	expression tag	UNP C4R3K1
u	-25	LEU	-	expression tag	UNP C4R3K1
u	-24	ILE	-	expression tag	UNP C4R3K1
u	-23	HIS	-	expression tag	UNP C4R3K1
u	-22	ASN	-	expression tag	UNP C4R3K1
u	-21	HIS	-	expression tag	UNP C4R3K1
u	-20	HIS	-	expression tag	UNP C4R3K1
u	-19	LYS	-	expression tag	UNP C4R3K1
u	-18	HIS	-	expression tag	UNP C4R3K1
u	-17	GLU	-	expression tag	UNP C4R3K1
u	-16	HIS	-	expression tag	UNP C4R3K1
u	-15	ALA	-	expression tag	UNP C4R3K1
u	-14	HIS	-	expression tag	UNP C4R3K1
u	-13	ALA	-	expression tag	UNP C4R3K1
u	-12	GLU	-	expression tag	UNP C4R3K1
u	-11	HIS	-	expression tag	UNP C4R3K1
u	-10	LEU	-	expression tag	UNP C4R3K1
u	-9	TYR	-	expression tag	UNP C4R3K1
u	-8	PHE	-	expression tag	UNP C4R3K1
u	-7	GLN	-	expression tag	UNP C4R3K1
u	-6	GLY	-	expression tag	UNP C4R3K1
u	-5	SER	-	expression tag	UNP C4R3K1
u	-4	SER	-	expression tag	UNP C4R3K1
u	-3	GLY	-	expression tag	UNP C4R3K1
u	-2	SER	-	expression tag	UNP C4R3K1
u	-1	SER	-	expression tag	UNP C4R3K1
u	0	GLY	-	expression tag	UNP C4R3K1

- Molecule 24 is a protein called RNAP II-associated protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	v	349	2878	1835	510	528	5	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
v	-2	GLY	-	expression tag	UNP C4R997

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
v	-1	SER	-	expression tag	UNP C4R997
v	0	ALA	-	expression tag	UNP C4R997

- Molecule 25 is a protein called Constituent of Paf1 complex with RNA polymerase II, Paf1p, Hpr1p, Ctr9, Leo1, Rtf1 and Ccr4p.

Mol	Chain	Residues	Atoms				AltConf	Trace	
			Total	C	N	O			S
25	x	205	1682	1086	287	307	2	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
x	-29	MET	-	initiating methionine	UNP C4R1E6
x	-28	LYS	-	expression tag	UNP C4R1E6
x	-27	ASP	-	expression tag	UNP C4R1E6
x	-26	HIS	-	expression tag	UNP C4R1E6
x	-25	LEU	-	expression tag	UNP C4R1E6
x	-24	ILE	-	expression tag	UNP C4R1E6
x	-23	HIS	-	expression tag	UNP C4R1E6
x	-22	ASN	-	expression tag	UNP C4R1E6
x	-21	HIS	-	expression tag	UNP C4R1E6
x	-20	HIS	-	expression tag	UNP C4R1E6
x	-19	LYS	-	expression tag	UNP C4R1E6
x	-18	HIS	-	expression tag	UNP C4R1E6
x	-17	GLU	-	expression tag	UNP C4R1E6
x	-16	HIS	-	expression tag	UNP C4R1E6
x	-15	ALA	-	expression tag	UNP C4R1E6
x	-14	HIS	-	expression tag	UNP C4R1E6
x	-13	ALA	-	expression tag	UNP C4R1E6
x	-12	GLU	-	expression tag	UNP C4R1E6
x	-11	HIS	-	expression tag	UNP C4R1E6
x	-10	LEU	-	expression tag	UNP C4R1E6
x	-9	TYR	-	expression tag	UNP C4R1E6
x	-8	PHE	-	expression tag	UNP C4R1E6
x	-7	GLN	-	expression tag	UNP C4R1E6
x	-6	GLY	-	expression tag	UNP C4R1E6
x	-5	SER	-	expression tag	UNP C4R1E6
x	-4	SER	-	expression tag	UNP C4R1E6
x	-3	GLY	-	expression tag	UNP C4R1E6
x	-2	SER	-	expression tag	UNP C4R1E6
x	-1	SER	-	expression tag	UNP C4R1E6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
x	0	GLY	-	expression tag	UNP C4R1E6

- Molecule 26 is a protein called Histone H3.3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
26	a	76	615	390	115	108	2	0	0
26	e	90	735	464	140	129	2	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	-3	GLY	-	expression tag	UNP P84243
a	-2	SER	-	expression tag	UNP P84243
a	-1	HIS	-	expression tag	UNP P84243
e	-3	GLY	-	expression tag	UNP P84243
e	-2	SER	-	expression tag	UNP P84243
e	-1	HIS	-	expression tag	UNP P84243

- Molecule 27 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	b	83	662	418	129	114	1	0	0
27	f	78	619	391	120	107	1	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
b	-3	GLY	-	expression tag	UNP P62805
b	-2	SER	-	expression tag	UNP P62805
b	-1	HIS	-	expression tag	UNP P62805
f	-3	GLY	-	expression tag	UNP P62805
f	-2	SER	-	expression tag	UNP P62805
f	-1	HIS	-	expression tag	UNP P62805

- Molecule 28 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
28	c	103	Total	C	N	O	0	0
			796	502	155	139		
28	g	70	Total	C	N	O	0	0
			544	342	109	93		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
c	-3	GLY	-	expression tag	UNP P04908
c	-2	SER	-	expression tag	UNP P04908
c	-1	HIS	-	expression tag	UNP P04908
g	-3	GLY	-	expression tag	UNP P04908
g	-2	SER	-	expression tag	UNP P04908
g	-1	HIS	-	expression tag	UNP P04908

- Molecule 29 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	d	95	Total	C	N	O	S	0	0
			746	468	136	140	2		
29	h	93	Total	C	N	O	S	0	0
			725	456	130	137	2		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
d	-6	GLY	-	expression tag	UNP P06899
d	-5	SER	-	expression tag	UNP P06899
d	-4	HIS	-	expression tag	UNP P06899
h	-6	GLY	-	expression tag	UNP P06899
h	-5	SER	-	expression tag	UNP P06899
h	-4	HIS	-	expression tag	UNP P06899

- Molecule 30 is a protein called FACT complex subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	j	466	Total	C	N	O	S	0	0
			3780	2401	658	708	13		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
j	-2	GLY	-	expression tag	UNP C4QYQ8
j	-1	PRO	-	expression tag	UNP C4QYQ8
j	0	GLY	-	expression tag	UNP C4QYQ8

- Molecule 31 is a protein called FACT complex subunit POB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
31	k	429	3502	2215	613	664	10	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
k	-2	GLY	-	expression tag	UNP F2QNN8
k	-1	PRO	-	expression tag	UNP F2QNN8
k	0	GLY	-	expression tag	UNP F2QNN8

- Molecule 32 is ZINC ION (three-letter code: ZN) (formula: Zn).



Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
32	A	2	2	2	0
32	B	1	1	1	0
32	C	1	1	1	0
32	I	2	2	2	0
32	J	1	1	1	0
32	L	1	1	1	0
32	M	1	1	1	0
32	V	1	1	1	0

- Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
33	A	1	1	1	0




- Molecule 3: RNA polymerase II third largest subunit B44, part of central core




- Molecule 4: RNA polymerase II subunit B32




- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1



- Molecule 6: RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III




- Molecule 7: RNA polymerase II subunit




- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3






- Molecule 9: DNA-directed RNA polymerase subunit




- Molecule 10: RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III




- Molecule 11: RNA polymerase II subunit B12.5




- Molecule 12: RNA polymerase subunit ABC10-alpha




- Molecule 13: Transcription elongation factor 1 homolog




- Molecule 14: DNA (198-MER)







● Molecule 15: RNA (5'-R(P*UP*GP*UP*AP*AP*UP*CP*CP*CP*CP*UP*UP*GP*GP*CP*GP*GP*UP*U)-3')




● Molecule 16: DNA (198-MER)




● Molecule 17: Transcription elongation factor SPT4



● Molecule 18: Transcription elongation factor SPT5





• Molecule 19: Transcription elongation factor Spt6








Table listing protein residues (amino acids) for EMD-33447, 7XTD. Residues are arranged in vertical columns. Green highlights indicate specific residues. Grey highlights indicate residues that are not visible in the structure.




- Molecule 25: Constituent of Paf1 complex with RNA polymerase II, Paf1p, Hpr1p, Ctr9, Leo1, Rtf1 and Ccr4p





- Molecule 26: Histone H3.3




- Molecule 26: Histone H3.3





• Molecule 27: Histone H4



• Molecule 27: Histone H4



• Molecule 28: Histone H2A type 1-B/E



• Molecule 28: Histone H2A type 1-B/E

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	38828	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.096	Depositor
Minimum map value	-0.052	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.009	Depositor
Map size (\AA)	356.16, 356.16, 356.16	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.484, 1.484, 1.484	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/11267	0.49	0/15222
2	B	0.34	0/9464	0.53	2/12763 (0.0%)
3	C	0.33	0/2139	0.49	0/2895
4	D	0.24	0/1361	0.46	0/1837
5	E	0.30	0/1773	0.50	0/2385
6	F	0.33	0/687	0.50	0/931
7	G	0.28	0/1354	0.48	0/1837
8	H	0.33	0/1070	0.49	0/1444
9	I	0.24	0/934	0.50	0/1257
10	J	0.36	0/563	0.51	0/753
11	K	0.32	0/953	0.50	0/1291
12	L	0.37	0/365	0.62	0/484
13	M	0.25	0/513	0.41	0/693
14	N	1.13	23/2575 (0.9%)	1.32	34/3978 (0.9%)
15	P	0.43	0/443	0.86	2/687 (0.3%)
16	T	1.14	17/2827 (0.6%)	1.29	41/4352 (0.9%)
17	V	0.24	0/840	0.50	0/1140
18	W	0.25	0/4300	0.49	0/5812
19	m	0.24	0/9925	0.45	0/13424
20	n	0.24	0/1132	0.42	0/1526
21	q	0.24	0/7689	0.40	0/10368
22	r	0.30	0/2169	0.53	0/2901
23	u	0.34	0/1740	0.57	2/2347 (0.1%)
24	v	0.25	0/2944	0.47	0/3973
25	x	0.26	0/1716	0.45	0/2310
26	a	0.29	0/622	0.54	0/834
26	e	0.33	0/743	0.61	1/996 (0.1%)
27	b	0.30	0/669	0.59	0/894
27	f	0.34	0/626	0.63	1/837 (0.1%)
28	c	0.29	0/806	0.55	0/1089
28	g	0.34	0/552	0.59	1/743 (0.1%)
29	d	0.30	0/757	0.51	0/1015

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
29	h	0.28	0/736	0.47	0/990
30	j	0.68	0/3855	0.86	4/5191 (0.1%)
31	k	0.68	0/3579	0.88	9/4833 (0.2%)
All	All	0.44	40/83688 (0.0%)	0.63	97/114032 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
14	N	0	5
16	T	0	2
30	j	0	1
31	k	0	2
All	All	0	10

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	N	-40	DC	C1'-N1	6.95	1.58	1.49
16	T	30	DT	C1'-N1	6.67	1.57	1.49
16	T	37	DC	C1'-N1	6.60	1.57	1.49
14	N	-96	DT	C1'-N1	6.50	1.57	1.49
14	N	-27	DC	C1'-N1	6.47	1.57	1.49
16	T	95	DC	C1'-N1	6.45	1.57	1.49
16	T	72	DA	C1'-N9	-6.44	1.38	1.47
14	N	-79	DG	C1'-N9	-6.44	1.38	1.47
14	N	-8	DG	C1'-N9	-6.43	1.38	1.47
16	T	33	DC	C1'-N1	6.41	1.57	1.49
16	T	61	DA	C1'-N9	-6.28	1.38	1.47
16	T	42	DC	C1'-N1	6.24	1.57	1.49
14	N	-10	DG	C1'-N9	-6.22	1.38	1.47
14	N	-7	DG	C1'-N9	-6.18	1.38	1.47
16	T	31	DT	C1'-N1	6.10	1.57	1.49
14	N	-94	DG	C1'-N9	-6.02	1.38	1.47
14	N	-5	DG	C1'-N9	-5.99	1.38	1.47
16	T	48	DG	C1'-N9	-5.96	1.39	1.47
16	T	44	DC	C1'-N1	5.90	1.56	1.49
14	N	-49	DG	C1'-N9	-5.89	1.39	1.47
14	N	-13	DA	C1'-N9	-5.89	1.39	1.47
16	T	80	DA	C1'-N9	-5.88	1.39	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	T	16	DA	C1'-N9	-5.81	1.39	1.47
16	T	77	DG	C1'-N9	-5.71	1.39	1.47
16	T	62	DA	C1'-N9	-5.71	1.39	1.47
14	N	-12	DA	C1'-N9	-5.66	1.39	1.47
14	N	2	DG	C1'-N9	-5.63	1.39	1.47
14	N	4	DG	C1'-N9	-5.59	1.39	1.47
14	N	-6	DG	C1'-N9	-5.51	1.39	1.47
16	T	58	DG	C1'-N9	-5.51	1.39	1.47
14	N	-90	DG	C1'-N9	-5.50	1.39	1.47
16	T	54	DG	C1'-N9	-5.39	1.39	1.47
14	N	-51	DG	C1'-N9	-5.36	1.39	1.47
14	N	-86	DG	C1'-N9	-5.36	1.39	1.47
14	N	6	DA	C1'-N9	-5.24	1.40	1.47
14	N	-14	DA	C1'-N9	-5.21	1.40	1.47
14	N	-4	DG	C1'-N9	-5.11	1.40	1.47
14	N	-3	DA	C1'-N9	-5.11	1.40	1.47
14	N	-44	DG	C1'-N9	-5.06	1.40	1.47
16	T	3	DT	C1'-N1	5.03	1.55	1.49

All (97) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
30	j	684	ARG	NE-CZ-NH1	9.40	125.00	120.30
30	j	836	ARG	NE-CZ-NH1	9.39	125.00	120.30
16	T	105	DA	N1-C6-N6	-9.10	113.14	118.60
16	T	31	DT	O4'-C1'-N1	9.04	114.32	108.00
15	P	10	U	OP1-P-O3'	8.79	124.55	105.20
16	T	110	DA	N1-C6-N6	-8.75	113.35	118.60
16	T	103	DA	N1-C6-N6	-8.57	113.46	118.60
14	N	-107	DA	N1-C6-N6	-8.44	113.54	118.60
16	T	109	DA	N1-C6-N6	-8.39	113.57	118.60
16	T	114	DA	N1-C6-N6	-7.80	113.92	118.60
14	N	-41	DT	O4'-C1'-N1	7.63	113.34	108.00
16	T	28	DG	O4'-C1'-N9	7.56	113.29	108.00
31	k	287	ARG	NE-CZ-NH2	-7.54	116.53	120.30
16	T	102	DA	C5-C6-N1	7.38	121.39	117.70
16	T	102	DA	N1-C6-N6	-7.33	114.20	118.60
16	T	114	DA	C5-C6-N1	7.24	121.32	117.70
16	T	105	DA	C5-C6-N1	7.23	121.32	117.70
16	T	115	DC	N3-C2-O2	-7.22	116.84	121.90
14	N	-107	DA	C5-C6-N1	7.04	121.22	117.70
15	P	11	U	OP1-P-OP2	-7.04	109.04	119.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	N	-113	DC	N3-C2-O2	-7.03	116.98	121.90
16	T	110	DA	C5-C6-N1	7.02	121.21	117.70
16	T	103	DA	C5-C6-N1	6.97	121.19	117.70
16	T	29	DA	O4'-C1'-N9	6.83	112.78	108.00
31	k	287	ARG	NE-CZ-NH1	6.78	123.69	120.30
16	T	101	DC	N3-C2-O2	-6.72	117.20	121.90
14	N	-30	DA	O4'-C1'-N9	6.69	112.68	108.00
16	T	109	DA	C5-C6-N1	6.67	121.04	117.70
16	T	97	DA	O4'-C1'-N9	6.60	112.62	108.00
14	N	-109	DC	O4'-C1'-N1	6.59	112.61	108.00
16	T	105	DA	C4-C5-C6	-6.56	113.72	117.00
16	T	102	DA	C4-C5-C6	-6.51	113.74	117.00
31	k	292	ARG	NE-CZ-NH1	6.49	123.55	120.30
16	T	109	DA	C4-C5-C6	-6.49	113.75	117.00
16	T	39	DA	O4'-C1'-N9	6.44	112.51	108.00
16	T	110	DA	C4-C5-C6	-6.38	113.81	117.00
14	N	-102	DT	O4'-C1'-N1	6.35	112.44	108.00
16	T	111	DT	O4'-C1'-N1	6.34	112.44	108.00
16	T	99	DC	O4'-C1'-N1	6.34	112.44	108.00
14	N	-105	DC	N3-C2-O2	-6.32	117.48	121.90
14	N	-107	DA	C4-C5-C6	-6.31	113.85	117.00
16	T	103	DA	C4-C5-C6	-6.24	113.88	117.00
16	T	114	DA	C4-C5-C6	-6.23	113.89	117.00
26	e	116	ARG	NE-CZ-NH1	6.12	123.36	120.30
14	N	-104	DT	C6-C5-C7	-6.07	119.26	122.90
16	T	111	DT	C6-C5-C7	-6.00	119.30	122.90
16	T	107	DC	N3-C2-O2	-5.99	117.71	121.90
16	T	108	DT	C6-C5-C7	-5.92	119.35	122.90
14	N	-106	DT	C6-C5-C7	-5.90	119.36	122.90
27	f	45	ARG	NE-CZ-NH1	5.89	123.25	120.30
16	T	112	DT	P-O3'-C3'	5.83	126.69	119.70
23	u	273	ARG	NE-CZ-NH1	5.82	123.21	120.30
14	N	-97	DT	O4'-C1'-N1	5.82	112.07	108.00
31	k	302	ARG	NE-CZ-NH1	5.80	123.20	120.30
14	N	-103	DT	C6-C5-C7	-5.76	119.44	122.90
16	T	112	DT	C6-C5-C7	-5.72	119.47	122.90
14	N	-100	DT	N3-C2-O2	-5.67	118.90	122.30
2	B	433	ARG	NE-CZ-NH1	5.67	123.13	120.30
14	N	-113	DC	N1-C2-O2	5.66	122.30	118.90
16	T	104	DG	N1-C6-O6	-5.65	116.51	119.90
14	N	-111	DG	N1-C6-O6	-5.65	116.51	119.90
14	N	-114	DT	C6-C5-C7	-5.62	119.53	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	N	-112	DT	C6-C5-C7	-5.58	119.55	122.90
16	T	103	DA	O4'-C1'-N9	5.55	111.89	108.00
14	N	-104	DT	N3-C2-O2	-5.55	118.97	122.30
14	N	-103	DT	N3-C2-O2	-5.55	118.97	122.30
16	T	112	DT	N3-C2-O2	-5.53	118.98	122.30
31	k	264	ARG	NE-CZ-NH1	5.51	123.06	120.30
16	T	115	DC	N1-C2-O2	5.50	122.20	118.90
14	N	-110	DG	N1-C6-O6	-5.49	116.61	119.90
14	N	-108	DC	N3-C2-O2	-5.47	118.07	121.90
31	k	58	ARG	NE-CZ-NH1	5.38	122.99	120.30
31	k	270	ARG	NE-CZ-NH1	5.36	122.98	120.30
16	T	111	DT	N3-C2-O2	-5.33	119.10	122.30
30	j	817	ARG	NE-CZ-NH1	5.30	122.95	120.30
2	B	427	ARG	NE-CZ-NH1	5.29	122.94	120.30
23	u	271	ARG	NE-CZ-NH1	5.28	122.94	120.30
14	N	-109	DC	N3-C2-O2	-5.27	118.21	121.90
14	N	-112	DT	N3-C2-O2	-5.26	119.14	122.30
14	N	-106	DT	N3-C2-O2	-5.26	119.14	122.30
16	T	106	DG	C5-C6-N1	5.23	114.12	111.50
16	T	94	DC	P-O3'-C3'	5.22	125.97	119.70
14	N	-100	DT	C6-C5-C7	-5.22	119.77	122.90
30	j	616	ARG	NE-CZ-NH1	5.21	122.91	120.30
28	g	77	ARG	NE-CZ-NH1	5.21	122.90	120.30
14	N	-81	DG	C3'-C2'-C1'	-5.16	96.31	102.50
14	N	-102	DT	C6-C5-C7	-5.14	119.82	122.90
14	N	-44	DG	O4'-C1'-N9	5.11	111.58	108.00
31	k	129	ARG	NE-CZ-NH1	5.09	122.85	120.30
14	N	-105	DC	N1-C2-O2	5.09	121.95	118.90
14	N	-106	DT	O4'-C1'-N1	5.08	111.56	108.00
14	N	-43	DT	O4'-C1'-N1	5.05	111.54	108.00
16	T	105	DA	O4'-C1'-N9	5.04	111.53	108.00
31	k	5	ASP	CB-CG-OD2	-5.03	113.78	118.30
14	N	-109	DC	C1'-O4'-C4'	-5.02	105.08	110.10
14	N	-108	DC	N3-C4-C5	5.02	123.91	121.90
16	T	31	DT	C1'-O4'-C4'	-5.01	105.09	110.10

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
14	N	-101	DG	Sidechain
14	N	-102	DT	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
14	N	-108	DC	Sidechain
14	N	-111	DG	Sidechain
14	N	-114	DT	Sidechain
16	T	103	DA	Sidechain
16	T	112	DT	Sidechain
30	j	616	ARG	Sidechain
31	k	292	ARG	Sidechain
31	k	58	ARG	Sidechain

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1392/1743 (80%)	1354 (97%)	37 (3%)	1 (0%)	51	84
2	B	1154/1227 (94%)	1117 (97%)	37 (3%)	0	100	100
3	C	261/304 (86%)	259 (99%)	2 (1%)	0	100	100
4	D	170/186 (91%)	166 (98%)	4 (2%)	0	100	100
5	E	211/214 (99%)	205 (97%)	6 (3%)	0	100	100
6	F	82/155 (53%)	80 (98%)	2 (2%)	0	100	100
7	G	169/171 (99%)	166 (98%)	3 (2%)	0	100	100
8	H	129/145 (89%)	125 (97%)	4 (3%)	0	100	100
9	I	109/115 (95%)	107 (98%)	2 (2%)	0	100	100
10	J	65/72 (90%)	65 (100%)	0	0	100	100
11	K	111/118 (94%)	110 (99%)	1 (1%)	0	100	100
12	L	43/72 (60%)	41 (95%)	2 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
13	M	62/113 (55%)	62 (100%)	0	0	100	100
17	V	104/108 (96%)	100 (96%)	4 (4%)	0	100	100
18	W	527/911 (58%)	506 (96%)	21 (4%)	0	100	100
19	m	1179/1503 (78%)	1157 (98%)	22 (2%)	0	100	100
20	n	137/417 (33%)	136 (99%)	1 (1%)	0	100	100
21	q	928/1084 (86%)	922 (99%)	6 (1%)	0	100	100
22	r	260/544 (48%)	254 (98%)	6 (2%)	0	100	100
23	u	206/459 (45%)	204 (99%)	2 (1%)	0	100	100
24	v	341/396 (86%)	327 (96%)	14 (4%)	0	100	100
25	x	201/395 (51%)	200 (100%)	1 (0%)	0	100	100
26	a	74/139 (53%)	70 (95%)	4 (5%)	0	100	100
26	e	88/139 (63%)	87 (99%)	1 (1%)	0	100	100
27	b	81/106 (76%)	78 (96%)	3 (4%)	0	100	100
27	f	76/106 (72%)	73 (96%)	3 (4%)	0	100	100
28	c	101/133 (76%)	100 (99%)	1 (1%)	0	100	100
28	g	68/133 (51%)	66 (97%)	2 (3%)	0	100	100
29	d	93/129 (72%)	89 (96%)	4 (4%)	0	100	100
29	h	91/129 (70%)	90 (99%)	1 (1%)	0	100	100
30	j	462/1008 (46%)	449 (97%)	12 (3%)	1 (0%)	47	79
31	k	423/531 (80%)	411 (97%)	12 (3%)	0	100	100
All	All	9398/13005 (72%)	9176 (98%)	220 (2%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	960	VAL
30	j	543	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1219/1528 (80%)	1216 (100%)	3 (0%)	93	96
2	B	1018/1077 (94%)	1016 (100%)	2 (0%)	93	96
3	C	236/264 (89%)	236 (100%)	0	100	100
4	D	149/160 (93%)	149 (100%)	0	100	100
5	E	196/197 (100%)	194 (99%)	2 (1%)	76	86
6	F	75/137 (55%)	75 (100%)	0	100	100
7	G	148/148 (100%)	148 (100%)	0	100	100
8	H	120/130 (92%)	120 (100%)	0	100	100
9	I	106/109 (97%)	106 (100%)	0	100	100
10	J	61/66 (92%)	61 (100%)	0	100	100
11	K	104/109 (95%)	104 (100%)	0	100	100
12	L	38/56 (68%)	38 (100%)	0	100	100
13	M	61/99 (62%)	60 (98%)	1 (2%)	62	79
17	V	90/92 (98%)	90 (100%)	0	100	100
18	W	480/796 (60%)	480 (100%)	0	100	100
19	m	1087/1354 (80%)	1086 (100%)	1 (0%)	93	97
20	n	125/361 (35%)	125 (100%)	0	100	100
21	q	824/962 (86%)	824 (100%)	0	100	100
22	r	239/485 (49%)	239 (100%)	0	100	100
23	u	192/406 (47%)	191 (100%)	1 (0%)	88	93
24	v	325/369 (88%)	325 (100%)	0	100	100
25	x	190/354 (54%)	190 (100%)	0	100	100
26	a	64/112 (57%)	64 (100%)	0	100	100
26	e	77/112 (69%)	77 (100%)	0	100	100
27	b	68/81 (84%)	68 (100%)	0	100	100
27	f	63/81 (78%)	63 (100%)	0	100	100
28	c	82/102 (80%)	82 (100%)	0	100	100
28	g	54/102 (53%)	54 (100%)	0	100	100
29	d	81/107 (76%)	81 (100%)	0	100	100
29	h	79/107 (74%)	79 (100%)	0	100	100
30	j	417/910 (46%)	417 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
31	k	392/474 (83%)	392 (100%)	0	100	100
All	All	8460/11447 (74%)	8450 (100%)	10 (0%)	93	97

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	408	ARG
1	A	831	LYS
1	A	1302	LYS
2	B	463	LYS
2	B	904	ARG
5	E	36	GLN
5	E	166	ARG
13	M	40	LYS
19	m	514	LYS
23	u	124	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	737	ASN
8	H	44	ASN
9	I	11	ASN
18	W	644	ASN
19	m	955	ASN
20	n	232	GLN
21	q	147	ASN
25	x	296	ASN
27	b	25	ASN
28	c	73	ASN
30	j	750	GLN
31	k	127	ASN
31	k	141	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
15	P	19/19 (100%)	10 (52%)	1 (5%)

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
15	P	-6	G
15	P	-5	U
15	P	-4	A
15	P	-3	A
15	P	-2	U
15	P	0	C
15	P	1	C
15	P	8	G
15	P	9	G
15	P	11	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
15	P	-7	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33447. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 120




Y Index: 120




Z Index: 120


6.2.2 Raw map



X Index: 120



Y Index: 120




Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 141




Y Index: 125




Z Index: 137


6.3.2 Raw map



X Index: 141



Y Index: 121




Z Index: 146

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.009. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 526 nm^3 ; this corresponds to an approximate mass of 475 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.90	-	-
Author-provided FSC curve	3.85	4.64	3.91
Unmasked-calculated*	4.19	7.59	4.34

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33447 and PDB model 7XTD. Per-residue inclusion information can be found in section 3 on page 16.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.009 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.009).









































































9.4 Atom inclusion [i](#)



At the recommended contour level, 60% of all backbone atoms, 47% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.009) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.4724	 0.2210
A	 0.8067	 0.4220
B	 0.7835	 0.4370
C	 0.8352	 0.4550
D	 0.5836	 0.2180
E	 0.8353	 0.4030
F	 0.8495	 0.4680
G	 0.6575	 0.3200
H	 0.8415	 0.4480
I	 0.5625	 0.2560
J	 0.8256	 0.4530
K	 0.8579	 0.4640
L	 0.7902	 0.4030
M	 0.4524	 0.1740
N	 0.3410	 0.0870
P	 0.7519	 0.3230
T	 0.3710	 0.1150
V	 0.5457	 0.1410
W	 0.4457	 0.2030
a	 0.0436	 0.0430
b	 0.0126	 0.0010
c	 0.0065	 0.0020
d	 0.0193	 0.0010
e	 0.0282	 0.0190
f	 0.0286	 -0.0230
g	 0.0038	 0.0030
h	 0.0042	 0.0010
j	 0.0480	 0.0270
k	 0.0064	 0.0010
m	 0.4212	 0.1310
n	 0.4850	 0.1470
q	 0.2850	 0.0790
r	 0.2053	 0.0950
u	 0.3132	 0.1560
v	 0.2016	 0.0990
x	 0.4436	 0.2670

