

# wwPDB EM Validation Summary Report (i)

Jul 2, 2024 – 07:27 PM JST

PDB ID : 8XT1

EMDB ID : EMD-38633

Title : Cryo-EM structure of the human 39S mitoribosome with 5uM Tigecycline

Authors : Li, X.; Wang, M.; Cheng, J.

Deposited on : 2024-01-10

Resolution : 3.10 Å(reported)

Based on initial model : 7A5I

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at  $\frac{\text{https://www.wwpdb.org/validation/2017/EMValidationReportHelp}}{\text{with specific help available everywhere you see the } \widehat{\textbf{i}} \text{ symbol.}$ 

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.13$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	L1	1559	72%	23% • •
2	L2	69	57% 23%	• 19%
3	LB	305	78%	22%
4	LC	348	87%	13%
5	LD	311	80%	20%
6	LI	267	36% 64%	
7	LJ	261	60%	39%
8	LK	192	90%	• 9%




 $Continued\ from\ previous\ page...$ 





 $Continued\ from\ previous\ page...$ 





## 2 Entry composition (i)

There are 55 unique types of molecules in this entry. The entry contains 100095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 16s rRNA.

Mol	Chain	Residues		I	AltConf	Trace			
1	L1	1500	Total 31847	C 14290	N 5750	O 10307	P 1500	0	0

• Molecule 2 is a RNA chain called Val tRNA.

Mol	Chain	Residues		$\mathbf{A}^{1}$	toms	AltConf	Trace		
2	L2	56	Total 1191	C 534	N 214	O 387	P 56	0	0

• Molecule 3 is a protein called Large ribosomal subunit protein uL2m.

Mol	Chain	Residues		At	AltConf	Trace			
3	LB	237	Total 1851	C 1151	N 375	O 316	S 9	0	0

• Molecule 4 is a protein called Large ribosomal subunit protein uL3m.

Mol	Chain	Residues		At	AltConf	Trace			
4	LC	304	Total 2393	C 1538	N 415	O 429	S 11	0	0

• Molecule 5 is a protein called Large ribosomal subunit protein uL4m.

Mol	Chain	Residues		Ato	AltConf	Trace			
5	LD	250	Total 2013	C 1294	N 365	O 348	S 6	0	0

• Molecule 6 is a protein called Large ribosomal subunit protein bL9m.

Mol	Chain	Residues		Ato	$\overline{\mathbf{m}}$ s	AltConf	Trace	
6	тт	05	Total	С	N	О	0	0
0	L	95	784	498	152	134	U	U



• Molecule 7 is a protein called Large ribosomal subunit protein uL10m.

N	Mol	Chain	Residues		A	toms	AltConf	Trace		
	7	LJ	158	Total 1283	C 828	N 235	O 210	S 10	0	0

• Molecule 8 is a protein called Large ribosomal subunit protein uL11m.

Mol	Chain	Residues		At	oms	AltConf	Trace		
0	1 1/	175	Total	С	N	О	S	0	0
0	LΙΧ	175	1330	847	237	244	2	U	U

• Molecule 9 is a protein called Large ribosomal subunit protein uL13m.

Mol	Chain	Residues		At	oms	AltConf	Trace		
9	LM	177	Total 1451	C 934	N 259	O 251	S 7	0	0

• Molecule 10 is a protein called Large ribosomal subunit protein uL14m.

Mol	Chain	Residues		At	oms	AltConf	Trace		
10	LN	115	Total	С	N	0	S	0	0
			889	559	171	154	Э		

• Molecule 11 is a protein called Large ribosomal subunit protein uL15m.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	LO	287	Total 2305	C 1472	N 425	O 402	S 6	0	0

• Molecule 12 is a protein called Large ribosomal subunit protein uL16m.

Mol	Chain	Residues		At	oms		AltConf	Trace	
12	LP	221	Total 1779	C 1138	N 325	O 306	S 10	0	0

• Molecule 13 is a protein called Large ribosomal subunit protein bL17m.

I	Mol	Chain	Residues		${f Atoms}$					Trace
	13	LQ	152	Total 1245	C 784	N 239	O 215	S 7	0	0

• Molecule 14 is a protein called Mitochondrial ribosomal protein L18, isoform CRA\_b.



Mol	Chain	Residues		Atoms					Trace
1.4	ΙD	146	Total	С	N	О	S	0	0
14	LIU	140	1189	743	226	215	5		U

• Molecule 15 is a protein called Large ribosomal subunit protein bL19m.

Mol	Chain	Residues		$\mathbf{Atoms}$					Trace
15	LS	219	Total 1822	C 1168	N 322	O 323	S 9	0	0

• Molecule 16 is a protein called Large ribosomal subunit protein bL20m.

Mol	Chain	Residues		${f Atoms}$					Trace
16	LT	140	Total	С	N	0	S	0	0
	_		1153	732	231	186	4		

• Molecule 17 is a protein called Large ribosomal subunit protein bL21m.

Mol	Chain	Residues		Atoms					Trace
17	T.IT	160	Total	С	N	О	S	0	0
11	LU	100	1284	829	226	225	4		

• Molecule 18 is a protein called 39S ribosomal protein L22, mitochondrial.

Mol	Chain	Residues	${f Atoms}$					AltConf	Trace
18	LV	166	Total 1368	C 875	N 254	O 232	S 7	0	0

• Molecule 19 is a protein called Large ribosomal subunit protein uL23m.

Mol	Chain	Residues		Atoms					Trace
10	LW	143	Total	С	N	О	S	0	0
19	LW	140	1188	752	224	208	4	U	U

• Molecule 20 is a protein called Large ribosomal subunit protein uL24m.

Mol	Chain	Residues		Ato	AltConf	Trace			
20	LX	202	Total 1652	C 1053	N 294	O 297	S 8	0	0

• Molecule 21 is a protein called Large ribosomal subunit protein bL27m.



Mol	Chain	Residues		At	oms			AltConf	Trace
91	La	111	Total	С	N	О	S	0	0
21	La	111	871	558	164	146	3	0	

• Molecule 22 is a protein called Large ribosomal subunit protein bL28m.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
22	Lb	243	Total 2035	C 1317	N 351	O 362	S 5	0	0

• Molecule 23 is a protein called Large ribosomal subunit protein uL29m.

Mol	Chain	Residues		At	oms			AltConf	Trace
23	Lu	176	Total 1517	C 970	N 291	O 252	S 4	0	0

• Molecule 24 is a protein called Large ribosomal subunit protein uL30m.

Mol	Chain	Residues		At	oms	AltConf	Trace		
24	1.4	120	Total	С	N	О	S	0	0
24	Lu	120	978	626	183	166	3	0	0

• Molecule 25 is a protein called Large ribosomal subunit protein bL32m.

Mol	Chain	Residues		At	oms	AltConf	Trace		
25	Lf	108	Total 880		N 172	O 157	S 6	0	0

• Molecule 26 is a protein called Large ribosomal subunit protein bL33m.

Mol	Chain	Residues		Ato	ms	AltConf	Trace		
26	I or	50	Total	С	N	О	S	0	0
20	Lg	52	433	278	83	70	2	0	U

• Molecule 27 is a protein called Large ribosomal subunit protein bL34m.

Mol	Chain	Residues		Ato	ms	AltConf	Trace		
27	Lh	46	Total 376		N 83	O 59	S 1	0	0

• Molecule 28 is a protein called Large ribosomal subunit protein bL35m.



Mol	Chain	Residues		At	oms		AltConf	Trace	
26	т;	95	Total	С	N	О	S	0	0
20	171	90	831	539	162	127	3		0

• Molecule 29 is a protein called Large ribosomal subunit protein bL36m.

Mol	Chain	Residues		Ato	ms	AltConf	Trace		
20	т;	20	Total	С	N	О	S	0	0
29	ப்	38	341	217	72	48	4	0	U

• Molecule 30 is a protein called Large ribosomal subunit protein mL37.

$\mathbf{Mol}$	Chain	Residues		At	AltConf	Trace			
30	Lk	394	Total 3210	C 2073	N 560	O 566	S 11	0	0

• Molecule 31 is a protein called Large ribosomal subunit protein mL38.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
31	Ll	354	Total 2947	C 1881	N 525	O 532	S 9	0	0

• Molecule 32 is a protein called Large ribosomal subunit protein mL39.

Mol	Chain	Residues		At	AltConf	Trace			
32	Lm	293	Total 2382	C 1525	N 404	O 435	S 18	0	0

• Molecule 33 is a protein called Large ribosomal subunit protein mL40.

Mol	Chain	Residues		At	oms	AltConf	Trace		
33	Ln	99	Total	С	N	О	S	0	0
33	Ln	99	836	535	144	155	2		U

• Molecule 34 is a protein called Large ribosomal subunit protein mL41.

Mol	Chain	Residues		At	oms	AltConf	Trace		
34	Lo	124	Total 997	C 644	N 170	O 181	S 2	0	0

• Molecule 35 is a protein called Large ribosomal subunit protein mL42.



Mol	Chain	Residues		At	oms	AltConf	Trace		
35	Lp	97	Total 815	C 514	N 147	O 149	S 5	0	0

• Molecule 36 is a protein called Large ribosomal subunit protein mL43.

Mol	Chain	Residues		At	oms	AltConf	Trace		
26	T a	148	Total	С	N	О	S	0	0
36	Lq	140	1178	733	229	213	3	0	U

• Molecule 37 is a protein called Large ribosomal subunit protein mL44.

Mo	Chain	Residues		Ato		AltConf	Trace		
37	Lr	275	Total 2217	C 1415	N 383	O 410	S 9	0	0

• Molecule 38 is a protein called Large ribosomal subunit protein mL45.

Mol	Chain	Residues		At	oms			AltConf	Trace
38	Ls	214	Total 1754	C 1117	N 304	O 320	S 13	0	0

• Molecule 39 is a protein called Large ribosomal subunit protein mL46.

Mol	Chain	Residues		Ato	AltConf	Trace			
39	Lt	217	Total 1762	C 1124	N 310	O 323	S 5	0	0

• Molecule 40 is a protein called Large ribosomal subunit protein mL48.

Mol	Chain	Residues		At	oms	AltConf	Trace		
40	Τ.,,	131	Total	С	N	О	S	0	0
40	Lv	191	1039	663	169	203	4	U	U

• Molecule 41 is a protein called Large ribosomal subunit protein mL49.

Mol	Chain	Residues		At	oms	AltConf	Trace		
41	Lw	132	Total 1097	C 710	N 191	O 194	S 2	0	0

• Molecule 42 is a protein called Large ribosomal subunit protein mL50.



Mol	Chain	Residues		At	oms	AltConf	Trace		
49	Lv	110	Total	С	N	О	S	0	0
42	LX	110	895	568	156	168	3	U	U

• Molecule 43 is a protein called Large ribosomal subunit protein mL51.

Mol	Chain	Residues		At	oms	AltConf	Trace		
43	Ly	97	Total 827	C 532	N 165	O 126	S	0	0

• Molecule 44 is a protein called 39S ribosomal protein L52, mitochondrial.

Mol	Chain	Residues		At	oms			AltConf	Trace
44	Lz	92	Total 732	C 454	N 142	O 134	S 2	0	0

• Molecule 45 is a protein called Large ribosomal subunit protein mL53.

Mol	Chain	Residues		At	oms	AltConf	Trace		
45	1.3	96	Total	С	N	О	S	0	0
40	ЦЭ	30	743	462	143	133	5		

• Molecule 46 is a protein called Large ribosomal subunit protein mL54.

Mol	Chain	Residues		At	oms	AltConf	Trace		
46	L4	83	Total 703	C 446	N 124	O 130	S 3	0	0

• Molecule 47 is a protein called Large ribosomal subunit protein mL55.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
47	L5	45	Total 372	C 232	N 76	O 62	S 2	0	0

• Molecule 48 is a protein called Large ribosomal subunit protein mL63.

Mol	Chain	Residues	Atoms				AltConf	Trace	
48	L6	94	Total 797	C 501	N 165	O 128	S 3	0	0

• Molecule 49 is a protein called Large ribosomal subunit protein mL62.



Mol	Chain	Residues	Atoms				AltConf	Trace	
49	L7	127	Total	C	N	0	S	0	0
			1058	661	201	192	$^4$		

• Molecule 50 is a protein called Large ribosomal subunit protein mL64.

Mol	Chain	Residues	Atoms				AltConf	Trace	
50	ΤO	128	Total	С	N	О	S	0	0
30	L8	120	1076	671	208	192	5	U	0

• Molecule 51 is a protein called Large ribosomal subunit protein mL66.

Mol	Chain	Residues		At	oms			AltConf	Trace
<b>E</b> 1	CD	146	Total	С	N	О	S	0	0
91	SN	140	1203	764	232	199	8	0	U

• Molecule 52 is a protein called Large ribosomal subunit protein mL65.


Mol	Chain	Residues	Atoms				AltConf	Trace	
52	Sf	370	Total 3036	C 1946	N 542	O 534	S 14	0	0

• Molecule 53 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
53	L1	90	Total Mg 90 90	0
53	LB	1	Total Mg 1 1	0
53	LC	1	Total Mg 1 1	0
53	La	1	Total Mg 1 1	0
53	Lw	1	Total Mg 1 1	0
53	L6	1	Total Mg 1 1	0

• Molecule 54 is TIGECYCLINE (three-letter code: T1C) (formula:  $C_{29}H_{41}N_5O_8$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf
54	Т 1	1	Total	С	N	О	0
04	171	1	42	29	5	8	U

• Molecule 55 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	AltConf
55	Lf	1	Total Zn 1 1	0
55	Lj	1	Total Zn 1 1	0
55	SR	1	Total Zn 1 1	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 16s rRNA













 $\bullet$  Molecule 13: Large ribosomal subunit protein bL17m













Chain Ll:







93%




• Molecule 37: Large ribosomal subunit protein mL44













# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	57266	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION; Relion	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.070	Depositor
Minimum map value	-0.017	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	446.88, 446.88, 446.88	wwPDB
Map dimensions	420, 420, 420	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.064, 1.064, 1.064	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, T1C, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	ond lengths	l I	Bond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	L1	0.29	0/35628	0.97	105/55448~(0.2%)
2	L2	0.28	0/1328	1.00	1/2056~(0.0%)
3	LB	0.27	0/1888	0.62	0/2538
4	LC	0.28	0/2462	0.55	0/3340
5	LD	0.34	0/2071	0.63	0/2817
6	LI	0.35	0/798	0.77	0/1073
7	LJ	0.39	1/1308 (0.1%)	0.83	2/1761 (0.1%)
8	LK	0.31	0/1348	0.63	1/1813 (0.1%)
9	LM	0.29	0/1495	0.57	0/2029
10	LN	0.29	0/904	0.59	0/1218
11	LO	0.29	0/2359	0.62	1/3185 (0.0%)
12	LP	0.28	0/1826	0.57	0/2458
13	LQ	0.30	0/1269	0.64	0/1708
14	LR	0.30	0/1215	0.68	2/1645 (0.1%)
15	LS	0.32	0/1863	0.60	0/2509
16	LT	0.30	0/1174	0.59	0/1572
17	LU	0.31	0/1311	0.64	0/1778
18	LV	0.30	0/1402	0.57	0/1886
19	LW	0.34	0/1217	0.68	1/1644~(0.1%)
20	LX	0.29	0/1697	0.62	0/2302
21	La	0.28	0/893	0.56	0/1204
22	Lb	0.28	0/2090	0.55	$1/2825 \ (0.0\%)$
23	Lu	0.30	0/1552	0.60	0/2079
24	Ld	0.32	0/1003	0.57	0/1354
25	Lf	0.32	0/895	0.60	0/1201
26	Lg	0.29	0/438	0.67	0/583
27	Lh	0.25	0/382	0.63	0/507
28	Li	0.27	0/852	0.60	1/1136 (0.1%)
29	Lj	0.25	0/349	0.60	0/461
30	Lk	0.28	0/3305	0.55	0/4502
31	Ll	0.32	0/3042	0.65	1/4140 (0.0%)
32	Lm	0.29	0/2439	0.55	0/3299



Mol	Chain	Во	ond lengths	I	Bond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
33	Ln	0.38	0/855	0.78	3/1152 (0.3%)
34	Lo	0.33	0/1025	0.64	0/1379
35	Lp	0.32	0/839	0.64	0/1136
36	Lq	0.28	0/1202	0.59	0/1626
37	Lr	0.29	0/2264	0.55	1/3059 (0.0%)
38	Ls	0.31	0/1800	0.62	0/2436
39	Lt	0.30	0/1797	0.68	1/2422 (0.0%)
40	Lv	0.30	0/1055	0.63	0/1427
41	Lw	0.34	0/1134	0.65	0/1547
42	Lx	0.33	0/918	0.64	0/1249
43	Ly	0.26	0/849	0.57	0/1135
44	Lz	0.29	0/747	0.57	0/1005
45	L3	0.35	0/754	0.74	1/1017 (0.1%)
46	L4	0.31	0/722	0.68	0/978
47	L5	0.27	0/379	0.76	1/510 (0.2%)
48	L6	0.32	0/818	0.62	0/1097
49	L7	0.28	0/1071	0.61	0/1433
50	L8	0.28	0/1107	0.58	0/1498
51	SR	0.33	0/1238	0.65	0/1676
52	Sf	0.29	0/3114	0.58	0/4225
All	All	0.30	1/105491 (0.0%)	0.77	123/150078 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
8	LK	0	2
11	LO	0	1
12	LP	0	1
13	LQ	0	1
18	LV	0	1
23	Lu	0	1
31	Ll	0	1
All	All	0	8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
7	LJ	163	GLU	CG-CD	-5.88	1.43	1.51



	$\overline{}$	C 1	$\Omega\Omega$	1 1	1	. 1.		1. 1	1 1
The worst	ה	$\cap$ t L	23	hond	angle	Outliers	are	listed	pelow.
THE WOLDS	$\circ$		40	DOM	angic	Outilities	COL C	mouca	DCIOW.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	L1	23	С	C2-N1-C1'	11.02	130.92	118.80
1	L1	23	С	N1-C2-O2	10.61	125.27	118.90
1	L1	23	С	N3-C2-O2	-8.84	115.72	121.90
1	L1	823	С	C2-N1-C1'	8.28	127.90	118.80
1	L1	23	С	C6-N1-C1'	-8.08	111.10	120.80

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
8	LK	57	THR	Peptide
8	LK	60	ILE	Peptide
11	LO	39	ARG	Sidechain
12	LP	73	ARG	Sidechain
13	LQ	110	ILE	Peptide

#### 5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percei	ntiles
3	LB	$235/305\ (77\%)$	227 (97%)	8 (3%)	0	100	100
4	LC	302/348~(87%)	277 (92%)	25 (8%)	0	100	100
5	LD	$248/311\ (80\%)$	238 (96%)	10 (4%)	0	100	100
6	LI	$93/267\ (35\%)$	88 (95%)	5 (5%)	0	100	100
7	LJ	154/261 (59%)	138 (90%)	16 (10%)	0	100	100
8	LK	173/192~(90%)	163 (94%)	10 (6%)	0	100	100



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
9	LM	$175/178\ (98\%)$	165 (94%)	10 (6%)	0	100	100
10	LN	$113/145\ (78\%)$	107 (95%)	6 (5%)	0	100	100
11	LO	$285/296\ (96\%)$	276 (97%)	9 (3%)	0	100	100
12	LP	$219/251\ (87\%)$	212 (97%)	7 (3%)	0	100	100
13	LQ	$150/175\ (86\%)$	138 (92%)	11 (7%)	1 (1%)	22	57
14	LR	144/179~(80%)	140 (97%)	4 (3%)	0	100	100
15	LS	$217/292\ (74\%)$	202 (93%)	15 (7%)	0	100	100
16	LT	138/149~(93%)	134 (97%)	4 (3%)	0	100	100
17	LU	158/205 (77%)	151 (96%)	7 (4%)	0	100	100
18	LV	164/212 (77%)	157 (96%)	7 (4%)	0	100	100
19	LW	139/153 (91%)	138 (99%)	1 (1%)	0	100	100
20	LX	200/216 (93%)	185 (92%)	15 (8%)	0	100	100
21	La	109/148 (74%)	104 (95%)	5 (5%)	0	100	100
22	Lb	241/256 (94%)	227 (94%)	14 (6%)	0	100	100
23	Lu	174/250 (70%)	172 (99%)	2 (1%)	0	100	100
24	Ld	118/161 (73%)	111 (94%)	7 (6%)	0	100	100
25	Lf	106/188 (56%)	101 (95%)	5 (5%)	0	100	100
26	Lg	50/65 (77%)	50 (100%)	0	0	100	100
27	Lh	44/92 (48%)	43 (98%)	1 (2%)	0	100	100
28	Li	93/188 (50%)	91 (98%)	2 (2%)	0	100	100
29	Lj	36/103 (35%)	36 (100%)	0	0	100	100
30	Lk	392/423 (93%)	382 (97%)	10 (3%)	0	100	100
31	Ll	352/380 (93%)	327 (93%)	25 (7%)	0	100	100
32	Lm	291/338 (86%)	283 (97%)	8 (3%)	0	100	100
33	Ln	97/206 (47%)	87 (90%)	10 (10%)	0	100	100
34	Lo	122/137 (89%)	115 (94%)	7 (6%)	0	100	100
35	Lp	93/142 (66%)	89 (96%)	4 (4%)	0	100	100
36	Lq	146/215 (68%)	135 (92%)	11 (8%)	0	100	100
37	Lr	271/332 (82%)	265 (98%)	6 (2%)	0	100	100
38	Ls	210/306 (69%)	204 (97%)	6 (3%)	0	100	100
39	Lt	211/279 (76%)	186 (88%)	23 (11%)	2 (1%)	17	52



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
40	Lv	125/212~(59%)	121 (97%)	4 (3%)	0	100	100
41	Lw	130/166~(78%)	122 (94%)	8 (6%)	0	100	100
42	Lx	108/158~(68%)	105 (97%)	3 (3%)	0	100	100
43	Ly	95/128 (74%)	90 (95%)	5 (5%)	0	100	100
44	Lz	90/123~(73%)	86 (96%)	4 (4%)	0	100	100
45	L3	94/112 (84%)	88 (94%)	6 (6%)	0	100	100
46	L4	81/138 (59%)	75 (93%)	6 (7%)	0	100	100
47	L5	43/128 (34%)	41 (95%)	2 (5%)	0	100	100
48	L6	92/102 (90%)	89 (97%)	3 (3%)	0	100	100
49	L7	119/206 (58%)	114 (96%)	5 (4%)	0	100	100
50	L8	126/222 (57%)	123 (98%)	3 (2%)	0	100	100
51	SR	140/196 (71%)	138 (99%)	2 (1%)	0	100	100
52	Sf	366/439 (83%)	353 (96%)	12 (3%)	1 (0%)	41	73
All	All	8072/10674 (76%)	7689 (95%)	379 (5%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
39	Lt	266	PRO
39	Lt	270	ALA
52	Sf	260	GLU
13	LQ	111	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	LB	191/245~(78%)	191 (100%)	0	100 100
4	LC	258/290 (89%)	258 (100%)	0	100 100
5	LD	$217/262\ (83\%)$	217 (100%)	0	100 100



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
6	LI	$86/228 \; (38\%)$	86 (100%)	0	100	100
7	LJ	145/232~(62%)	145 (100%)	0	100	100
8	LK	138/150 (92%)	138 (100%)	0	100	100
9	LM	155/156 (99%)	155 (100%)	0	100	100
10	LN	98/124 (79%)	98 (100%)	0	100	100
11	LO	245/249 (98%)	245 (100%)	0	100	100
12	LP	188/211 (89%)	188 (100%)	0	100	100
13	LQ	133/150~(89%)	132 (99%)	1 (1%)	81	92
14	LR	128/154~(83%)	128 (100%)	0	100	100
15	LS	201/256 (78%)	200 (100%)	1 (0%)	88	94
16	LT	118/126 (94%)	117 (99%)	1 (1%)	81	92
17	LU	145/180 (81%)	144 (99%)	1 (1%)	84	93
18	LV	146/182 (80%)	146 (100%)	0	100	100
19	LW	128/135 (95%)	128 (100%)	0	100	100
20	LX	180/191 (94%)	179 (99%)	1 (1%)	86	94
21	La	91/119 (76%)	91 (100%)	0	100	100
22	Lb	219/229 (96%)	219 (100%)	0	100	100
23	Lu	159/223~(71%)	159 (100%)	0	100	100
24	Ld	111/147 (76%)	110 (99%)	1 (1%)	78	91
25	Lf	97/164~(59%)	97 (100%)	0	100	100
26	Lg	49/60 (82%)	49 (100%)	0	100	100
27	Lh	40/72 (56%)	40 (100%)	0	100	100
28	Li	88/166 (53%)	88 (100%)	0	100	100
29	Lj	37/89~(42%)	37 (100%)	0	100	100
30	Lk	353/368 (96%)	353 (100%)	0	100	100
31	Ll	313/332 (94%)	313 (100%)	0	100	100
32	Lm	269/303~(89%)	269 (100%)	0	100	100
33	Ln	91/190 (48%)	91 (100%)	0	100	100
34	Lo	$104/112 \ (93\%)$	102 (98%)	2 (2%)	57	81
35	Lp	93/133 (70%)	92 (99%)	1 (1%)	73	89
36	Lq	130/186 (70%)	130 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
37	$\operatorname{Lr}$	241/288~(84%)	241 (100%)	0	100	100
38	Ls	196/274~(72%)	196 (100%)	0	100	100
39	$\operatorname{Lt}$	188/236~(80%)	184 (98%)	4 (2%)	53	79
40	Lv	117/188~(62%)	117 (100%)	0	100	100
41	Lw	122/148~(82%)	122 (100%)	0	100	100
42	Lx	104/148~(70%)	103 (99%)	1 (1%)	76	90
43	Ly	86/110 (78%)	86 (100%)	0	100	100
44	Lz	73/97~(75%)	73 (100%)	0	100	100
45	L3	81/90 (90%)	81 (100%)	0	100	100
46	L4	78/116~(67%)	78 (100%)	0	100	100
47	L5	40/113~(35%)	39 (98%)	1 (2%)	47	75
48	L6	80/87~(92%)	80 (100%)	0	100	100
49	L7	117/181~(65%)	117 (100%)	0	100	100
50	L8	110/178~(62%)	110 (100%)	0	100	100
51	SR	133/169 (79%)	132 (99%)	1 (1%)	81	92
52	Sf	326/381 (86%)	325 (100%)	1 (0%)	92	96
All	All	7236/9218 (78%)	7219 (100%)	17 (0%)	93	97

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
47	L5	58	LYS
52	Sf	161	ARG
34	Lo	116	LYS
35	Lp	99	LYS
39	Lt	127	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
45	L3	15	GLN
52	Sf	373	GLN
52	Sf	375	ASN
16	LT	94	GLN
33	Ln	143	GLN



#### 5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	L1	1491/1559~(95%)	346 (23%)	15 (1%)
2	L2	51/69 (73%)	17 (33%)	0
All	All	1542/1628 (94%)	363 (23%)	15 (0%)

5 of 363 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	L1	5	A
1	L1	6	A
1	L1	8	С
1	L1	9	U
1	L1	11	G

5 of 15 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	L1	575	A
1	L1	1319	G
1	L1	837	A
1	L1	1371	U
1	L1	930	A

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 99 ligands modelled in this entry, 98 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res Link		Bo	nd leng	ths	В	ond ang	les
		туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
	54	T1C	L1	1690	53	44,45,45	1.18	4 (9%)	53,72,72	1.07	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
54	T1C	L1	1690	53	-	11/22/80/80	0/4/4/4

#### All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)
54	L1	1690	T1C	C21-N21	5.45	1.47	1.33
54	L1	1690	T1C	O11-C11	2.21	1.27	1.23
54	L1	1690	T1C	C7-N7	2.16	1.48	1.42
54	L1	1690	T1C	C4-N4	2.15	1.52	1.47

#### All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
54	L1	1690	T1C	C1-C1C-C12	3.11	113.53	109.88
54	L1	1690	T1C	C11-C1B-C12	2.86	121.07	118.80
54	L1	1690	T1C	O1C-C1C-C12	-2.71	105.81	110.14
54	L1	1690	T1C	C1C-C41-C4	2.69	115.31	111.64
54	L1	1690	T1C	C1C-C1-C2	2.12	119.12	115.75

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
54	L1	1690	T1C	C92-C91-N9-C9
54	L1	1690	T1C	C41-C4-N4-C43
54	L1	1690	T1C	C3-C4-N4-C43
54	L1	1690	T1C	C3-C4-N4-C42
54	L1	1690	T1C	C3-C2-C21-O21

There are no ring outliers.



No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-38633. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

#### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map





The images above show the map projected in three orthogonal directions.




## 6.2 Central slices (i)

#### 6.2.1 Primary map









X Index: 210


Y Index: 210

Z Index: 210

#### 6.2.2 Raw map







X Index: 210

Y Index: 210


Z Index: 210


The images above show central slices of the map in three orthogonal directions.




# 6.3 Largest variance slices (i)

#### 6.3.1 Primary map









 ${\rm X~Index:~208}$ 


Y Index: 201

Z Index: 211

#### 6.3.2 Raw map







X Index: 208

Y Index: 201


Z Index: 211

The images above show the largest variance slices of the map in three orthogonal directions.




# 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



#### 6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

# 6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


# 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



## 7.2 Volume estimate (i)




The volume at the recommended contour level is  $1215~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $1097~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



# 7.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.323  $\rm \AA^{-1}$ 



# 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

## 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.323  $\rm \mathring{A}^{-1}$ 



# 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
(A)	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.94	6.62	4.05


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.94 differs from the reported value 3.1 by more than 10 %





# 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-38633 and PDB model 8XT1. Per-residue inclusion information can be found in section 3 on page 14.

# 9.1 Map-model overlay (i)








The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




# 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).



# 9.4 Atom inclusion (i)



At the recommended contour level, 98% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.



# 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9240	0.4720
L1	0.9850	0.5040
L2	0.9430	0.2800
L3	0.8810	0.3790
L4	0.7640	0.2940
L5	0.6690	0.1720
L6	0.9620	0.5170
L7	0.8810	0.4380
L8	0.8690	0.4140
LB	0.9290	0.5130
LC	0.9410	0.5090
LD	0.9510	0.5220
LI	0.8900	0.4540
LJ	0.8350	0.3700
LK	0.7500	0.2480
LM	0.9580	0.5160
LN	0.9160	0.5050
LO	0.9450	0.5150
LP	0.9300	0.5070
LQ	0.9330	0.5110
LR	0.9160	0.4580
LS	0.8890	0.4800
$\operatorname{LT}$	0.9400	0.5120
LU	0.9270	0.5130
LV	0.9440	0.5170
LW	0.9160	0.4950
LX	0.8800	0.4520
La	0.9640	0.5360
Lb	0.9080	0.4800
Ld	0.9380	0.5210
Lf	0.9250	0.5010
Lg	0.9450	0.4890
Lh	0.9860	0.5490
Li	0.9760	0.5470
Lj	0.9720	0.5370



Continued on next page...



## $Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
Lk	0.9060	0.4860
Ll	0.8850	0.4080
Lm	0.8790	0.4460
Ln	0.6680	0.1900
Lo	0.9050	0.4810
Lp	0.9000	0.4880
Lq	0.9500	0.5270
Lr	0.9040	0.4820
Ls	0.8260	0.4260
Lt	0.5790	0.1240
Lu	0.9230	0.4930
Lv	0.7080	0.2640
Lw	0.9360	0.4980
Lx	0.8830	0.4500
Ly	0.9670	0.5370
Lz	0.9100	0.4680
SR	0.9600	0.5070
Sf	0.9220	0.5040

