

Full wwPDB X-ray Structure Validation Report (i)

Nov 13, 2023 – 05:39 PM JST

PDB ID : 5XUX

Title : Crystal structure of Rib7 from Methanosarcina mazei

Authors: Yeh, T.M.; Chen, S.C.; Chang, T.H.; Huang, M.F.; Liaw, S.H.

Deposited on : 2017-06-26

Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Wiedite	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	240	94%	5% •
1	В	240	89%	5% 7%
1	С	240	5% 89%	• 7%
1	D	240	6%	5% 7%
1	E	240	88%	5% 7%
1	F	240	92%	7% •



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAP	F	301	-	-	-	X



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 11043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Conserved protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	С	N	О	S	0	0	0
1	A	230	1808	1136	320	344	8	0	U	
1	В	224	Total	С	N	О	S	0	0	0
1	D	224	1698	1072	291	327	8		U	
1	С	224	Total	С	N	О	S	0	0	0
1		224	1694	1069	291	327	7	U		
1	D	D 223	Total	С	N	О	S	0	0	0
1	D	229	1690	1067	290	326	7		0	
1	Е	223	Total	С	N	О	S	0	0	0
1	12	229	1690	1067	290	326	7		0	
1	F	238	Total	С	N	О	S	0	0	0
1	Г	230	1808	1136	320	344	8	U	0	

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q8PYN5
A	-10	ARG	-	expression tag	UNP Q8PYN5
A	-9	GLY	-	expression tag	UNP Q8PYN5
A	-8	SER	-	expression tag	UNP Q8PYN5
A	-7	HIS	-	expression tag	UNP Q8PYN5
A	-6	HIS	-	expression tag	UNP Q8PYN5
A	-5	HIS	-	expression tag	UNP Q8PYN5
A	-4	HIS	-	expression tag	UNP Q8PYN5
A	-3	HIS	-	expression tag	UNP Q8PYN5
A	-2	HIS	-	expression tag	UNP Q8PYN5
A	-1	GLY	-	expression tag	UNP Q8PYN5
A	0	SER	-	expression tag	UNP Q8PYN5
В	-11	MET	-	expression tag	UNP Q8PYN5
В	-10	ARG	-	expression tag	UNP Q8PYN5
В	-9	GLY	-	expression tag	UNP Q8PYN5
В	-8	SER	-	expression tag	UNP Q8PYN5
В	-7	HIS	-	expression tag	UNP Q8PYN5



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
В	-6	HIS	-	expression tag	UNP Q8PYN5
В	-5	HIS	-	expression tag	UNP Q8PYN5
В	-4	HIS	-	expression tag	UNP Q8PYN5
В	-3	HIS	-	expression tag	UNP Q8PYN5
В	-2	HIS	-	expression tag	UNP Q8PYN5
В	-1	GLY	_	expression tag	UNP Q8PYN5
В	0	SER	-	expression tag	UNP Q8PYN5
С	-11	MET	-	expression tag	UNP Q8PYN5
С	-10	ARG	-	expression tag	UNP Q8PYN5
С	-9	GLY	_	expression tag	UNP Q8PYN5
С	-8	SER	-	expression tag	UNP Q8PYN5
С	-7	HIS	_	expression tag	UNP Q8PYN5
С	-6	HIS	_	expression tag	UNP Q8PYN5
С	-5	HIS	-	expression tag	UNP Q8PYN5
С	-4	HIS	-	expression tag	UNP Q8PYN5
С	-3	HIS	_	expression tag	UNP Q8PYN5
С	-2	HIS	-	expression tag	UNP Q8PYN5
С	-1	GLY	-	expression tag	UNP Q8PYN5
С	0	SER	-	expression tag	UNP Q8PYN5
D	-11	MET	-	expression tag	UNP Q8PYN5
D	-10	ARG	-	expression tag	UNP Q8PYN5
D	-9	GLY	-	expression tag	UNP Q8PYN5
D	-8	SER	-	expression tag	UNP Q8PYN5
D	-7	HIS	-	expression tag	UNP Q8PYN5
D	-6	HIS	-	expression tag	UNP Q8PYN5
D	-5	HIS	-	expression tag	UNP Q8PYN5
D	-4	HIS	-	expression tag	UNP Q8PYN5
D	-3	HIS	-	expression tag	UNP Q8PYN5
D	-2	HIS	-	expression tag	UNP Q8PYN5
D	-1	GLY	-	expression tag	UNP Q8PYN5
D	0	SER	-	expression tag	UNP Q8PYN5
Е	-11	MET	-	expression tag	UNP Q8PYN5
Е	-10	ARG	-	expression tag	UNP Q8PYN5
Е	-9	GLY	-	expression tag	UNP Q8PYN5
Е	-8	SER	-	expression tag	UNP Q8PYN5
Е	-7	HIS	-	expression tag	UNP Q8PYN5
Е	-6	HIS	-	expression tag	UNP Q8PYN5
Е	-5	HIS	-	expression tag	UNP Q8PYN5
Е	-4	HIS	-	expression tag	UNP Q8PYN5
Е	-3	HIS	-	expression tag	UNP Q8PYN5
Е	-2	HIS	-	expression tag	UNP Q8PYN5
Е	-1	GLY		expression tag	UNP Q8PYN5



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Е	0	SER	-	expression tag	UNP Q8PYN5
F	-11	MET	-	expression tag	UNP Q8PYN5
F	-10	ARG	-	expression tag	UNP Q8PYN5
F	-9	GLY	-	expression tag	UNP Q8PYN5
F	-8	SER	-	expression tag	UNP Q8PYN5
F	-7	HIS	-	expression tag	UNP Q8PYN5
F	-6	HIS	-	expression tag	UNP Q8PYN5
F	-5	HIS	-	expression tag	UNP Q8PYN5
F	-4	HIS	-	expression tag	UNP Q8PYN5
F	-3	HIS	-	expression tag	UNP Q8PYN5
F	-2	HIS	-	expression tag	UNP Q8PYN5
F	-1	GLY	-	expression tag	UNP Q8PYN5
F	0	SER	-	expression tag	UNP Q8PYN5

• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0
2	A	1	48	21	7	17	3	U	U
2	В	1	Total	С	N	О	Р	0	0
2	Б	1	48	21	7	17	3	U	0
2	C	1	Total	С	N	О	Р	0	0
2		1	48	21	7	17	3	U	0
2	D	1	Total	С	N	О	Р	0	0
	ש	1	48	21	7	17	3	U	U



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Ŀ	1	Total	С	N	О	Р	0	0
2	E	1	48	21	7	17	3	U	
9	E	1	Total	С	N	О	Р	0	0
2	Г	1	48	21	7	17	3	U	0

$\bullet\,$ Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	72	Total O 72 72	0	0
3	В	83	Total O 83 83	0	0
3	С	30	Total O 30 30	0	0
3	D	27	Total O 27 27	0	0
3	Е	84	Total O 84 84	0	0
3	F	71	Total O 71 71	0	0



Chain E:

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Conserved protein • Molecule 1: Conserved protein Chain B: • Molecule 1: Conserved protein Chain C: 89% 7% • Molecule 1: Conserved protein Chain D: 7% • Molecule 1: Conserved protein

88%





• Molecule 1: Conserved protein







4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants	118.46Å 118.46Å 375.55Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 - 2.27	Depositor
resolution (A)	47.54 - 2.27	EDS
% Data completeness	99.7 (50.00-2.27)	Depositor
(in resolution range)	99.8 (47.54-2.27)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.10 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.198 , 0.228	Depositor
it, it _{free}	0.204 , 0.230	DCC
R_{free} test set	4451 reflections $(4.91%)$	wwPDB-VP
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 23.3	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.480 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11043	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.43	0/1834	0.67	0/2462	
1	В	0.44	0/1718	0.67	$2/2308 \; (0.1\%)$	
1	С	0.42	0/1714	0.64	0/2303	
1	D	0.42	0/1710	0.62	0/2298	
1	Е	0.43	0/1710	0.67	$1/2298 \; (0.0\%)$	
1	F	0.44	0/1834	0.69	$2/2462 \ (0.1\%)$	
All	All	0.43	0/10520	0.66	5/14131 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	1
1	Е	0	2
1	F	0	2
All	All	0	6

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	F	3	ARG	NE-CZ-NH2	7.33	123.97	120.30
1	F	3	ARG	NE-CZ-NH1	-6.91	116.84	120.30
1	Е	23	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	В	23	ARG	NE-CZ-NH1	-5.61	117.49	120.30
1	В	23	ARG	NE-CZ-NH2	5.06	122.83	120.30



There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	156	GLU	Peptide
1	В	157	GLY	Peptide
1	Е	156	GLU	Peptide
1	Е	157	GLY	Peptide
1	F	156	GLU	Peptide
1	F	157	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1859	6	0
1	В	1698	0	1766	4	0
1	С	1694	0	1757	5	0
1	D	1690	0	1754	6	0
1	Е	1690	0	1754	5	0
1	F	1808	0	1859	6	0
2	A	48	0	25	1	0
2	В	48	0	25	0	0
2	С	48	0	25	0	0
2	D	48	0	25	1	0
2	Ε	48	0	25	0	0
2	F	48	0	25	0	0
3	A	72	0	0	1	0
3	В	83	0	0	0	0
3	С	30	0	0	0	0
3	D	27	0	0	0	0
3	Е	84	0	0	0	0
3	F	71	0	0	0	0
All	All	11043	0	10899	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



A	A., 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
1:D:18:LEU:HD21	1:D:191:PHE:HA	1.85	0.58
1:A:-5:HIS:ND1	3:A:401:HOH:O	2.32	0.57
1:A:107:ILE:HD11	1:A:148:MET:HE3	1.86	0.57
1:F:29:SER:HB2	1:F:33:ASP:HB3	1.87	0.57
1:B:5:PHE:CD1	1:C:76:GLY:HA3	2.42	0.55
1:C:6:ILE:HD13	1:C:153:LEU:HB3	1.90	0.54
1:E:29:SER:HB2	1:E:33:ASP:HB3	1.90	0.52
1:D:76:GLY:HA3	1:E:5:PHE:CD1	2.44	0.52
1:A:29:SER:HB2	1:A:33:ASP:HB3	1.94	0.49
1:D:41:ARG:NH2	1:D:156:GLU:OE2	2.45	0.49
1:B:49:VAL:HG22	1:B:53:THR:HB	1.95	0.48
1:E:119:ILE:HG23	1:E:128:VAL:HG11	1.94	0.48
1:C:6:ILE:CD1	1:C:153:LEU:HD23	2.44	0.47
1:E:186:LYS:HB3	1:F:193:ASP:HB3	1.96	0.47
1:F:148:MET:HB3	1:F:148:MET:HE2	1.79	0.47
1:B:29:SER:HB2	1:B:33:ASP:HB3	1.96	0.47
1:A:82:VAL:HG21	1:A:150:ILE:HD11	1.96	0.46
1:D:18:LEU:HB3	2:D:301:NAP:H2N	1.98	0.46
1:A:88:SER:N	2:A:301:NAP:O2X	2.48	0.45
1:F:3:ARG:NH1	1:F:172:ASP:OD2	2.50	0.45
1:A:32:LEU:HD12	1:A:214:GLU:HG3	1.99	0.44
1:D:48:MET:HB2	1:D:153:LEU:HD11	1.99	0.44
1:C:48:MET:HB2	1:C:153:LEU:HD11	2.00	0.43
1:C:51:ILE:HD11	1:C:87:ASP:HA	1.99	0.43
1:F:48:MET:HB2	1:F:153:LEU:HD11	2.00	0.42
1:B:111:SER:HA	1:B:131:THR:O	2.21	0.40
1:D:5:PHE:CD1	1:E:76:GLY:HA3	2.57	0.40
1:F:110:VAL:O	1:F:130:LYS:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	236/240 (98%)	233 (99%)	3 (1%)	0	100	100
1	В	222/240 (92%)	218 (98%)	4 (2%)	0	100	100
1	С	222/240 (92%)	216 (97%)	6 (3%)	0	100	100
1	D	221/240 (92%)	213 (96%)	7 (3%)	1 (0%)	29	34
1	E	221/240 (92%)	217 (98%)	4 (2%)	0	100	100
1	F	236/240 (98%)	229 (97%)	7 (3%)	0	100	100
All	All	1358/1440 (94%)	1326 (98%)	31 (2%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	133	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	193/195 (99%)	192 (100%)	1 (0%)	88 94
1	В	182/195 (93%)	180 (99%)	2 (1%)	73 84
1	С	181/195 (93%)	177 (98%)	4 (2%)	52 66
1	D	181/195 (93%)	177 (98%)	4 (2%)	52 66
1	E	181/195 (93%)	179 (99%)	2 (1%)	73 84
1	F	193/195 (99%)	187 (97%)	6 (3%)	40 53
All	All	1111/1170 (95%)	1092 (98%)	19 (2%)	60 74

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LYS
1	В	21	LYS
1	В	97	ASP
1	С	83	ARG



Continued from previous page...

Mol	Chain	Res	Type
1	С	94	LEU
1	С	101	LYS
1	С	112	ASN
1	D	91	ARG
1	D	97	ASP
1	D	144	LYS
1	D	186	LYS
1	Е	97	ASP
1	Е	176	THR
1	F	-6	HIS
1	F	3	ARG
1	F	21	LYS
1	F	97	ASP
1	F	199	GLU
1	F	226	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	fol Type Chain Res		Link	Bond lengths			Bond angles			
MIOI	vioi Type Chain	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	В	301	-	45,52,52	0.91	2 (4%)	56,80,80	1.25	5 (8%)
2	NAP	С	301	-	45,52,52	0.89	3 (6%)	56,80,80	1.17	6 (10%)
2	NAP	A	301	-	45,52,52	0.92	2 (4%)	56,80,80	1.12	4 (7%)
2	NAP	F	301	-	45,52,52	0.91	2 (4%)	56,80,80	1.15	4 (7%)
2	NAP	Е	301	-	45,52,52	0.90	1 (2%)	56,80,80	1.14	5 (8%)
2	NAP	D	301	-	45,52,52	0.87	3 (6%)	56,80,80	1.28	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	В	301	-	-	4/31/67/67	0/5/5/5
2	NAP	С	301	-	-	3/31/67/67	0/5/5/5
2	NAP	A	301	-	-	12/31/67/67	0/5/5/5
2	NAP	F	301	-	-	12/31/67/67	0/5/5/5
2	NAP	E	301	-	-	2/31/67/67	0/5/5/5
2	NAP	D	301	-	-	7/31/67/67	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\mathring{\mathrm{A}})$	Ideal(Å)
2	F	301	NAP	C5A-C4A	2.85	1.48	1.40
2	A	301	NAP	C5A-C4A	2.81	1.48	1.40
2	С	301	NAP	C5A-C4A	2.58	1.47	1.40
2	Ε	301	NAP	C5A-C4A	2.56	1.47	1.40
2	D	301	NAP	C5A-C4A	2.56	1.47	1.40
2	В	301	NAP	C5A-C4A	2.40	1.47	1.40
2	A	301	NAP	C2A-N3A	2.34	1.35	1.32
2	С	301	NAP	O4D-C1D	2.32	1.44	1.41
2	F	301	NAP	C2A-N3A	2.31	1.35	1.32
2	В	301	NAP	P2B-O2B	2.26	1.63	1.59
2	D	301	NAP	O4D-C1D	2.25	1.44	1.41
2	С	301	NAP	C2A-N3A	2.17	1.35	1.32
2	D	301	NAP	C2A-N3A	2.15	1.35	1.32



All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
2	В	301	NAP	N3A-C2A-N1A	-3.87	122.63	128.68
2	Е	301	NAP	N3A-C2A-N1A	-3.60	123.06	128.68
2	D	301	NAP	N3A-C2A-N1A	-3.59	123.07	128.68
2	В	301	NAP	C3N-C7N-N7N	3.44	121.88	117.75
2	F	301	NAP	N3A-C2A-N1A	-3.41	123.35	128.68
2	A	301	NAP	N3A-C2A-N1A	-3.40	123.36	128.68
2	С	301	NAP	N3A-C2A-N1A	-3.31	123.51	128.68
2	Е	301	NAP	C3N-C7N-N7N	3.23	121.62	117.75
2	D	301	NAP	C6N-N1N-C2N	-3.19	119.07	121.97
2	С	301	NAP	C6N-N1N-C2N	-2.57	119.64	121.97
2	F	301	NAP	C3D-C2D-C1D	2.55	104.81	100.98
2	D	301	NAP	O2A-PA-O1A	2.50	124.61	112.24
2	В	301	NAP	O2A-PA-O1A	2.48	124.52	112.24
2	A	301	NAP	C3D-C2D-C1D	2.48	104.71	100.98
2	D	301	NAP	O2B-P2B-O1X	-2.47	99.85	109.39
2	D	301	NAP	C4A-C5A-N7A	-2.38	106.92	109.40
2	С	301	NAP	O2A-PA-O1A	2.37	123.96	112.24
2	С	301	NAP	C3D-C2D-C1D	2.32	104.48	100.98
2	A	301	NAP	C4A-C5A-N7A	-2.32	106.99	109.40
2	С	301	NAP	C4A-C5A-N7A	-2.26	107.04	109.40
2	Е	301	NAP	O7N-C7N-N7N	-2.21	119.44	122.58
2	D	301	NAP	C3D-C2D-C1D	2.20	104.30	100.98
2	F	301	NAP	C4A-C5A-N7A	-2.19	107.11	109.40
2	В	301	NAP	C3N-C2N-N1N	2.17	122.55	120.43
2	Е	301	NAP	O2A-PA-O1A	2.15	122.89	112.24
2	A	301	NAP	O3X-P2B-O2X	2.13	115.79	107.64
2	D	301	NAP	C3N-C7N-N7N	2.09	120.26	117.75
2	F	301	NAP	O4B-C4B-C5B	2.09	116.23	109.37
2	D	301	NAP	C3N-C2N-N1N	2.08	122.46	120.43
2	В	301	NAP	O7N-C7N-N7N	-2.05	119.66	122.58
2	С	301	NAP	C3N-C7N-N7N	2.03	120.18	117.75
2	Е	301	NAP	C2A-N1A-C6A	2.02	122.21	118.75

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	NAP	C5D-O5D-PN-O2N
2	A	301	NAP	O4D-C4D-C5D-O5D
2	В	301	NAP	O4D-C1D-N1N-C6N
2	D	301	NAP	C2B-O2B-P2B-O3X



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	D	301	NAP	O4D-C1D-N1N-C6N
2	F	301	NAP	C2B-O2B-P2B-O1X
2	F	301	NAP	O4D-C1D-N1N-C6N
2	A	301	NAP	C3B-C4B-C5B-O5B
2	A	301	NAP	C3D-C4D-C5D-O5D
2	С	301	NAP	C3B-C4B-C5B-O5B
2	В	301	NAP	C1B-C2B-O2B-P2B
2	F	301	NAP	C1B-C2B-O2B-P2B
2	Е	301	NAP	C1B-C2B-O2B-P2B
2	A	301	NAP	O4B-C4B-C5B-O5B
2	F	301	NAP	C4N-C3N-C7N-O7N
2	D	301	NAP	C2N-C3N-C7N-O7N
2	F	301	NAP	C2N-C3N-C7N-O7N
2	F	301	NAP	C4N-C3N-C7N-N7N
2	D	301	NAP	C4N-C3N-C7N-O7N
2	С	301	NAP	O4B-C4B-C5B-O5B
2	F	301	NAP	C3B-C2B-O2B-P2B
2	D	301	NAP	C4N-C3N-C7N-N7N
2	F	301	NAP	C2N-C3N-C7N-N7N
2	D	301	NAP	C2N-C3N-C7N-N7N
2	A	301	NAP	C2B-O2B-P2B-O2X
2	A	301	NAP	C5D-O5D-PN-O3
2	A	301	NAP	C5D-O5D-PN-O1N
2	D	301	NAP	C3B-C4B-C5B-O5B
2	Е	301	NAP	C3B-C2B-O2B-P2B
2	A	301	NAP	C4N-C3N-C7N-O7N
2	F	301	NAP	C3B-C4B-C5B-O5B
2	A	301	NAP	C4N-C3N-C7N-N7N
2	В	301	NAP	C3B-C2B-O2B-P2B
2	F	301	NAP	O4B-C4B-C5B-O5B
2	A	301	NAP	C2N-C3N-C7N-O7N
2	A	301	NAP	C2N-C3N-C7N-N7N
2	В	301	NAP	PN-O3-PA-O1A
2	F	301	NAP	PN-O3-PA-O2A
2	С	301	NAP	C4N-C3N-C7N-O7N
2	F	301	NAP	C5D-O5D-PN-O1N

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAP	1	0




Continued from previous page...


\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	NAP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	238/240 (99%)	-0.13	3 (1%) 77 81	28, 42, 69, 89	0
1	В	224/240 (93%)	-0.17	2 (0%) 84 87	29, 40, 59, 108	0
1	С	224/240 (93%)	0.38	12 (5%) 25 31	39, 60, 95, 113	0
1	D	223/240 (92%)	0.40	14 (6%) 20 24	39, 59, 92, 110	0
1	E	223/240 (92%)	-0.16	0 100 100	29, 40, 58, 68	0
1	F	238/240 (99%)	-0.14	2 (0%) 86 89	28, 42, 69, 97	0
All	All	1370/1440 (95%)	0.03	33 (2%) 59 65	28, 47, 83, 113	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	122	LEU	4.0
1	D	131	THR	3.9
1	D	118	LYS	3.6
1	D	197	PHE	3.4
1	D	121	MET	3.4
1	С	134	PHE	3.3
1	С	120	ARG	3.2
1	В	1	MET	2.9
1	D	134	PHE	2.8
1	С	98	ILE	2.8
1	С	112	ASN	2.8
1	С	118	LYS	2.8
1	F	228	ASN	2.7
1	D	120	ARG	2.6
1	D	157	GLY	2.6
1	D	110	VAL	2.6
1	С	225	GLY	2.5
1	С	124	GLU	2.4
1	A	-9	GLY	2.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	117	GLU	2.4
1	С	183	ILE	2.4
1	F	134	PHE	2.4
1	D	57	ASP	2.3
1	С	130	LYS	2.3
1	D	144	LYS	2.3
1	D	86	VAL	2.2
1	С	94	LEU	2.2
1	A	34	PHE	2.2
1	D	111	SER	2.2
1	D	129	ILE	2.1
1	A	120	ARG	2.1
1	С	133	ALA	2.1
1	В	34	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q < 0.9
2	NAP	A	301	48/48	0.65	0.39	99,125,136,137	0
2	NAP	F	301	48/48	0.74	0.41	91,130,140,141	0
2	NAP	D	301	48/48	0.91	0.18	57,73,101,104	0
2	NAP	С	301	48/48	0.91	0.18	57,74,95,96	0
2	NAP	Ε	301	48/48	0.98	0.12	31,34,46,49	0
2	NAP	В	301	48/48	0.98	0.11	30,36,46,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers




as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAP D 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP C 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAP E 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP B 301:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

