



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 05:03 AM EDT

PDB ID : 6XUQ  
Title : Human Ecto-5'-nucleotidase (CD73) in complex with A1618 (compound 1b in publication) in the closed state in crystal form III  
Authors : Strater, N.  
Deposited on : 2020-01-20  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-nucleotidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	514	4003	2546	680	759	18	0	1	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	initiating methionine	UNP P21589
A	9	GLU	-	expression tag	UNP P21589
A	10	TRP	-	expression tag	UNP P21589
A	11	SER	-	expression tag	UNP P21589
A	12	TRP	-	expression tag	UNP P21589
A	13	VAL	-	expression tag	UNP P21589
A	14	PHE	-	expression tag	UNP P21589
A	15	LEU	-	expression tag	UNP P21589
A	16	PHE	-	expression tag	UNP P21589
A	17	PHE	-	expression tag	UNP P21589
A	18	LEU	-	expression tag	UNP P21589
A	19	SER	-	expression tag	UNP P21589
A	20	VAL	-	expression tag	UNP P21589
A	21	THR	-	expression tag	UNP P21589
A	22	THR	-	expression tag	UNP P21589
A	23	GLY	-	expression tag	UNP P21589
A	24	VAL	-	expression tag	UNP P21589
A	25	HIS	-	expression tag	UNP P21589
A	26	SER	-	expression tag	UNP P21589
A	53	ASP	ASN	conflict	UNP P21589
A	311	ASP	ASN	conflict	UNP P21589
A	333	ASP	ASN	conflict	UNP P21589
A	403	ASP	ASN	conflict	UNP P21589
A	550	GLY	-	expression tag	UNP P21589
A	551	GLY	-	expression tag	UNP P21589
A	552	GLY	-	expression tag	UNP P21589
A	553	GLY	-	expression tag	UNP P21589

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	554	ALA	-	expression tag	UNP P21589
A	555	GLY	-	expression tag	UNP P21589
A	556	GLY	-	expression tag	UNP P21589
A	557	GLY	-	expression tag	UNP P21589
A	558	GLY	-	expression tag	UNP P21589
A	559	HIS	-	expression tag	UNP P21589
A	560	HIS	-	expression tag	UNP P21589
A	561	HIS	-	expression tag	UNP P21589
A	562	HIS	-	expression tag	UNP P21589
A	563	HIS	-	expression tag	UNP P21589
A	564	HIS	-	expression tag	UNP P21589


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is [[(2 {R},3 {S},4 {R},5 {R})-5-[6-chloranyl-4-[(1 {S})-1-(4-fluorophenyl)ethyl]amino]pyrazolo[3,4-b]pyridin-1-yl]-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]methylphosphonic acid (three-letter code: O1T) (formula: C<sub>20</sub>H<sub>24</sub>ClFN<sub>4</sub>O<sub>9</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	Cl	F	N	O			P
4	A	1	37	20	1	1	4	9	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	218	Total	O	0	2
			218	218		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-nucleotidase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.09Å 93.85Å 235.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.21 – 1.97 27.35 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.21-1.97) 99.9 (27.35-1.97)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 1.96Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.212 , 0.239 0.226 , 0.261	Depositor DCC
$R_{free}$ test set	1102 reflections (2.62%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.7	Xtrriage
Anisotropy	0.322	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 57.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.023 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4261	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, O1T, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/4087	0.72	0/5533

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4003	0	3995	23	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	37	0	0	0	0
5	A	218	0	0	1	0
All	All	4261	0	3995	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:VAL:HG12	1:A:494:LYS:HB3	1.72	0.71
1:A:40:ARG:HD3	1:A:283:PHE:HB3	1.79	0.65
1:A:138:PRO:HG3	1:A:162:LYS:HG2	1.84	0.59
1:A:179:LYS:HD3	1:A:195:ASP:HA	1.87	0.56
1:A:425:GLN:HB2	1:A:517:ARG:HB3	1.90	0.54
1:A:342:THR:HB	1:A:406:ILE:HD11	1.90	0.52
1:A:164:LEU:HD13	1:A:173:ILE:HD12	1.92	0.52
1:A:38:HIS:CE1	1:A:85:ASP:HB3	2.45	0.51
1:A:58:MET:HG2	1:A:311:ASP:HA	1.93	0.51
1:A:488:LYS:HD2	1:A:491:GLU:HG3	1.93	0.50
1:A:175:GLY:HA2	1:A:218:LEU:O	2.13	0.48
1:A:338:GLU:HG3	1:A:405:THR:HG23	1.95	0.48
1:A:356:ARG:HD2	5:A:746:HOH:O	2.16	0.46
1:A:431:LEU:HD21	1:A:495:VAL:HG21	1.99	0.45
1:A:328:ARG:HA	1:A:331:LEU:HD12	1.99	0.45
1:A:35:ASN:HB3	1:A:83:ALA:HB3	1.99	0.44
1:A:205:ASP:O	1:A:209:THR:HG23	2.19	0.43
1:A:31:ILE:HD12	1:A:292:ILE:HD11	2.00	0.42
1:A:87:TYR:CD2	1:A:119:GLU:HA	2.54	0.42
1:A:498:PRO:HD2	1:A:501:LEU:HD12	2.02	0.42
1:A:346:LEU:HD12	1:A:398:ILE:HB	2.02	0.41
1:A:32:LEU:HD21	1:A:71:ILE:HG21	2.03	0.40
1:A:32:LEU:HB3	1:A:287:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	511/557 (92%)	490 (96%)	19 (4%)	2 (0%)	34 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	517	ARG
1	A	88	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	443/475 (93%)	429 (97%)	14 (3%)	39 28

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLU
1	A	45	SER
1	A	125	GLU
1	A	170	VAL
1	A	206	LYS
1	A	211	ASN
1	A	328	ARG
1	A	365	CYS
1	A	403	ASP
1	A	490	ASP
1	A	497	LEU
1	A	516	LEU
1	A	534	LYS
1	A	536	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	O1T	A	604	2	35,40,40	0.57	0	44,61,61	1.03	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	O1T	A	604	2	-	2/20/40/40	0/4/4/4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	604	O1T	C11-N3-C12	3.03	130.52	124.37
4	A	604	O1T	C10-C11-N3	-2.98	117.36	121.96
4	A	604	O1T	C9-N2-C8	-2.53	115.38	116.61

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	604	O1T	C7-C11-N3-C12
4	A	604	O1T	C10-C11-N3-C12

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	514/557 (92%)	1.06	108 (21%) <b>1</b> <b>0</b>	48, 66, 91, 113	7 (1%)

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	VAL	7.6
1	A	431	LEU	6.6
1	A	174	VAL	6.4
1	A	81	LEU	6.0
1	A	34	THR	5.9
1	A	373	LEU	5.5
1	A	402	ASN	5.4
1	A	240	VAL	5.4
1	A	452	VAL	5.3
1	A	388	ILE	5.2
1	A	389	LEU	5.2
1	A	83	ALA	5.1
1	A	218	LEU	5.0
1	A	408	TRP	4.9
1	A	37	VAL	4.6
1	A	403	ASP	4.5
1	A	114	ALA	4.4
1	A	150	LEU	4.2
1	A	450	LEU	4.2
1	A	329	ILE	4.2
1	A	216	ILE	4.1
1	A	284	GLY	4.0
1	A	455	ILE	4.0
1	A	39	SER	4.0
1	A	108	LEU	4.0
1	A	394	ILE	4.0
1	A	287	LEU	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	89	GLY	3.8
1	A	35	ASN	3.7
1	A	330	LYS	3.6
1	A	115	LEU	3.6
1	A	217	ALA	3.6
1	A	404	GLY	3.6
1	A	82	ASP	3.5
1	A	282	ALA	3.5
1	A	502	ALA	3.5
1	A	153	GLN	3.5
1	A	41	LEU	3.5
1	A	393	GLY	3.5
1	A	283	PHE	3.5
1	A	364	ILE	3.4
1	A	384	VAL	3.4
1	A	490	ASP	3.4
1	A	112	ALA	3.4
1	A	113	MET	3.4
1	A	477	THR	3.3
1	A	33	HIS	3.3
1	A	173	ILE	3.3
1	A	280	ALA	3.3
1	A	84	GLY	3.2
1	A	239	VAL	3.2
1	A	278	VAL	3.2
1	A	129	GLU	3.1
1	A	238	VAL	3.1
1	A	40	ARG	2.9
1	A	495	VAL	2.9
1	A	54	ALA	2.8
1	A	32	LEU	2.8
1	A	336	THR	2.8
1	A	149	PRO	2.7
1	A	453	GLY	2.7
1	A	140	LEU	2.7
1	A	360	MET	2.7
1	A	74	ALA	2.7
1	A	387	CYS	2.7
1	A	31	ILE	2.6
1	A	334	TYR	2.6
1	A	55	SER	2.6
1	A	363	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	327	TRP	2.5
1	A	365	CYS	2.5
1	A	401	ARG	2.5
1	A	509	GLN	2.5
1	A	86	GLN	2.5
1	A	337	GLN	2.5
1	A	478	LYS	2.5
1	A	516	LEU	2.5
1	A	520	SER	2.5
1	A	496	ILE	2.5
1	A	219	GLY	2.4
1	A	52	VAL	2.4
1	A	288	GLY	2.4
1	A	519	ASP	2.4
1	A	116	GLY	2.4
1	A	451	GLN	2.4
1	A	413	ALA	2.3
1	A	326	LYS	2.3
1	A	189	THR	2.3
1	A	183	PHE	2.3
1	A	80	LEU	2.3
1	A	461	LEU	2.2
1	A	341	LYS	2.2
1	A	372	ASN	2.2
1	A	417	PHE	2.2
1	A	414	VAL	2.1
1	A	362	ASN	2.1
1	A	501	LEU	2.1
1	A	383	HIS	2.1
1	A	90	THR	2.1
1	A	462	SER	2.1
1	A	258	VAL	2.1
1	A	512	LYS	2.1
1	A	241	GLY	2.1
1	A	242	GLY	2.0
1	A	443	GLY	2.0
1	A	139	ILE	2.0
1	A	252	ASN	2.0
1	A	546	ILE	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	O1T	A	604	37/37	0.92	0.18	55,66,92,93	0
3	CA	A	603	1/1	0.97	0.05	86,86,86,86	0
2	ZN	A	601	1/1	0.98	0.03	68,68,68,68	0
2	ZN	A	602	1/1	0.98	0.10	58,58,58,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.