



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 10:14 PM BST

PDB ID : 6Y3Z
Title : Crystal structure of the Pby1 ATP-grasp enzyme bound to the *S. cerevisiae* mRNA decapping complex (Dcp1-Dcp2-Edc3)
Authors : Graille, M.
Deposited on : 2020-02-19
Resolution : 3.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	
2	B	236	
3	C	71	
4	P	429	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called m7GpppN-mRNA hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	243	2044	1323	338	374	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	272	HIS	-	expression tag	UNP P53550
A	273	HIS	-	expression tag	UNP P53550
A	274	HIS	-	expression tag	UNP P53550
A	275	HIS	-	expression tag	UNP P53550
A	276	HIS	-	expression tag	UNP P53550
A	277	HIS	-	expression tag	UNP P53550

- Molecule 2 is a protein called mRNA-decapping enzyme subunit 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	18	159	104	28	27	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP Q12517
B	-3	PRO	-	expression tag	UNP Q12517
B	-2	LEU	-	expression tag	UNP Q12517
B	-1	GLY	-	expression tag	UNP Q12517
B	0	SER	-	expression tag	UNP Q12517

- Molecule 3 is a protein called Enhancer of mRNA-decapping protein 3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	63	470	299	84	87	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	GLY	-	expression tag	UNP P39998
C	-3	PRO	-	expression tag	UNP P39998
C	-2	LEU	-	expression tag	UNP P39998
C	-1	GLY	-	expression tag	UNP P39998
C	0	SER	-	expression tag	UNP P39998

- Molecule 4 is a protein called Probable tubulin-tyrosine ligase PB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	P	348	2896	1868	461	558	9	0	0	0

There are 5 discrepancies between the modelled and reference sequences:





Chain	Residue	Modelled	Actual	Comment	Reference
P	325	GLY	-	expression tag	UNP P38254
P	326	PRO	-	expression tag	UNP P38254
P	327	LEU	-	expression tag	UNP P38254
P	328	GLY	-	expression tag	UNP P38254
P	329	SER	-	expression tag	UNP P38254

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	P	1	1	1	0	0



● Molecule 4: Probable tubulin-tyrosine ligase PBY1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.21Å 90.60Å 194.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.50 – 3.49 48.50 – 3.49	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.50-3.49) 99.5 (48.50-3.49)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 3.48Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.246 , 0.321 0.274 , 0.368	Depositor DCC
R_{free} test set	767 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	145.9	Xtrriage
Anisotropy	0.024	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 181.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5570	wwPDB-VP
Average B, all atoms (Å ²)	193.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/2086	0.71	0/2804
2	B	0.42	0/162	0.61	0/216
3	C	0.43	0/473	0.79	0/629
4	P	0.54	0/2961	0.83	0/4011
All	All	0.50	0/5682	0.78	0/7660

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	2050	38	0
2	B	159	0	154	2	0
3	C	470	0	503	18	0
4	P	2896	0	2804	77	0
5	P	1	0	0	0	0
All	All	5570	0	5511	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:P:585:LEU:HD23	4:P:627:VAL:HG12	1.53	0.90
3:C:19:ILE:CG2	3:C:36:VAL:CG2	2.57	0.82
3:C:19:ILE:CG2	3:C:36:VAL:HG22	2.10	0.81
1:A:252:GLU:HG2	3:C:59:VAL:HG12	1.62	0.81
1:A:88:ARG:HB3	1:A:91:GLU:HB3	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/277 (85%)	214 (91%)	17 (7%)	4 (2%)	9	42
2	B	16/236 (7%)	15 (94%)	1 (6%)	0	100	100
3	C	61/71 (86%)	53 (87%)	8 (13%)	0	100	100
4	P	338/429 (79%)	295 (87%)	38 (11%)	5 (2%)	10	45
All	All	650/1013 (64%)	577 (89%)	64 (10%)	9 (1%)	11	46

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	P	704	GLY
1	A	158	LEU
1	A	214	MET
4	P	346	ILE
4	P	505	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/262 (88%)	180 (78%)	50 (22%)	1	5
2	B	16/211 (8%)	11 (69%)	5 (31%)	0	2
3	C	50/56 (89%)	38 (76%)	12 (24%)	0	4
4	P	327/398 (82%)	250 (76%)	77 (24%)	1	4
All	All	623/927 (67%)	479 (77%)	144 (23%)	1	4

5 of 144 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	56	ASP
4	P	395	LEU
4	P	710	LEU
4	P	335	LEU
4	P	360	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
4	P	383	HIS
4	P	732	ASN
4	P	400	ASN
1	A	254	GLN
4	P	672	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	243/277 (87%)	0.64	35 (14%) 2 3	136, 231, 289, 296	0
2	B	18/236 (7%)	0.54	1 (5%) 24 22	254, 263, 276, 279	0
3	C	63/71 (88%)	0.26	6 (9%) 8 8	172, 214, 255, 264	0
4	P	348/429 (81%)	0.21	20 (5%) 23 21	90, 157, 219, 270	0
All	All	672/1013 (66%)	0.38	62 (9%) 9 9	90, 188, 279, 296	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	PHE	5.4
1	A	61	PRO	5.3
1	A	142	ASP	5.1
1	A	68	ILE	4.9
1	A	75	ILE	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MG	P	801	1/1	0.88	0.20	128,128,128,128	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.