

wwPDB EM Validation Summary Report (i)

Jun 26, 2023 – 02:11 PM JST

:	7Y35
:	EMD-33588
:	Cryo-EM structure of the Abaloparatide-bound human PTH1R-Gs complex
:	Zhai, X.; Mao, C.; Shen, Q.; Zang, S.; Shen, D.; Zhang, H.; Chen, Z.; Wang,
	G.; Zhang, C.; Zhang, Y.; Liu, Z.
:	2022-06-09
:	2.90 Å(reported)
	: : :

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


EMDB validation analysis	:	0.0.1. dev 50
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive}\ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length		Quality of cha	in
1	R	727	44%	7%	49%
2	Р	34	65%	,	35%
3	А	380	52%	7%	41%
4	В	380		76%	13% 11%
5	G	70		74%	9% 17%
6	N	126		88%	12%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Parathyroid hormone/parathyroid hormone-related peptide receptor.

Mol	Chain	Residues	Atoms				AltConf	Trace	
1	R	371	Total 3053	C 2016	N 512	O 505	S 20	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	188	ALA	GLY	conflict	UNP Q03431

• Molecule 2 is a protein called Abaloparatide.

Mol	Chain	Residues	Atoms			AltConf	Trace	
2	Р	34	Total 274	C 171	N 55	O 48	0	1

 \bullet Molecule 3 is a protein called Isoform Gnas-2 of Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms				AltConf	Trace	
3	А	224	Total 1858	C 1173	N 336	0 343	S 6	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	226	ALA	GLY	engineered mutation	UNP P63092-2
А	268	ALA	GLU	engineered mutation	UNP P63092-2
А	271	LYS	ASN	engineered mutation	UNP P63092-2
А	274	ASP	LYS	engineered mutation	UNP P63092-2
А	280	LYS	ARG	engineered mutation	UNP P63092-2
А	284	ASP	THR	engineered mutation	UNP P63092-2
А	285	THR	ILE	engineered mutation	UNP P63092-2



- Molecule 4 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
4	В	338	Total 2600	C 1604	N 467	O 508	S 21	0	0

There are 41 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-13	MET	-	initiating methionine	UNP P54311
В	-12	HIS	-	expression tag	UNP P54311
В	-11	HIS	-	expression tag	UNP P54311
В	-10	HIS	-	expression tag	UNP P54311
В	-9	HIS	-	expression tag	UNP P54311
В	-8	HIS	-	expression tag	UNP P54311
В	-7	HIS	-	expression tag	UNP P54311
В	-6	LEU	-	expression tag	UNP P54311
В	-5	GLU	-	expression tag	UNP P54311
В	-4	VAL	-	expression tag	UNP P54311
В	-3	LEU	-	expression tag	UNP P54311
В	-2	PHE	-	expression tag	UNP P54311
В	-1	GLN	-	expression tag	UNP P54311
В	0	GLY	-	expression tag	UNP P54311
В	1	PRO	-	expression tag	UNP P54311
В	341	GLY	-	expression tag	UNP P54311
В	342	SER	-	expression tag	UNP P54311
В	343	SER	-	expression tag	UNP P54311
В	344	GLY	-	expression tag	UNP P54311
В	345	GLY	-	expression tag	UNP P54311
В	346	GLY	-	expression tag	UNP P54311
В	347	GLY	-	expression tag	UNP P54311
В	348	SER	-	expression tag	UNP P54311
В	349	GLY	-	expression tag	UNP P54311
В	350	GLY	-	expression tag	UNP P54311
В	351	GLY	-	expression tag	UNP P54311
В	352	GLY	-	expression tag	UNP P54311
В	353	SER	-	expression tag	UNP P54311
В	354	SER	-	expression tag	UNP P54311
В	355	GLY	-	expression tag	UNP P54311
В	356	VAL	-	expression tag	UNP P54311
В	357	SER	-	expression tag	UNP P54311
В	358	GLY	-	expression tag	UNP P54311
В	359	TRP	-	expression tag	UNP P54311
В	360	ARG	_	expression tag	UNP P54311

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
В	361	LEU	-	expression tag	UNP P54311
В	362	PHE	-	expression tag	UNP P54311
В	363	LYS	-	expression tag	UNP P54311
В	364	LYS	-	expression tag	UNP P54311
В	365	ILE	-	expression tag	UNP P54311
В	366	SER	-	expression tag	UNP P54311

Continued from previous page...

- Molecule 5 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms				AltConf	Trace	
5	G	58	Total 445	C 278	N 78	O 86	${ m S} { m 3}$	0	0

• Molecule 6 is a protein called NanoBody 35.


Mol	Chain	Residues	Atoms				AltConf	Trace	
6	Ν	126	Total 961	C 599	N 168	0 188	S 6	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Parathyroid hormone/parathyroid hormone-related peptide receptor



• Molecule 3: Isoform Gnas-2 of Guanine nucleotide-binding protein G(s) subunit alpha isoforms short





• Molecule 6: NanoBody 35

Chain N:	88%	12%
q1 F29 V37 V37 V37 F29 F10 F10 F10 F10 F10 F10 F10 F100 F107 C107	Y115 1125 V126	



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	456840	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	64	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	49310	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.772	Depositor
Minimum map value	-0.003	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.038	Depositor
Recommended contour level	0.05	Depositor
Map size (Å)	227.13602, 227.13602, 227.13602	wwPDB
Map dimensions	224, 224, 224	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.014, 1.014, 1.014	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AIB, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	ond angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	R	0.56	0/3143	0.52	3/4269~(0.1%)
2	Р	0.71	0/269	0.40	0/357
3	А	0.45	0/1892	0.47	0/2543
4	В	0.47	0/2647	0.56	0/3589
5	G	0.34	0/451	0.45	0/609
6	Ν	0.55	0/981	0.52	0/1329
All	All	0.51	0/9383	0.52	3/12696~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	R	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	R	172	LYS	CA-C-N	-7.64	100.39	117.20
1	R	172	LYS	O-C-N	6.54	133.17	122.70
1	R	172	LYS	C-N-CA	5.84	136.29	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	R	172	LYS	Mainchain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	3053	0	3046	34	0
2	Р	274	0	296	10	0
3	А	1858	0	1835	18	0
4	В	2600	0	2505	28	0
5	G	445	0	454	5	0
6	N	961	0	928	10	0
All	All	9191	0	9064	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:444:GLU:OE1	2:P:3:SER:HB2	1.73	0.86
1:R:448:ASN:ND2	2:P:4:GLU:OE2	2.22	0.70
4:B:49:ARG:NH1	5:G:61:PHE:O	2.23	0.69
1:R:422:ILE:HA	1:R:425:MET:HG2	1.75	0.67
3:A:27:GLU:OE2	4:B:89:LYS:NZ	2.31	0.64

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	363/727~(50%)	348 (96%)	15 (4%)	0	100	100

Continued on next page...



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
2	Р	31/34~(91%)	30~(97%)	1 (3%)	0	100	100
3	А	216/380~(57%)	207~(96%)	9~(4%)	0	100	100
4	В	336/380~(88%)	321 (96%)	15~(4%)	0	100	100
5	G	56/70~(80%)	56 (100%)	0	0	100	100
6	Ν	124/126~(98%)	118 (95%)	6~(5%)	0	100	100
All	All	1126/1717~(66%)	1080 (96%)	46 (4%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	R	325/620~(52%)	324 (100%)	1 (0%)	92	98
2	Р	30/30~(100%)	30 (100%)	0	100	100
3	А	200/340~(59%)	198~(99%)	2(1%)	76	92
4	В	281/311 (90%)	278~(99%)	3 (1%)	73	92
5	G	47/57~(82%)	47 (100%)	0	100	100
6	Ν	104/104 (100%)	104 (100%)	0	100	100
All	All	987/1462 (68%)	981 (99%)	6 (1%)	86	96

5 of 6 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
4	В	23	LYS
4	В	105	TYR
4	В	234	PHE
3	А	265	ARG
1	R	183	VAL

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such side chains are listed below:



Mol	Chain	Res	Type
1	R	402	GLN
1	R	420	HIS
2	Р	9	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	gths	В	ond ang	gles
WIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	AIB	Р	29	2	$1,\!5,\!6$	1.20	0	2,7,9	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AIB	Р	29	2	-	0/2/3/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Р	29	AIB	1	0



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-33588. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map



X Index: 112




Y Index: 112



Z Index: 112

6.2.2 Raw map



X Index: 112

Y Index: 112

Z Index: 112

The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 122




Y Index: 124



Z Index: 148

6.3.2 Raw map



X Index: 122

Y Index: 125




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 102 nm^3 ; this corresponds to an approximate mass of 92 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.345 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.345 $\mathrm{\AA^{-1}}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estim	Estimation criterion (FSC cut-of				
Resolution estimate (A)	0.143	0.5	Half-bit			
Reported by author	2.90	-	-			
Author-provided FSC curve	-	-	-			
Unmasked-calculated*	3.14	3.66	3.21			


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-33588 and PDB model 7Y35. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).



9.4 Atom inclusion (i)



At the recommended contour level, 99% of all backbone atoms, 94% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9370	0.4790
А	0.9250	0.4860
В	0.9640	0.5390
G	0.8970	0.4330
Ν	0.9600	0.5440
Р	0.9100	0.4050
R	0.9220	0.4180

