



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 4, 2024 – 07:12 pm GMT

PDB ID : 4Y77  
Title : Yeast 20S proteasome in complex with Ac-LAF-ep  
Authors : Huber, E.M.; Groll, M.  
Deposited on : 2015-02-13  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 5% 98% 6% 5%
1	O	250	 4% 97% 6% 5%
2	B	258	 6% 89% 6% 5%
2	P	258	 6% 89% 5% 6%
3	C	254	 8% 87% 6% 6%





*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	12% 87% 6% • 6%
4	D	260	2% 86% • 10%
4	R	260	4% 85% 5% 10%
5	E	234	4% 92% 6% •
5	S	234	6% 91% 7% •
6	F	288	4% 80% 5% 16%
6	T	288	5% 80% 5% 16%
7	G	252	4% 90% 6% •
7	U	252	3% 90% 6% •
8	H	232	2% 91% • •
8	V	232	3% 91% • •
9	I	205	% 92% 7%
9	W	205	% 93% 5% •
10	J	198	2% 90% 7% • •
10	X	198	2% 91% 6% • •
11	K	212	95% 5%
11	Y	212	% 95% 5%
12	L	222	95% 5%
12	Z	222	% 95% 5%
13	M	246	2% 86% • • 9%
13	a	246	2% 88% • 9%
14	N	196	2% 95% 5%
14	b	196	% 98% •
15	c	5	80% 20%
15	d	5	80% 20%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
15	e	5	 80% 20%
15	f	5	 80% 20%
15	g	5	 80% 20%
15	h	5	 80% 20%

## 2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 50341 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total 1915	C 1219	N 315	O 377	S 4	0	0	0
1	O	250	Total 1915	C 1219	N 315	O 377	S 4	0	0	0

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	Total 1904	C 1201	N 321	O 379	S 3	0	0	0
2	P	244	Total 1904	C 1201	N 321	O 379	S 3	0	0	0

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	240	Total 1881	C 1176	N 329	O 372	S 4	0	0	0
3	Q	240	Total 1881	C 1176	N 329	O 372	S 4	0	0	0

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	235	Total 1813	C 1136	N 304	O 366	S 7	0	0	0
4	R	235	Total 1813	C 1136	N 304	O 366	S 7	0	0	0

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			
5	S	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			
6	T	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			
7	U	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			
8	V	222	Total	C	N	O	S	0	1	0
			1691	1066	295	323	7			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	224	Total	C	N	O	S	0	0	0
			1753	1108	300	338	7			
13	a	223	Total	C	N	O	S	0	0	0
			1745	1104	299	335	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is a protein called Ac-LAF-ep.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	c	5	Total	C	N	O	0	0	0
			31	23	3	5			
15	d	5	Total	C	N	O	0	0	0
			31	23	3	5			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	e	5	Total 31	C 23	N 3	O 5	0	0	0
15	f	5	Total 31	C 23	N 3	O 5	0	0	0
15	g	5	Total 31	C 23	N 3	O 5	0	0	0
15	h	5	Total 31	C 23	N 3	O 5	0	0	0

- Molecule 16 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	G	1	Total 1	Mg 1	0	0
16	I	2	Total 2	Mg 2	0	0
16	K	1	Total 1	Mg 1	0	0
16	L	1	Total 1	Mg 1	0	0
16	N	1	Total 1	Mg 1	0	0
16	Z	1	Total 1	Mg 1	0	0

- Molecule 17 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	G	1	Total 1	Cl 1	0	0
17	U	1	Total 1	Cl 1	0	0

- Molecule 18 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
18	K	1	12	6	1	4	1	0	0
18	Y	1	12	6	1	4	1	0	0

- Molecule 19 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
19	A	43	Total	O	0	0
			43	43		
19	B	33	Total	O	0	0
			33	33		
19	C	24	Total	O	0	0
			24	24		
19	D	24	Total	O	0	0
			24	24		
19	E	15	Total	O	0	0
			15	15		
19	F	32	Total	O	0	0
			32	32		
19	G	43	Total	O	0	0
			43	43		
19	H	40	Total	O	0	0
			40	40		
19	I	49	Total	O	0	0
			49	49		
19	J	42	Total	O	0	0
			42	42		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
19	K	57	Total O 57 57	0	0
19	L	42	Total O 42 42	0	0
19	M	30	Total O 30 30	0	0
19	N	29	Total O 29 29	0	0
19	O	26	Total O 26 26	0	0
19	P	23	Total O 23 23	0	0
19	Q	13	Total O 13 13	0	0
19	R	18	Total O 18 18	0	0
19	S	13	Total O 13 13	0	0
19	T	24	Total O 24 24	0	0
19	U	36	Total O 36 36	0	0
19	V	34	Total O 34 34	0	0
19	W	40	Total O 40 40	0	0
19	X	47	Total O 47 47	0	0
19	Y	50	Total O 50 50	0	0
19	Z	47	Total O 47 47	0	0
19	a	47	Total O 47 47	0	0
19	b	38	Total O 38 38	0	0
19	c	3	Total O 3 3	0	0
19	d	1	Total O 1 1	0	0
19	e	1	Total O 1 1	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
19	f	1	Total O 1 1	0	0
19	g	2	Total O 2 2	0	0
19	h	2	Total O 2 2	0	0



GLN  
ASP  
LYS  
LYS  
LYS  
LYS  
SER  
ASN  
HIS

- Molecule 3: Proteasome subunit alpha type-4

Chain Q: 12% 87% 6% 6%




- Molecule 4: Proteasome subunit alpha type-5

Chain D: 2% 86% 10%



- Molecule 4: Proteasome subunit alpha type-5

Chain R: 4% 85% 5% 10%



ALA  
ASP  
VAL  
GLU  
MET  
SER

- Molecule 5: Proteasome subunit alpha type-6

Chain E: 4% 92% 6%



- Molecule 5: Proteasome subunit alpha type-6

Chain S: 6% 91% 7%



- Molecule 6: Probable proteasome subunit alpha type-7

Chain F: 4% 80% 5% 16%








- Molecule 9: Proteasome subunit beta type-3



- Molecule 10: Proteasome subunit beta type-4



- Molecule 10: Proteasome subunit beta type-4



- Molecule 11: Proteasome subunit beta type-5




- Molecule 11: Proteasome subunit beta type-5





- Molecule 12: Proteasome subunit beta type-6




- Molecule 12: Proteasome subunit beta type-6




• Molecule 13: Proteasome subunit beta type-7




• Molecule 13: Proteasome subunit beta type-7




• Molecule 14: Proteasome subunit beta type-1




• Molecule 14: Proteasome subunit beta type-1



• Molecule 15: Ac-LAF-ep




• Molecule 15: Ac-LAF-ep




• Molecule 15: Ac-LAF-ep



Chain e:  80% 20%




• Molecule 15: Ac-LAF-ep

Chain f:  80% 20%




• Molecule 15: Ac-LAF-ep

Chain g:  80% 20%



• Molecule 15: Ac-LAF-ep

Chain h:  80% 20%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.45Å 300.30Å 145.22Å 90.00° 112.90° 90.00°	Depositor
Resolution (Å)	15.00 – 2.50 15.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.2 (15.00-2.50) 98.2 (15.00-2.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.47 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.199 , 0.215 0.202 , 0.217	Depositor DCC
$R_{free}$ test set	17933 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.2	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 35.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	50341	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: POL, MG, CL, PHL, ACE, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1952	0.48	0/2642
1	O	0.27	0/1952	0.47	0/2642
2	B	0.27	0/1934	0.50	0/2618
2	P	0.27	0/1934	0.49	0/2618
3	C	0.28	0/1910	0.51	0/2586
3	Q	0.27	0/1910	0.51	0/2586
4	D	0.27	0/1837	0.48	0/2475
4	R	0.27	0/1837	0.48	0/2475
5	E	0.27	0/1800	0.48	0/2433
5	S	0.27	0/1800	0.48	0/2433
6	F	0.27	0/1932	0.46	0/2609
6	T	0.27	0/1932	0.46	0/2609
7	G	0.27	0/1945	0.48	0/2634
7	U	0.27	0/1945	0.48	0/2634
8	H	0.35	0/1715	0.51	0/2326
8	V	0.29	0/1726	0.49	0/2341
9	I	0.27	0/1611	0.48	0/2174
9	W	0.28	0/1611	0.49	0/2174
10	J	0.26	0/1589	0.49	0/2142
10	X	0.26	0/1589	0.49	0/2142
11	K	0.29	0/1681	0.52	1/2274 (0.0%)
11	Y	0.29	0/1681	0.52	1/2274 (0.0%)
12	L	0.27	0/1795	0.48	0/2420
12	Z	0.32	0/1795	0.48	0/2420
13	M	0.29	0/1783	0.52	0/2420
13	a	0.29	0/1775	0.52	0/2409
14	N	0.33	0/1541	0.51	1/2087 (0.0%)
14	b	0.33	0/1541	0.50	0/2087
15	c	0.78	0/13	1.33	0/17
15	d	1.14	0/13	1.11	0/17
15	e	1.17	0/13	1.37	0/17
15	f	0.66	0/13	1.26	0/17

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
15	g	1.15	0/13	1.19	0/17
15	h	0.98	0/13	1.12	0/17
All	All	0.29	0/50131	0.49	3/67786 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
11	Y	1	THR	N-CA-C	5.88	126.87	111.00
11	K	1	THR	N-CA-C	5.40	125.57	111.00
14	N	1	THR	N-CA-C	5.34	125.42	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	2	0
1	O	1915	0	1929	3	0
2	B	1904	0	1904	5	0
2	P	1904	0	1904	5	0
3	C	1881	0	1895	9	0
3	Q	1881	0	1895	9	0
4	D	1813	0	1797	3	0
4	R	1813	0	1797	5	0
5	E	1773	0	1775	4	0
5	S	1773	0	1775	6	0
6	F	1892	0	1883	3	0
6	T	1892	0	1883	2	0
7	G	1907	0	1901	4	0
7	U	1907	0	1901	4	0
8	H	1684	0	1685	3	0
8	V	1691	0	1692	4	0
9	I	1581	0	1574	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	W	1581	0	1574	9	0
10	J	1561	0	1569	10	0
10	X	1561	0	1569	8	0
11	K	1644	0	1592	5	0
11	Y	1644	0	1592	5	0
12	L	1757	0	1711	3	0
12	Z	1757	0	1711	3	0
13	M	1753	0	1754	4	0
13	a	1745	0	1750	0	0
14	N	1512	0	1478	2	0
14	b	1512	0	1478	0	0
15	c	31	0	34	0	0
15	d	31	0	33	0	0
15	e	31	0	34	0	0
15	f	31	0	34	0	0
15	g	31	0	33	0	0
15	h	31	0	34	0	0
16	G	1	0	0	0	0
16	I	2	0	0	0	0
16	K	1	0	0	0	0
16	L	1	0	0	0	0
16	N	1	0	0	0	0
16	Z	1	0	0	0	0
17	G	1	0	0	0	0
17	U	1	0	0	0	0
18	K	12	0	13	0	0
18	Y	12	0	13	0	0
19	A	43	0	0	0	0
19	B	33	0	0	0	0
19	C	24	0	0	0	0
19	D	24	0	0	0	0
19	E	15	0	0	0	0
19	F	32	0	0	1	0
19	G	43	0	0	0	0
19	H	40	0	0	0	0
19	I	49	0	0	0	0
19	J	42	0	0	1	0
19	K	57	0	0	0	0
19	L	42	0	0	0	0
19	M	30	0	0	0	0
19	N	29	0	0	0	0
19	O	26	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
19	P	23	0	0	0	0
19	Q	13	0	0	0	0
19	R	18	0	0	0	0
19	S	13	0	0	0	0
19	T	24	0	0	0	0
19	U	36	0	0	0	0
19	V	34	0	0	0	0
19	W	40	0	0	0	0
19	X	47	0	0	0	0
19	Y	50	0	0	0	0
19	Z	47	0	0	0	0
19	a	47	0	0	0	0
19	b	38	0	0	0	0
19	c	3	0	0	0	0
19	d	1	0	0	0	0
19	e	1	0	0	0	0
19	f	1	0	0	0	0
19	g	2	0	0	0	0
19	h	2	0	0	0	0
All	All	50341	0	49125	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:J:1:MET:O	10:J:2:ASP:HB2	1.85	0.75
10:X:1:MET:O	10:X:2:ASP:HB2	1.86	0.75
8:V:50:ALA:HB3	9:W:126:ILE:HD12	1.73	0.69
12:Z:13:LEU:HD13	12:Z:150:LEU:HD21	1.77	0.67
12:L:13:LEU:HD13	12:L:150:LEU:HD21	1.76	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	242 (98%)	6 (2%)	0	100	100
1	O	248/250 (99%)	242 (98%)	6 (2%)	0	100	100
2	B	242/258 (94%)	234 (97%)	7 (3%)	1 (0%)	34	54
2	P	242/258 (94%)	234 (97%)	7 (3%)	1 (0%)	34	54
3	C	238/254 (94%)	230 (97%)	5 (2%)	3 (1%)	12	21
3	Q	238/254 (94%)	230 (97%)	5 (2%)	3 (1%)	12	21
4	D	231/260 (89%)	227 (98%)	4 (2%)	0	100	100
4	R	231/260 (89%)	227 (98%)	4 (2%)	0	100	100
5	E	229/234 (98%)	224 (98%)	5 (2%)	0	100	100
5	S	229/234 (98%)	224 (98%)	5 (2%)	0	100	100
6	F	241/288 (84%)	234 (97%)	7 (3%)	0	100	100
6	T	241/288 (84%)	234 (97%)	7 (3%)	0	100	100
7	G	239/252 (95%)	235 (98%)	4 (2%)	0	100	100
7	U	239/252 (95%)	235 (98%)	4 (2%)	0	100	100
8	H	220/232 (95%)	213 (97%)	7 (3%)	0	100	100
8	V	221/232 (95%)	214 (97%)	7 (3%)	0	100	100
9	I	202/205 (98%)	195 (96%)	7 (4%)	0	100	100
9	W	202/205 (98%)	195 (96%)	7 (4%)	0	100	100
10	J	193/198 (98%)	189 (98%)	3 (2%)	1 (0%)	29	48
10	X	193/198 (98%)	189 (98%)	3 (2%)	1 (0%)	29	48
11	K	210/212 (99%)	207 (99%)	3 (1%)	0	100	100
11	Y	210/212 (99%)	207 (99%)	3 (1%)	0	100	100
12	L	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
12	Z	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
13	M	222/246 (90%)	216 (97%)	6 (3%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
13	a	221/246 (90%)	216 (98%)	5 (2%)	0	100	100
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
14	b	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
15	c	2/5 (40%)	2 (100%)	0	0	100	100
15	d	2/5 (40%)	2 (100%)	0	0	100	100
15	e	2/5 (40%)	2 (100%)	0	0	100	100
15	f	2/5 (40%)	2 (100%)	0	0	100	100
15	g	2/5 (40%)	2 (100%)	0	0	100	100
15	h	2/5 (40%)	2 (100%)	0	0	100	100
All	All	6270/6644 (94%)	6115 (98%)	145 (2%)	10 (0%)	47	68

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	51	VAL
3	C	202	GLN
10	J	2	ASP
2	P	51	VAL
3	Q	202	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	206 (99%)	3 (1%)	67	86
1	O	209/209 (100%)	206 (99%)	3 (1%)	67	86
2	B	203/216 (94%)	194 (96%)	9 (4%)	28	52
2	P	203/216 (94%)	194 (96%)	9 (4%)	28	52
3	C	212/226 (94%)	204 (96%)	8 (4%)	33	58
3	Q	212/226 (94%)	204 (96%)	8 (4%)	33	58
4	D	194/215 (90%)	185 (95%)	9 (5%)	27	50

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	R	194/215 (90%)	185 (95%)	9 (5%)	27	50
5	E	190/193 (98%)	181 (95%)	9 (5%)	26	49
5	S	190/193 (98%)	181 (95%)	9 (5%)	26	49
6	F	201/239 (84%)	190 (94%)	11 (6%)	21	41
6	T	201/239 (84%)	190 (94%)	11 (6%)	21	41
7	G	206/210 (98%)	198 (96%)	8 (4%)	32	57
7	U	206/210 (98%)	198 (96%)	8 (4%)	32	57
8	H	181/190 (95%)	174 (96%)	7 (4%)	32	57
8	V	182/190 (96%)	175 (96%)	7 (4%)	33	58
9	I	172/173 (99%)	167 (97%)	5 (3%)	42	69
9	W	172/173 (99%)	166 (96%)	6 (4%)	36	62
10	J	173/175 (99%)	167 (96%)	6 (4%)	36	62
10	X	173/175 (99%)	167 (96%)	6 (4%)	36	62
11	K	169/169 (100%)	164 (97%)	5 (3%)	41	68
11	Y	169/169 (100%)	164 (97%)	5 (3%)	41	68
12	L	185/185 (100%)	178 (96%)	7 (4%)	33	58
12	Z	185/185 (100%)	178 (96%)	7 (4%)	33	58
13	M	192/208 (92%)	185 (96%)	7 (4%)	35	61
13	a	191/208 (92%)	185 (97%)	6 (3%)	40	67
14	N	162/162 (100%)	158 (98%)	4 (2%)	47	73
14	b	162/162 (100%)	159 (98%)	3 (2%)	57	80
15	c	1/1 (100%)	1 (100%)	0	100	100
15	d	1/1 (100%)	1 (100%)	0	100	100
15	e	1/1 (100%)	1 (100%)	0	100	100
15	f	1/1 (100%)	1 (100%)	0	100	100
15	g	1/1 (100%)	1 (100%)	0	100	100
15	h	1/1 (100%)	1 (100%)	0	100	100
All	All	5304/5546 (96%)	5109 (96%)	195 (4%)	34	60

5 of 195 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	Q	169	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
6	T	203	ASN
4	R	40	LEU
5	S	29	LYS
7	U	154	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 106 such sidechains are listed below:

Mol	Chain	Res	Type
2	P	123	GLN
5	S	116	GLN
13	a	48	ASN
3	Q	38	ASN
4	R	15	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
15	PHL	e	4	15,14	11,11,11	1.88	1 (9%)	11,13,13	1.49	2 (18%)
15	PHL	d	4	11,15	11,11,11	1.66	1 (9%)	11,13,13	1.66	3 (27%)
15	PHL	c	4	15,8	11,11,11	1.64	1 (9%)	11,13,13	1.47	3 (27%)
15	PHL	f	4	15,8	11,11,11	1.63	1 (9%)	11,13,13	1.42	2 (18%)
15	PHL	g	4	11,15	11,11,11	1.75	1 (9%)	11,13,13	1.47	3 (27%)
15	PHL	h	4	15,14	11,11,11	1.91	1 (9%)	11,13,13	1.88	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	PHL	e	4	15,14	-	3/6/6/6	0/1/1/1
15	PHL	d	4	11,15	-	2/6/6/6	0/1/1/1
15	PHL	c	4	15,8	-	4/6/6/6	0/1/1/1
15	PHL	f	4	15,8	-	4/6/6/6	0/1/1/1
15	PHL	g	4	11,15	-	2/6/6/6	0/1/1/1
15	PHL	h	4	15,14	-	4/6/6/6	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	h	4	PHL	CB-CG	-5.85	1.37	1.51
15	e	4	PHL	CB-CG	-5.66	1.37	1.51
15	g	4	PHL	CB-CG	-5.35	1.38	1.51
15	c	4	PHL	CB-CG	-5.33	1.38	1.51
15	f	4	PHL	CB-CG	-5.23	1.38	1.51

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	h	4	PHL	CB-CA-C	-3.69	105.55	112.21
15	h	4	PHL	CG-CB-CA	3.64	120.19	113.24
15	e	4	PHL	CB-CA-C	-3.33	106.19	112.21
15	d	4	PHL	CG-CB-CA	3.28	119.50	113.24
15	g	4	PHL	CB-CA-C	-3.06	106.68	112.21

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	d	4	PHL	CA-CB-CG-CD1
15	g	4	PHL	CA-CB-CG-CD1
15	d	4	PHL	CA-CB-CG-CD2
15	g	4	PHL	CA-CB-CG-CD2
15	c	4	PHL	CA-CB-CG-CD2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 9 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
18	MES	K	302	-	12,12,12	2.26	1 (8%)	14,16,16	1.27	2 (14%)
18	MES	Y	301	-	12,12,12	2.25	1 (8%)	14,16,16	1.29	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
18	MES	K	302	-	-	0/6/14/14	0/1/1/1
18	MES	Y	301	-	-	0/6/14/14	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	K	302	MES	C8-S	-7.58	1.66	1.77
18	Y	301	MES	C8-S	-7.56	1.66	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
18	K	302	MES	O3S-S-C8	3.06	110.72	105.77
18	Y	301	MES	O3S-S-C8	2.90	110.46	105.77
18	Y	301	MES	O2S-S-C8	2.74	110.21	106.92
18	K	302	MES	O2S-S-C8	2.38	109.78	106.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	250/250 (100%)	-0.19	12 (4%) 30 32	35, 50, 89, 128	0
1	O	250/250 (100%)	-0.08	11 (4%) 34 37	41, 57, 102, 132	0
2	B	244/258 (94%)	0.00	16 (6%) 18 19	36, 54, 108, 166	0
2	P	244/258 (94%)	0.09	15 (6%) 21 22	40, 57, 110, 166	0
3	C	240/254 (94%)	0.10	21 (8%) 10 10	35, 57, 121, 159	0
3	Q	240/254 (94%)	0.47	31 (12%) 3 3	45, 75, 157, 213	0
4	D	235/260 (90%)	-0.11	5 (2%) 63 66	41, 61, 94, 141	0
4	R	235/260 (90%)	0.01	10 (4%) 35 38	46, 65, 103, 153	0
5	E	231/234 (98%)	-0.00	9 (3%) 39 42	44, 65, 99, 140	0
5	S	231/234 (98%)	0.19	13 (5%) 24 25	46, 74, 118, 155	0
6	F	243/288 (84%)	-0.15	11 (4%) 33 36	40, 60, 110, 142	0
6	T	243/288 (84%)	0.17	14 (5%) 23 24	40, 72, 125, 164	0
7	G	241/252 (95%)	-0.18	10 (4%) 37 40	36, 52, 96, 150	0
7	U	241/252 (95%)	-0.14	7 (2%) 51 55	39, 55, 89, 140	0
8	H	222/232 (95%)	-0.03	4 (1%) 68 71	36, 51, 78, 121	0
8	V	222/232 (95%)	-0.04	6 (2%) 54 58	39, 54, 78, 132	0
9	I	204/205 (99%)	-0.50	3 (1%) 73 75	31, 44, 73, 95	0
9	W	204/205 (99%)	-0.43	3 (1%) 73 75	32, 47, 77, 100	0
10	J	195/198 (98%)	-0.40	3 (1%) 73 75	32, 44, 70, 117	0
10	X	195/198 (98%)	-0.35	3 (1%) 73 75	35, 47, 73, 125	0
11	K	212/212 (100%)	-0.42	1 (0%) 91 91	31, 45, 69, 90	0
11	Y	212/212 (100%)	-0.37	3 (1%) 75 77	34, 47, 74, 99	0
12	L	222/222 (100%)	-0.38	0 100 100	33, 49, 75, 101	0
12	Z	222/222 (100%)	-0.31	3 (1%) 75 77	35, 49, 77, 105	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	M	224/246 (91%)	-0.36	6 (2%) 54 58	30, 49, 74, 94	0
13	a	223/246 (90%)	-0.37	4 (1%) 68 71	32, 49, 73, 93	0
14	N	196/196 (100%)	-0.38	4 (2%) 65 68	32, 45, 73, 100	0
14	b	196/196 (100%)	-0.38	2 (1%) 82 84	33, 46, 75, 96	0
15	c	2/5 (40%)	-0.06	0 100 100	67, 67, 67, 69	0
15	d	2/5 (40%)	-0.48	0 100 100	62, 62, 62, 71	0
15	e	2/5 (40%)	-0.62	0 100 100	53, 53, 53, 64	0
15	f	2/5 (40%)	0.87	0 100 100	70, 70, 70, 76	0
15	g	2/5 (40%)	0.17	0 100 100	62, 62, 62, 73	0
15	h	2/5 (40%)	-0.39	0 100 100	56, 56, 56, 68	0
All	All	6329/6644 (95%)	-0.15	230 (3%) 42 46	30, 54, 101, 213	0

The worst 5 of 230 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	P	219	ALA	12.3
8	V	222	ASP	11.1
8	H	222	ASP	9.6
2	P	222	GLY	9.6
2	B	219	ALA	9.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
15	PHL	c	4	11/11	0.90	0.17	72,77,80,80	0
15	PHL	f	4	11/11	0.90	0.19	73,79,82,82	0
15	PHL	e	4	11/11	0.95	0.16	54,57,60,60	0
15	PHL	g	4	11/11	0.95	0.20	64,68,73,75	0
15	PHL	d	4	11/11	0.96	0.22	60,63,73,73	0
15	PHL	h	4	11/11	0.97	0.17	57,59,64,64	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
16	MG	Z	301	1/1	0.92	0.42	65,65,65,65	0
18	MES	Y	301	12/12	0.92	0.18	50,80,84,86	0
18	MES	K	302	12/12	0.93	0.20	48,80,87,89	0
16	MG	I	301	1/1	0.94	0.30	61,61,61,61	0
16	MG	K	301	1/1	0.95	0.08	42,42,42,42	0
16	MG	N	201	1/1	0.96	0.07	41,41,41,41	0
16	MG	G	301	1/1	0.97	0.04	44,44,44,44	0
17	CL	G	302	1/1	0.98	0.15	30,30,30,30	0
17	CL	U	301	1/1	0.98	0.17	30,30,30,30	0
16	MG	I	302	1/1	0.98	0.05	43,43,43,43	0
16	MG	L	301	1/1	0.98	0.04	49,49,49,49	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.