

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 13, 2023 – 09:53 PM JST

PDB ID	:	5Y9N
Title	:	Crystal structure of Pyrococcus furiosus PbaA (monoclinic form), an archaeal
		homolog of proteasome-assembly chaperone
Authors	:	Yagi-Utsumi, M.; Sikdar, A.; Kozai, T.; Inoue, R.; Sugiyama, M.; Uchihashi,
		T.; Satoh, T.; Kato, K.
Deposited on	:	2017-08-26
Resolution	:	2.55  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$	
$R_{free}$	130704	$1284 \ (2.56-2.52)$	
Clashscore	141614	$1332 \ (2.56-2.52)$	
Ramachandran outliers	138981	1315 (2.56-2.52)	
Sidechain outliers	138945	1315 (2.56-2.52)	
RSRZ outliers	127900	$1272 \ (2.56-2.52)$	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	245	2% <b>8</b> 9%	7% • •
1	В	245	% 90%	5% • •
1	С	245	88%	7% • •
1	D	245	3% 	8% • •
1	Е	245	2% <b>8</b> 6%	9% • •
1	F	245	<mark>6%</mark> 88%	8% • •



Conti	nued from	<i>i</i> previous	page	
Mol	Chain	Length	Quality of chain	
1	G	245	88%	8% ••
1	Н	245	87%	8% •••
1	Ι	245	87%	9% •••
1	J	245	87%	8% • •



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 18821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace	
1	Δ	028	Total	С	Ν	0	S	0	0	0	
1	Л	230	1867	1206	305	346	10	0	0	0	
1	В	238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0	
1	D	200	1867	1206	305	346	10	0	0	0	
1	С	238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0	
1	0	200	1867	1206	305	346	10	0	0	0	
1	Л	228	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0	
1		200	1867	1206	305	346	10		0	0	
1	E	E 238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0	
-			1867	1206	305	346	10				
1	F	F	F 238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0
-	1	200	1867	1206	305	346	10	0	0	0	
1	G	238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0 0			
-	ŭ	200	1867	1206	305	346	10	0	0	0	
1	н	238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0		
1	11	200	1867	1206	305	346	10	0	0	0	
1	т	238	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	0	0	
	1	200	1867	1206	305	346	10	0	U	0	
1	T	238	Total	$\mathbf{C}$	N	0	S	0	0		
	5	230	1867	1206	305	346	10	0	0	0	

• Molecule 1 is a protein called PbaA.

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-2	GLY	-	expression tag	UNP Q8U4Q9
А	-1	SER	-	expression tag	UNP Q8U4Q9
А	0	HIS	-	expression tag	UNP Q8U4Q9
В	-2	GLY	-	expression tag	UNP Q8U4Q9
В	-1	SER	-	expression tag	UNP Q8U4Q9
В	0	HIS	-	expression tag	UNP Q8U4Q9
С	-2	GLY	-	expression tag	UNP Q8U4Q9
С	-1	SER	-	expression tag	UNP Q8U4Q9
С	0	HIS	-	expression tag	UNP Q8U4Q9



Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	GLY	-	expression tag	UNP Q8U4Q9
D	-1	SER	-	expression tag	UNP Q8U4Q9
D	0	HIS	-	expression tag	UNP Q8U4Q9
Е	-2	GLY	-	expression tag	UNP Q8U4Q9
E	-1	SER	-	expression tag	UNP Q8U4Q9
E	0	HIS	-	expression tag	UNP Q8U4Q9
F	-2	GLY	-	expression tag	UNP Q8U4Q9
F	-1	SER	-	expression tag	UNP Q8U4Q9
F	0	HIS	-	expression tag	UNP Q8U4Q9
G	-2	GLY	-	expression tag	UNP Q8U4Q9
G	-1	SER	-	expression tag	UNP Q8U4Q9
G	0	HIS	-	expression tag	UNP Q8U4Q9
Н	-2	GLY	-	expression tag	UNP Q8U4Q9
Н	-1	SER	-	expression tag	UNP Q8U4Q9
Н	0	HIS	-	expression tag	UNP Q8U4Q9
Ι	-2	GLY	-	expression tag	UNP Q8U4Q9
Ι	-1	SER	-	expression tag	UNP Q8U4Q9
Ι	0	HIS	-	expression tag	UNP Q8U4Q9
J	-2	GLY	-	expression tag	UNP Q8U4Q9
J	-1	SER	-	expression tag	UNP Q8U4Q9
J	0	HIS	-	expression tag	UNP Q8U4Q9

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Cl 1 1	0	0
2	В	1	Total Cl 1 1	0	0
2	С	2	Total Cl 2 2	0	0
2	D	2	Total Cl 2 2	0	0
2	Е	2	Total Cl 2 2	0	0
2	G	1	Total Cl 1 1	0	0
2	Н	1	Total Cl 1 1	0	0

• Molecule 3 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	11	Total O 11 11	0	0
3	В	16	Total         O           16         16	0	0
3	С	19	Total         O           19         19	0	0
3	D	12	Total         O           12         12	0	0
3	Ε	10	Total O 10 10	0	0
3	F	10	Total O 10 10	0	0
3	G	20	Total O 20 20	0	0
3	Н	16	Total O 16 16	0	0
3	Ι	15	$\begin{array}{cc} \text{Total} & \text{O} \\ 15 & 15 \end{array}$	0	0
3	J	12	TotalO1212	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: PbaA









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	92.69Å 200.94Å $92.76$ Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $110.96^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{Posolution} \left( \overset{\circ}{\mathbf{A}} \right)$	20.00 - 2.55	Depositor
Resolution (A)	19.99 - 2.55	EDS
% Data completeness	99.0 (20.00-2.55)	Depositor
(in resolution range)	99.2(19.99-2.55)	EDS
R <sub>merge</sub>	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.64 (at 2.56 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
P. P.	0.208 , $0.240$	Depositor
$n, n_{free}$	0.214 , $0.244$	DCC
$R_{free}$ test set	5085 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	76.3	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , $55.2$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	18821	wwPDB-VP
Average B, all atoms $(Å^2)$	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	B	ond angles
MOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.58	0/1902	0.76	0/2567
1	В	0.63	0/1902	0.79	3/2567~(0.1%)
1	С	0.64	0/1902	0.77	0/2567
1	D	0.59	0/1902	0.77	2/2567~(0.1%)
1	Е	0.57	0/1902	0.77	2/2567~(0.1%)
1	F	0.60	0/1902	0.79	2/2567~(0.1%)
1	G	0.60	0/1902	0.78	2/2567~(0.1%)
1	Н	0.61	0/1902	0.80	4/2567~(0.2%)
1	Ι	0.59	0/1902	0.76	1/2567~(0.0%)
1	J	0.58	0/1902	0.77	2/2567~(0.1%)
All	All	0.60	0/19020	0.78	18/25670~(0.1%)

There are no bond length outliers.

All	(18)	bond	angle	outliers	are	listed	below:
-----	------	------	-------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	Н	230	ARG	NE-CZ-NH1	-8.15	116.22	120.30
1	Н	5	LYS	CA-CB-CG	7.72	130.38	113.40
1	Ι	5	LYS	CA-CB-CG	7.55	130.02	113.40
1	Н	230	ARG	NE-CZ-NH2	6.98	123.79	120.30
1	F	138	LYS	CD-CE-NZ	6.91	127.58	111.70
1	J	5	LYS	CD-CE-NZ	6.75	127.23	111.70
1	Е	104	LYS	CA-CB-CG	6.53	127.76	113.40
1	J	222	MET	CG-SD-CE	6.25	110.19	100.20
1	В	214	MET	CG-SD-CE	5.95	109.72	100.20
1	F	5	LYS	CD-CE-NZ	5.71	124.83	111.70
1	G	228	GLU	OE1-CD-OE2	-5.67	116.49	123.30
1	Н	221	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	D	220	ARG	NE-CZ-NH1	5.29	122.94	120.30
1	В	221	ARG	NE-CZ-NH1	5.28	122.94	120.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	Е	221	ARG	NE-CZ-NH2	-5.19	117.71	120.30
1	D	231	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	В	100	ASN	CB-CG-OD1	-5.10	111.40	121.60
1	G	221	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1867	0	1916	13	0
1	В	1867	0	1916	7	0
1	С	1867	0	1916	13	0
1	D	1867	0	1916	15	0
1	Е	1867	0	1916	15	0
1	F	1867	0	1916	13	0
1	G	1867	0	1916	11	0
1	Н	1867	0	1916	13	0
1	Ι	1867	0	1916	13	0
1	J	1867	0	1916	11	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Е	2	0	0	0	0
2	G	1	0	0	0	0
2	Н	1	0	0	0	0
3	А	11	0	0	0	0
3	В	16	0	0	0	0
3	С	19	0	0	1	0
3	D	12	0	0	0	0
3	Е	10	0	0	0	0
3	F	10	0	0	1	0
3	G	20	0	0	1	0
3	Н	16	0	0	3	0



	f = f = f = f = f = f = f = f = f = f =									
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes				
3	Ι	15	0	0	0	0				
3	J	12	0	0	1	0				
All	All	18821	0	19160	103	0				

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:E:5:LYS:HB3	1:E:6:PRO:HD2	1.75	0.67	
1:J:86:LEU:HD22	1:J:90:LEU:HD23	1.80	0.63	
1:I:5:LYS:HD3	1:I:8:ASN:OD1	1.99	0.63	
1:H:86:LEU:HD22	1:H:90:LEU:HD23	1.82	0.62	
1:D:118:ILE:HD12	1:F:231:ARG:HD3	1.82	0.62	
1:J:37:LEU:HD11	1:J:195:LEU:HD23	1.83	0.61	
1:A:86:LEU:HD22	1:A:90:LEU:HD23	1.83	0.60	
1:B:37:LEU:HD11	1:B:195:LEU:HD23	1.84	0.59	
1:F:37:LEU:HD11	1:F:195:LEU:HD23	1.84	0.59	
1:D:86:LEU:HD22	1:D:90:LEU:HD23	1.85	0.59	
1:E:37:LEU:HD11	1:E:195:LEU:HD23	1.84	0.59	
1:D:150:TYR:CD1	1:F:239:THR:HG22	2.38	0.59	
1:H:235:ARG:O	1:H:239:THR:HG23	2.03	0.58	
1:B:235:ARG:O	1:B:239:THR:HG23	2.03	0.58	
1:G:235:ARG:O	1:G:239:THR:HG23	2.03	0.58	
1:F:86:LEU:HD22	1:F:90:LEU:HD23	1.86	0.58	
1:E:235:ARG:O	1:E:239:THR:HG23	2.04	0.58	
1:D:235:ARG:O	1:D:239:THR:HG23	2.03	0.58	
1:I:235:ARG:O	1:I:239:THR:HG23	2.03	0.58	
1:D:37:LEU:HD11	1:D:195:LEU:HD23	1.85	0.58	
1:C:235:ARG:O	1:C:239:THR:HG23	2.03	0.57	
1:F:235:ARG:O	1:F:239:THR:HG23	2.03	0.57	
1:I:37:LEU:HD11	1:I:195:LEU:HD23	1.86	0.57	
1:H:121:PHE:N	3:H:401:HOH:O	2.29	0.57	
1:I:86:LEU:HD22	1:I:90:LEU:HD23	1.85	0.57	
1:A:235:ARG:O	1:A:239:THR:HG23	2.03	0.57	
1:J:235:ARG:O	1:J:239:THR:HG23	2.03	0.57	
1:E:86:LEU:HD22	1:E:90:LEU:HD23	1.85	0.56	
1:H:5:LYS:HD2	1:H:8:ASN:OD1	2.07	0.55	
1:A:231:ARG:HD3	1:I:118:ILE:HD12	1.89	0.54	
1:G:86:LEU:HD22	1:G:90:LEU:HD23	1.89	0.54	



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:G:37:LEU:HD11	1:G:195:LEU:HD23	1.91	0.53	
1:H:122:LYS:HD2	3:H:402:HOH:O	2.07	0.53	
1:D:240:MET:SD	1:J:215:ILE:HD11	2.50	0.52	
1:A:118:ILE:HD12	1:I:231:ARG:HD3	1.90	0.52	
1:E:5:LYS:CB	1:E:6:PRO:HD2	2.40	0.52	
1:A:37:LEU:HD11	1:A:195:LEU:HD23	1.92	0.52	
1:C:215:ILE:HD11	1:F:240:MET:SD	2.50	0.51	
1:H:37:LEU:HD11	1:H:195:LEU:HD23	1.91	0.51	
1:J:166:ARG:NH2	3:J:301:HOH:O	2.42	0.50	
1:E:231:ARG:HD3	1:J:118:ILE:HD12	1.93	0.50	
1:C:67:PRO:HG3	1:D:162:TRP:CE2	2.46	0.50	
1:G:15:GLU:HG3	3:G:416:HOH:O	2.11	0.49	
1:C:109:LYS:HE3	3:C:411:HOH:O	2.12	0.49	
1:C:231:ARG:HD3	1:G:118:ILE:HD12	1.94	0.48	
1:C:37:LEU:HD11	1:C:195:LEU:HD23	1.94	0.48	
1:C:86:LEU:HD22	1:C:90:LEU:HD23	1.94	0.48	
1:F:170:LYS:HA	3:F:303:HOH:O	2.15	0.47	
1:H:120:PHE:N	3:H:401:HOH:O	2.43	0.47	
1:I:60:ILE:HD12	1:J:130:ILE:HG12	1.96	0.47	
1:B:86:LEU:HD22	1:B:90:LEU:HD23	1.97	0.46	
1:B:118:ILE:O	1:B:118:ILE:CG2	2.63	0.46	
1:F:118:ILE:O	1:F:118:ILE:CG2	2.64	0.46	
1:G:118:ILE:CG2	1:G:118:ILE:O	2.64	0.46	
1:J:118:ILE:O	1:J:118:ILE:CG2	2.64	0.46	
1:A:118:ILE:CG2	1:A:118:ILE:O	2.64	0.46	
1:E:118:ILE:O	1:E:118:ILE:CG2	2.64	0.46	
1:E:240:MET:SD	1:I:215:ILE:HD11	2.56	0.46	
1:I:118:ILE:CG2	1:I:118:ILE:O	2.64	0.46	
1:C:118:ILE:CG2	1:C:118:ILE:O	2.64	0.46	
1:D:231:ARG:HD3	1:F:118:ILE:HD12	1.96	0.46	
1:B:118:ILE:O	1:B:118:ILE:HG22	2.16	0.46	
1:D:118:ILE:O	1:D:118:ILE:CG2	2.64	0.45	
1:H:118:ILE:CG2	1:H:118:ILE:O	2.64	0.45	
1:A:162:TRP:CE2	1:E:67:PRO:HG3	2.51	0.45	
1:D:239:THR:HG22	1:F:150:TYR:CD1	2.51	0.45	
1:G:118:ILE:O	1:G:118:ILE:HG22	2.16	0.45	
1:D:118:ILE:HD13	1:D:118:ILE:HA	1.93	0.45	
1:A:118:ILE:O	1:A:118:ILE:HG22	2.17	0.45	
1:C:37:LEU:CD1	1:C:195:LEU:HD23	2.47	0.45	
1:D:118:ILE:O	1:D:118:ILE:HG22	2.17	0.45	
1:H:15:GLU:HG2	1:H:16:ASN:ND2	2.31	0.45	



A 4 1	A + 0	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:F:118:ILE:O	1:F:118:ILE:HG22	2.17	0.44
1:B:118:ILE:HD12	1:H:231:ARG:HD3	2.00	0.44
1:E:118:ILE:O	1:E:118:ILE:HG22	2.17	0.44
1:J:99:VAL:HG21	1:J:163:GLU:HB3	2.00	0.44
1:H:118:ILE:O	1:H:118:ILE:HG22	2.17	0.44
1:A:37:LEU:CD1	1:A:195:LEU:HD23	2.48	0.44
1:I:196:ASN:HD22	1:I:201:LEU:HB2	1.83	0.44
1:G:196:ASN:HD22	1:G:201:LEU:HB2	1.83	0.43
1:C:196:ASN:HD22	1:C:201:LEU:HB2	1.83	0.43
1:B:118:ILE:HD13	1:B:118:ILE:HA	1.92	0.43
1:A:161:LEU:HD21	1:E:65:PRO:HG3	2.01	0.43
1:C:118:ILE:O	1:C:118:ILE:HG22	2.19	0.43
1:J:118:ILE:O	1:J:118:ILE:HG22	2.19	0.43
1:I:118:ILE:O	1:I:118:ILE:HG22	2.18	0.43
1:C:67:PRO:HG3	1:D:162:TRP:CZ2	2.54	0.43
1:A:130:ILE:HD11	1:E:63:GLY:O	2.20	0.42
1:H:37:LEU:CD1	1:H:195:LEU:HD23	2.49	0.42
1:A:196:ASN:HD22	1:A:201:LEU:HB2	1.84	0.42
1:E:118:ILE:HD13	1:E:118:ILE:HA	1.93	0.42
1:D:150:TYR:CG	1:F:239:THR:HG22	2.55	0.42
1:F:118:ILE:HD13	1:F:118:ILE:HA	1.92	0.42
1:I:118:ILE:HD13	1:I:118:ILE:HA	1.93	0.42
1:J:196:ASN:HD22	1:J:201:LEU:HB2	1.85	0.42
1:G:118:ILE:HD13	1:G:118:ILE:HA	1.93	0.41
1:C:118:ILE:HD12	1:G:231:ARG:HD3	2.02	0.41
1:D:196:ASN:HD22	1:D:201:LEU:HB2	1.85	0.41
1:G:60:ILE:HD12	1:H:130:ILE:HG12	2.02	0.41
1:E:5:LYS:CB	1:E:6:PRO:CD	2.99	0.41
1:E:37:LEU:CD1	1:E:195:LEU:HD23	2.49	0.41
1:I:37:LEU:CD1	1:I:195:LEU:HD23	2.51	0.41
1:A:118:ILE:HD13	1:A:118:ILE:HA	1.93	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	В	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	С	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	D	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	Ε	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	F	236/245~(96%)	228~(97%)	6(2%)	2(1%)	19 27
1	G	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	Н	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	Ι	236/245~(96%)	229~(97%)	5(2%)	2(1%)	19 27
1	J	236/245~(96%)	229 (97%)	5 (2%)	2 (1%)	19 27
All	All	2360/2450~(96%)	2289 (97%)	51 (2%)	20 (1%)	19 27

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (20) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	75	ASN
1	В	75	ASN
1	С	75	ASN
1	D	75	ASN
1	Е	75	ASN
1	F	75	ASN
1	G	75	ASN
1	Н	75	ASN
1	Ι	75	ASN
1	J	75	ASN
1	Е	118	ILE
1	А	118	ILE
1	В	118	ILE
1	С	118	ILE
1	D	118	ILE
1	F	118	ILE
1	G	118	ILE
1	Н	118	ILE
1	Ι	118	ILE
1	J	118	ILE



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	201/207~(97%)	192~(96%)	9~(4%)	27	37
1	В	201/207~(97%)	189 (94%)	12 (6%)	19	25
1	С	201/207~(97%)	189 (94%)	12 (6%)	19	25
1	D	201/207~(97%)	190 (94%)	11 (6%)	21	29
1	Ε	201/207~(97%)	186 (92%)	15 (8%)	13	17
1	F	201/207~(97%)	188 (94%)	13 (6%)	17	23
1	G	201/207~(97%)	190 (94%)	11 (6%)	21	29
1	Н	201/207~(97%)	188 (94%)	13 (6%)	17	23
1	Ι	201/207~(97%)	187~(93%)	14 (7%)	15	19
1	J	201/207~(97%)	186 (92%)	15 (8%)	13	17
All	All	2010/2070~(97%)	1885 (94%)	125 (6%)	18	24

All (125) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	35	ASN
1	А	75	ASN
1	А	76	ASN
1	А	118	ILE
1	А	122	LYS
1	А	124	THR
1	А	166	ARG
1	А	195	LEU
1	А	210	LYS
1	В	35	ASN
1	В	64	ARG
1	В	75	ASN
1	В	76	ASN
1	В	104	LYS
1	В	118	ILE
1	В	124	THR



Mol	Chain	Res	Type
1	В	166	ARG
1	В	195	LEU
1	В	210	LYS
1	В	221	ARG
1	В	235	ARG
1	С	5	LYS
1	С	35	ASN
1	С	39	LYS
1	С	75	ASN
1	С	76	ASN
1	С	104	LYS
1	С	118	ILE
1	С	122	LYS
1	С	166	ARG
1	С	195	LEU
1	С	210	LYS
1	С	235	ARG
1	D	35	ASN
1	D	39	LYS
1	D	75	ASN
1	D	76	ASN
1	D	118	ILE
1	D	122	LYS
1	D	124	THR
1	D	166	ARG
1	D	195	LEU
1	D	210	LYS
1	D	213	GLU
1	Е	5	LYS
1	Е	35	ASN
1	Е	39	LYS
1	E	75	ASN
1	E	76	ASN
1	E	96	LYS
1	E	104	LYS
1	E	118	ILE
1	Е	122	LYS
1	E	124	THR
1	Е	166	ARG
1	Е	195	LEU
1	E	210	LYS
1	E	214	MET



Mol	Chain	Res	Type
1	Е	233	MET
1	F	5	LYS
1	F	35	ASN
1	F	39	LYS
1	F	75	ASN
1	F	76	ASN
1	F	104	LYS
1	F	118	ILE
1	F	122	LYS
1	F	124	THR
1	F	166	ARG
1	F	195	LEU
1	F	210	LYS
1	F	232	LYS
1	G	35	ASN
1	G	75	ASN
1	G	76	ASN
1	G	104	LYS
1	G	118	ILE
1	G	122	LYS
1	G	124	THR
1	G	166	ARG
1	G	195	LEU
1	G	210	LYS
1	G	233	MET
1	Н	5	LYS
1	Н	35	ASN
1	Н	39	LYS
1	Н	75	ASN
1	Н	76	ASN
1	Н	106	ASN
1	Н	118	ILE
1	H	122	LYS
1	Н	124	THR
1	Н	166	ARG
1	H	195	LEU
1	H	210	LYS
1	Н	230	ARG
1	Ι	5	LYS
1	Ι	35	ASN
1	Ι	39	LYS
1	Ι	75	ASN



Mol	Chain	Res	Type
1	Ι	76	ASN
1	Ι	104	LYS
1	Ι	118	ILE
1	Ι	122	LYS
1	Ι	124	THR
1	Ι	166	ARG
1	Ι	195	LEU
1	Ι	210	LYS
1	Ι	214	MET
1	Ι	224	GLU
1	J	35	ASN
1	J	39	LYS
1	J	75	ASN
1	J	76	ASN
1	J	104	LYS
1	J	118	ILE
1	J	122	LYS
1	J	124	THR
1	J	163	GLU
1	J	166	ARG
1	J	195	LEU
1	J	210	LYS
1	J	214	MET
1	J	221	ARG
1	J	222	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (50) such sidechains are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	35	ASN
1	А	76	ASN
1	А	196	ASN
1	А	218	GLN
1	А	236	GLN
1	В	35	ASN
1	В	76	ASN
1	В	196	ASN
1	В	218	GLN
1	В	236	GLN
1	С	35	ASN
1	С	76	ASN
1	С	196	ASN



		1	r ng m
Mol	Chain	Res	Type
1	С	218	GLN
1	С	236	GLN
1	D	35	ASN
1	D	76	ASN
1	D	196	ASN
1	D	218	GLN
1	D	236	GLN
1	Е	35	ASN
1	Е	76	ASN
1	Е	196	ASN
1	Е	218	GLN
1	F	35	ASN
1	F	76	ASN
1	F	196	ASN
1	F	218	GLN
1	F	236	GLN
1	G	35	ASN
1	G	76	ASN
1	G	196	ASN
1	G	236	GLN
1	Н	16	ASN
1	Н	35	ASN
1	Н	76	ASN
1	Н	196	ASN
1	Н	218	GLN
1	Н	236	GLN
1	Ι	35	ASN
1	Ι	76	ASN
1	Ι	196	ASN
1	Ι	218	GLN
1	Ι	236	GLN
1	J	35	ASN
1	J	76	ASN
1	J	196	ASN
1	J	218	GLN
1	J	225	GLN
1	J	236	GLN

Continued from previous page...

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RS	SRZ>	>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q < 0.9
1	А	238/245~(97%)	0.08	5(2%)	63	70	58, 82, 114, 154	0
1	В	238/245~(97%)	0.05	3(1%)	77	82	49, 72, 109, 130	0
1	С	238/245~(97%)	0.02	7(2%)	51	59	49, 67, 106, 132	0
1	D	238/245~(97%)	0.22	7 (2%)	51	59	52, 83, 120, 160	0
1	Ε	238/245~(97%)	0.19	6 (2%)	57	63	61, 86, 113, 137	0
1	F	238/245~(97%)	0.35	15~(6%)	20	23	58, 86, 121, 172	0
1	G	238/245~(97%)	0.04	3(1%)	77	82	53, 73, 104, 136	0
1	Η	238/245~(97%)	0.09	9~(3%)	40	47	54, 73, 113, 126	0
1	Ι	238/245~(97%)	0.23	10 (4%)	36	42	60, 83, 117, 146	0
1	J	$23\overline{8}/245~(97\%)$	0.20	13 (5%)	25	30	59, 87, 116, 154	0
All	All	2380/2450~(97%)	0.15	78 (3%)	46	53	49, 80, 115, 172	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	123	ASP	5.5
1	J	123	ASP	4.7
1	Ι	123	ASP	4.1
1	F	120	PHE	3.9
1	G	123	ASP	3.8
1	D	123	ASP	3.8
1	J	121	PHE	3.6
1	Н	121	PHE	3.6
1	А	121	PHE	3.5
1	J	120	PHE	3.4
1	В	121	PHE	3.4
1	С	220	ARG	3.4
1	F	172	TYR	3.2



5Y9	Ν
-----	---

Mol	Chain	Res	Type	RSRZ
1	А	5	LYS	3.1
1	D	120	PHE	3.1
1	Ι	124	THR	3.0
1	J	124	THR	3.0
1	D	121	PHE	2.9
1	F	138	LYS	2.9
1	F	202	ASN	2.8
1	Ι	121	PHE	2.8
1	Е	124	THR	2.8
1	J	122	LYS	2.8
1	Е	188	ALA	2.8
1	Ι	219	LEU	2.8
1	В	120	PHE	2.7
1	Н	20	ILE	2.7
1	С	123	ASP	2.7
1	F	130	ILE	2.7
1	В	123	ASP	2.7
1	Ι	220	ARG	2.6
1	F	174	LEU	2.6
1	F	220	ARG	2.6
1	F	85	PHE	2.6
1	Ι	75	ASN	2.5
1	D	221	ARG	2.5
1	D	124	THR	2.4
1	Ι	122	LYS	2.4
1	F	117	GLY	2.4
1	Е	20	ILE	2.4
1	А	123	ASP	2.4
1	D	129	GLY	2.4
1	А	220	ARG	2.3
1	А	19	PHE	2.3
1	Н	120	PHE	2.3
1	J	220	ARG	2.3
1	Н	124	THR	2.3
1	J	20	ILE	2.3
1	Ι	80	ALA	2.3
1	С	120	PHE	2.2
1	J	173	VAL	2.2
1	Н	144	GLY	2.2
1	J	174	LEU	2.2
1	J	219	LEU	2.2
1	J	232	LYS	2.1



Mol	Chain	Res	Type	RSRZ
1	С	129	GLY	2.1
1	F	122	LYS	2.1
1	J	98	ILE	2.1
1	С	138	LYS	2.1
1	G	130	ILE	2.1
1	Н	220	ARG	2.1
1	Н	202	ASN	2.1
1	J	129	GLY	2.1
1	С	121	PHE	2.1
1	G	120	PHE	2.1
1	D	122	LYS	2.1
1	F	222	MET	2.1
1	Н	123	ASP	2.1
1	Е	168	GLY	2.1
1	Е	123	ASP	2.0
1	Е	173	VAL	2.0
1	Ι	34	ALA	2.0
1	С	174	LEU	2.0
1	F	121	PHE	2.0
1	Ι	120	PHE	2.0
1	Н	135	GLU	2.0
1	F	124	THR	2.0
1	F	112	SER	2.0

Continued from previous page...

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
Mol	Type	Chain	$\mathbf{Res}$	Atoms	RSCC	RSR	$B-factors(Å^2)$	$Q{<}0.9$
2	CL	С	302	1/1	0.91	0.16	89,89,89,89	0
2	CL	Е	302	1/1	0.94	0.16	93,93,93,93	0
2	CL	Н	301	1/1	0.94	0.22	91,91,91,91	0
2	CL	D	302	1/1	0.95	0.15	94,94,94,94	0
2	CL	В	301	1/1	0.95	0.23	83,83,83,83	0
2	CL	А	301	1/1	0.95	0.23	86,86,86,86	0
2	CL	С	301	1/1	0.96	0.28	84,84,84,84	0
2	CL	D	301	1/1	0.97	0.23	97,97,97,97	0
2	CL	G	301	1/1	0.97	0.25	78,78,78,78	0
2	CL	Е	301	1/1	0.97	0.21	87,87,87,87	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

