



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 22, 2023 – 11:43 PM JST

PDB ID : 7YGL
Title : Crystal Structure of the ring nuclease Sso2081 from *Saccharolobus solfataricus* in complex with A4>p cleavage intermediate
Authors : Lin, Z.; Du, L.; Luo, Z.
Deposited on : 2022-07-11
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)
RNA backbone	3102	1008 (2.84-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	 23% 84% 12% ••
1	B	184	 24% 83% 16% •
2	C	4	 25% 75% 50% 25%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CRISPR system ring nuclease SSO2081.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	178	1414	896	240	270	8	0	0	0
1	B	184	1442	914	243	277	8	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	LEU	-	expression tag	UNP Q7LYJ6
A	-4	GLY	-	expression tag	UNP Q7LYJ6
A	-3	SER	-	expression tag	UNP Q7LYJ6
A	-2	GLY	-	expression tag	UNP Q7LYJ6
A	-1	ARG	-	expression tag	UNP Q7LYJ6
A	0	PRO	-	expression tag	UNP Q7LYJ6
A	11	ALA	SER	engineered mutation	UNP Q7LYJ6
B	-5	LEU	-	expression tag	UNP Q7LYJ6
B	-4	GLY	-	expression tag	UNP Q7LYJ6
B	-3	SER	-	expression tag	UNP Q7LYJ6
B	-2	GLY	-	expression tag	UNP Q7LYJ6
B	-1	ARG	-	expression tag	UNP Q7LYJ6
B	0	PRO	-	expression tag	UNP Q7LYJ6
B	11	ALA	SER	engineered mutation	UNP Q7LYJ6

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*AP*AP*(A23))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	4	88	40	20	24	4	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total O 4 4	0	0
3	B	1	Total O 1 1	0	0
3	C	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CRISPR system ring nuclease SSO2081



- Molecule 1: CRISPR system ring nuclease SSO2081



- Molecule 2: RNA (5'-R(*AP*AP*AP*(A23))-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.51Å 39.14Å 74.09Å 90.00° 105.18° 90.00°	Depositor
Resolution (Å)	28.91 – 2.50 28.91 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.8 (28.91-2.50) 98.9 (28.91-2.50)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.258 , 0.279 0.257 , 0.272	Depositor DCC
R_{free} test set	683 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	81.1	Xtrriage
Anisotropy	0.554	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 87.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2950	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A23

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/1431	0.79	0/1921
1	B	0.52	0/1460	0.85	0/1962
2	C	1.08	0/71	1.53	0/109
All	All	0.53	0/2962	0.85	0/3992

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1414	0	1465	13	0
1	B	1442	0	1485	26	0
2	C	88	0	44	4	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
All	All	2950	0	2994	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:VAL:HA	1:B:58:GLN:HB2	1.66	0.77
1:B:50:ARG:HD3	1:B:149:ILE:HG22	1.67	0.76
1:B:16:ILE:HD11	1:B:49:VAL:HG13	1.71	0.71
1:B:51:LEU:HD21	1:B:147:PRO:HG2	1.80	0.64
1:B:16:ILE:HG23	1:B:53:PHE:CE1	2.37	0.59
1:B:63:LYS:HG3	1:B:64:VAL:HG23	1.85	0.57
1:B:2:VAL:HG22	1:B:97:ASP:HA	1.88	0.56
1:B:50:ARG:HG2	1:B:51:LEU:N	2.23	0.54
1:B:51:LEU:HD22	1:B:163:TYR:CE2	2.45	0.52
1:A:105:ARG:NH1	2:C:4:A23:OP1	2.43	0.52
1:A:93:LEU:HD22	1:A:121:VAL:HG21	1.91	0.51
1:B:10:THR:H	2:C:2:A:HO2'	1.60	0.50
1:A:63:LYS:HG3	1:A:64:VAL:HG23	1.93	0.49
1:B:2:VAL:CG1	1:B:29:ILE:HG12	2.42	0.49
1:B:41:GLU:OE2	1:B:144:LYS:NZ	2.46	0.49
1:B:90:GLU:OE2	1:B:118:ASN:ND2	2.36	0.48
1:A:76:ILE:HG22	1:B:174:ILE:HD13	1.95	0.47
1:A:50:ARG:HG2	1:A:149:ILE:HG22	1.97	0.46
1:B:53:PHE:O	1:B:58:GLN:N	2.49	0.46
1:A:168:VAL:HB	1:A:171:ALA:HB2	1.98	0.45
1:B:39:ASN:HB3	1:B:42:VAL:HG22	1.99	0.44
1:B:51:LEU:CD2	1:B:147:PRO:HG2	2.49	0.43
1:B:128:ILE:HG22	2:C:3:A:O2'	2.18	0.43
1:A:14:GLY:HA3	1:A:126:SER:HB2	2.00	0.43
1:A:22:LEU:O	1:A:25:LYS:N	2.52	0.43
1:B:54:VAL:HG22	1:B:58:GLN:NE2	2.34	0.43
1:B:59:GLU:HG3	1:B:152:ARG:HG2	2.00	0.43
1:B:101:ILE:H	1:B:101:ILE:HG13	1.71	0.43
1:B:106:LYS:HG3	2:C:2:A:OP2	2.18	0.43
1:A:168:VAL:HG12	1:A:170:ASP:H	1.83	0.42
1:B:54:VAL:CA	1:B:58:GLN:HB2	2.44	0.42
1:B:53:PHE:O	1:B:57:ILE:N	2.51	0.42
1:A:39:ASN:HB3	1:A:42:VAL:HG22	2.03	0.41
1:A:18:SER:O	1:A:22:LEU:HG	2.21	0.41
1:A:156:ARG:HA	1:A:156:ARG:HD2	1.68	0.41
1:A:141:ARG:HH11	1:A:141:ARG:HD2	1.76	0.40
1:B:117:LYS:HB2	1:B:178:ILE:HD12	2.02	0.40
1:B:53:PHE:HD1	1:B:53:PHE:HA	1.77	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/184 (96%)	168 (96%)	8 (4%)	0	100	100
1	B	182/184 (99%)	176 (97%)	6 (3%)	0	100	100
All	All	358/368 (97%)	344 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/162 (98%)	154 (98%)	4 (2%)	47	73
1	B	160/162 (99%)	156 (98%)	4 (2%)	47	73
All	All	318/324 (98%)	310 (98%)	8 (2%)	47	73

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	TYR
1	A	102	THR
1	A	105	ARG
1	A	169	GLN
1	B	50	ARG
1	B	51	LEU
1	B	53	PHE
1	B	107	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	3/4 (75%)	3 (100%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	A
2	C	3	A
2	C	4	A23

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A23	C	4	2	19,28,29	0.99	1 (5%)	19,43,46	2.14	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A23	C	4	2	-	2/3/35/36	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	A23	C4-N3	2.95	1.39	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	A23	O2'-PC-O1C	-5.36	101.61	115.76
2	C	4	A23	O2C-PC-O1C	4.27	123.69	109.89
2	C	4	A23	O4'-C4'-C3'	2.38	109.97	104.87
2	C	4	A23	C5-C6-N6	2.34	123.91	120.35
2	C	4	A23	N3-C2-N1	-2.23	125.19	128.68
2	C	4	A23	O5'-C5'-C4'	-2.06	101.98	108.99

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	4	A23	C3'-C4'-C5'-O5'
2	C	4	A23	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4	A23	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/184 (96%)	1.20	43 (24%) 0 0	71, 114, 157, 180	0
1	B	184/184 (100%)	1.22	44 (23%) 0 0	62, 116, 164, 197	0
2	C	3/4 (75%)	2.73	3 (100%) 0 0	89, 89, 92, 102	0
All	All	365/372 (98%)	1.22	90 (24%) 0 0	62, 115, 160, 197	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	CYS	8.4
1	B	149	ILE	7.2
1	A	61	PHE	6.2
1	B	153	GLY	6.2
1	B	102	THR	6.1
1	A	153	GLY	5.6
1	A	13	GLY	5.5
1	B	150	LYS	5.3
1	A	103	GLY	5.1
1	A	14	GLY	5.0
1	B	144	LYS	4.9
1	A	102	THR	4.8
1	B	104	GLY	4.8
1	B	13	GLY	4.7
1	B	100	ASP	4.7
1	A	104	GLY	4.6
1	A	156	ARG	4.6
1	A	12	PRO	4.2
1	B	-2	GLY	4.2
1	B	9	GLY	4.2
1	B	103	GLY	4.0
1	A	8	LEU	3.9
1	A	100	ASP	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	106	LYS	3.9
1	B	22	LEU	3.9
1	B	8	LEU	3.8
1	A	59	GLU	3.8
1	A	149	ILE	3.6
1	B	12	PRO	3.6
1	B	7	THR	3.6
1	A	106	LYS	3.6
1	B	170	ASP	3.5
1	A	77	TYR	3.5
1	A	150	LYS	3.4
1	B	15	VAL	3.4
1	B	148	GLU	3.4
1	B	142	GLU	3.3
1	B	143	LEU	3.3
1	A	64	VAL	3.3
1	A	146	ILE	3.2
1	B	101	ILE	3.2
1	B	159	MET	3.2
1	A	63	LYS	3.1
1	A	72	ASP	3.1
1	A	19	PHE	3.0
1	A	107	SER	3.0
1	B	109	SER	2.9
1	B	135	LYS	2.9
1	A	101	ILE	2.9
2	C	2	A	2.9
1	A	24	LYS	2.9
2	C	3	A	2.8
1	A	160	LYS	2.8
1	B	77	TYR	2.8
1	A	7	THR	2.7
1	B	151	ASN	2.7
1	B	138	LYS	2.6
1	A	9	GLY	2.6
1	A	118	ASN	2.6
2	C	1	A	2.5
1	B	28	ASN	2.5
1	A	169	GLN	2.5
1	A	11	ALA	2.5
1	A	38	SER	2.4
1	B	5	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	155	CYS	2.4
1	B	107	SER	2.4
1	B	156	ARG	2.4
1	B	93	LEU	2.4
1	A	15	VAL	2.3
1	A	108	MET	2.3
1	B	128	ILE	2.3
1	B	31	GLU	2.3
1	A	154	GLU	2.3
1	B	-1	ARG	2.3
1	A	21	TYR	2.2
1	B	125	THR	2.1
1	A	60	LYS	2.1
1	A	65	GLU	2.1
1	B	57	ILE	2.1
1	B	63	LYS	2.1
1	B	154	GLU	2.1
1	B	6	ALA	2.1
1	A	151	ASN	2.1
1	A	10	THR	2.0
1	A	6	ALA	2.0
1	A	62	PRO	2.0
1	A	167	ILE	2.0
1	B	146	ILE	2.0
1	B	52	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A23	C	4	25/26	0.93	0.26	53,78,99,120	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.