



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:39 pm BST

PDB ID : 4YIG  
Title : vaccinia virus D4/A20(1-50) in complex with dsDNA containing an abasic site and free uracyl  
Authors : tarbouriech, N.; burmeister, W.P.; iseni, F.  
Deposited on : 2015-03-02  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	
1	E	232	
1	I	232	
2	B	52	
2	F	52	
2	J	52	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	10	 70% 30%
3	G	10	 90% 10%
3	K	10	 80% 20%
4	D	10	 50% 50%
4	H	10	 80% 20%
4	L	10	 100%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uracil-DNA glycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	Total 1788	C 1158	N 295	O 329	S 6	0	0	0
1	E	220	Total 1788	C 1158	N 295	O 329	S 6	0	0	0
1	I	220	Total 1788	C 1158	N 295	O 329	S 6	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP P20536
A	-12	GLY	-	expression tag	UNP P20536
A	-11	SER	-	expression tag	UNP P20536
A	-10	SER	-	expression tag	UNP P20536
A	-9	HIS	-	expression tag	UNP P20536
A	-8	HIS	-	expression tag	UNP P20536
A	-7	HIS	-	expression tag	UNP P20536
A	-6	HIS	-	expression tag	UNP P20536
A	-5	HIS	-	expression tag	UNP P20536
A	-4	HIS	-	expression tag	UNP P20536
A	-3	SER	-	expression tag	UNP P20536
A	-2	GLN	-	expression tag	UNP P20536
A	-1	ASP	-	expression tag	UNP P20536
A	0	PRO	-	expression tag	UNP P20536
E	-13	MET	-	initiating methionine	UNP P20536
E	-12	GLY	-	expression tag	UNP P20536
E	-11	SER	-	expression tag	UNP P20536
E	-10	SER	-	expression tag	UNP P20536
E	-9	HIS	-	expression tag	UNP P20536
E	-8	HIS	-	expression tag	UNP P20536
E	-7	HIS	-	expression tag	UNP P20536
E	-6	HIS	-	expression tag	UNP P20536
E	-5	HIS	-	expression tag	UNP P20536

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-4	HIS	-	expression tag	UNP P20536
E	-3	SER	-	expression tag	UNP P20536
E	-2	GLN	-	expression tag	UNP P20536
E	-1	ASP	-	expression tag	UNP P20536
E	0	PRO	-	expression tag	UNP P20536
I	-13	MET	-	initiating methionine	UNP P20536
I	-12	GLY	-	expression tag	UNP P20536
I	-11	SER	-	expression tag	UNP P20536
I	-10	SER	-	expression tag	UNP P20536
I	-9	HIS	-	expression tag	UNP P20536
I	-8	HIS	-	expression tag	UNP P20536
I	-7	HIS	-	expression tag	UNP P20536
I	-6	HIS	-	expression tag	UNP P20536
I	-5	HIS	-	expression tag	UNP P20536
I	-4	HIS	-	expression tag	UNP P20536
I	-3	SER	-	expression tag	UNP P20536
I	-2	GLN	-	expression tag	UNP P20536
I	-1	ASP	-	expression tag	UNP P20536
I	0	PRO	-	expression tag	UNP P20536

- Molecule 2 is a protein called DNA polymerase processivity factor component A20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	50	Total	C	N	O	S	0	0	0
			398	256	63	78	1			
2	F	50	Total	C	N	O	S	0	0	0
			398	256	63	78	1			
2	J	50	Total	C	N	O	S	0	0	0
			398	256	63	78	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P20995
B	0	ALA	-	expression tag	UNP P20995
B	1	MET	-	expression tag	UNP P20995
B	2	ALA	-	expression tag	UNP P20995
F	-1	GLY	-	expression tag	UNP P20995
F	0	ALA	-	expression tag	UNP P20995
F	1	MET	-	expression tag	UNP P20995
F	2	ALA	-	expression tag	UNP P20995
J	-1	GLY	-	expression tag	UNP P20995

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
J	0	ALA	-	expression tag	UNP P20995
J	1	MET	-	expression tag	UNP P20995
J	2	ALA	-	expression tag	UNP P20995


- Molecule 3 is a DNA chain called DNA (5'-D(\*CP\*TP\*GP\*TP\*(ORP)P\*AP\*TP\*CP\*TP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	10	Total 190	C 93	N 26	O 62	P 9	0	0	0
3	G	10	Total 190	C 93	N 26	O 62	P 9	0	0	0
3	K	10	Total 190	C 93	N 26	O 62	P 9	0	0	0

- Molecule 4 is a DNA chain called DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*AP\*AP\*CP\*AP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	10	Total 206	C 99	N 45	O 53	P 9	0	0	0
4	H	10	Total 206	C 99	N 45	O 53	P 9	0	0	0
4	L	10	Total 206	C 99	N 45	O 53	P 9	0	0	0

- Molecule 5 is URACIL (three-letter code: URA) (formula: C<sub>4</sub>H<sub>4</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	2	2		
5	E	1	Total	C	N	O	0	0
			8	4	2	2		
5	I	1	Total	C	N	O	0	0
			8	4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	19	Total	O	0	0
			19	19		
6	B	2	Total	O	0	0
			2	2		
6	C	3	Total	O	0	0
			3	3		
6	E	10	Total	O	0	0
			10	10		
6	F	1	Total	O	0	0
			1	1		
6	G	1	Total	O	0	0
			1	1		
6	H	1	Total	O	0	0
			1	1		
6	I	13	Total	O	0	0
			13	13		
6	J	3	Total	O	0	0
			3	3		

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	K	1	Total	O	0	0
			1	1		



### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uracil-DNA glycosylase

Chain A: 




- Molecule 1: Uracil-DNA glycosylase

Chain E: 




- Molecule 1: Uracil-DNA glycosylase

Chain I: 





- Molecule 2: DNA polymerase processivity factor component A20

Chain B: 




- Molecule 2: DNA polymerase processivity factor component A20

Chain F:  88% 8%



- Molecule 2: DNA polymerase processivity factor component A20

Chain J:  81% 15%



- Molecule 3: DNA (5'-D(\*CP\*TP\*GP\*TP\*(ORP)P\*AP\*TP\*CP\*TP\*T)-3')

Chain C:  70% 30%



- Molecule 3: DNA (5'-D(\*CP\*TP\*GP\*TP\*(ORP)P\*AP\*TP\*CP\*TP\*T)-3')

Chain G:  90% 10%



- Molecule 3: DNA (5'-D(\*CP\*TP\*GP\*TP\*(ORP)P\*AP\*TP\*CP\*TP\*T)-3')

Chain K:  80% 20%




- Molecule 4: DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*AP\*AP\*CP\*AP\*G)-3')

Chain D:  50% 50%



- Molecule 4: DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*AP\*AP\*CP\*AP\*G)-3')

Chain H:  80% 20%



- Molecule 4: DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*AP\*AP\*CP\*AP\*G)-3')

Chain L:  100%

There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.09Å 136.09Å 161.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.86 – 2.70 48.86 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.86-2.70) 99.9 (48.86-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.55 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.180 , 0.228 0.187 , 0.231	Depositor DCC
$R_{free}$ test set	2227 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	71.0	Xtrriage
Anisotropy	0.002	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 38.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.044 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: URA, ORP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/1838	0.94	4/2500 (0.2%)
1	E	0.75	1/1838 (0.1%)	0.92	2/2500 (0.1%)
1	I	0.72	0/1838	0.88	3/2500 (0.1%)
2	B	0.87	0/405	0.88	0/545
2	F	0.64	0/405	0.79	0/545
2	J	0.87	0/405	0.84	0/545
3	C	0.72	0/196	1.04	0/298
3	G	0.72	0/196	1.04	0/298
3	K	0.47	0/196	0.99	1/298 (0.3%)
4	D	0.53	0/233	1.09	2/358 (0.6%)
4	H	0.50	0/233	0.88	0/358
4	L	0.50	0/233	0.92	0/358
All	All	0.73	1/8016 (0.0%)	0.92	12/11103 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	119	TRP	CB-CG	-5.04	1.41	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	ASN	CB-CA-C	-8.72	92.96	110.40
4	D	22	DA	O4'-C4'-C3'	-8.69	100.79	106.00
3	K	1	DC	O4'-C4'-C3'	-8.26	101.05	106.00
1	A	83	ASN	N-CA-C	6.17	127.65	111.00
4	D	25	DA	O5'-P-OP1	-5.94	100.35	105.70
1	I	-1	ASP	CB-CG-OD1	5.70	123.43	118.30
1	E	83	ASN	CB-CA-C	-5.68	99.04	110.40
1	E	39	ARG	CB-CA-C	-5.54	99.33	110.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	192	ASP	CB-CG-OD2	-5.23	113.59	118.30
1	I	205	ASP	CB-CG-OD2	-5.15	113.67	118.30
1	A	37	LEU	CA-CB-CG	5.03	126.86	115.30
1	A	18	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1788	0	1781	10	0
1	E	1788	0	1781	10	0
1	I	1788	0	1781	13	0
2	B	398	0	404	1	0
2	F	398	0	404	3	0
2	J	398	0	404	1	0
3	C	190	0	114	2	0
3	G	190	0	114	1	0
3	K	190	0	114	1	0
4	D	206	0	113	2	0
4	H	206	0	113	1	0
4	L	206	0	113	0	0
5	A	8	0	3	0	0
5	E	8	0	3	0	0
5	I	8	0	3	0	0
6	A	19	0	0	2	0
6	B	2	0	0	0	0
6	C	3	0	0	0	0
6	E	10	0	0	0	0
6	F	1	0	0	0	0
6	G	1	0	0	0	0
6	H	1	0	0	0	0
6	I	13	0	0	0	0
6	J	3	0	0	0	0
6	K	1	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7824	0	7245	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:114:ASP:OD1	1:E:214:GLN:NE2	2.30	0.63
1:A:192:ASP:HB3	6:A:408:HOH:O	1.99	0.62
1:E:173:PRO:HB3	2:F:43:TRP:CE2	2.35	0.61
1:E:83:ASN:HB3	1:E:85:THR:HG23	1.86	0.57
1:A:109:ASN:ND2	6:A:401:HOH:O	2.38	0.55
1:I:36:TRP:O	1:I:39:ARG:HD3	2.06	0.54
1:A:7:SER:HB2	1:A:8:HIS:CD2	2.42	0.54
1:I:83:ASN:O	1:I:85:THR:HG23	2.09	0.52
2:B:1:MET:HG3	2:B:2:ALA:N	2.25	0.51
1:A:187:ARG:NH1	4:D:29:DC:OP1	2.43	0.51
3:C:3:DG:H1'	3:C:4:DT:H5''	1.94	0.50
1:A:184:ALA:HB1	1:A:188:GLN:NE2	2.27	0.49
4:H:27:DA:H2'	4:H:28:DA:C8	2.46	0.49
1:A:83:ASN:HB3	1:A:85:THR:HG23	1.93	0.49
1:I:48:LYS:O	1:I:49:PHE:C	2.52	0.47
1:E:173:PRO:HA	2:F:43:TRP:CZ3	2.49	0.47
1:I:6:VAL:HG12	1:I:27:VAL:HG13	1.96	0.47
4:D:27:DA:H2'	4:D:28:DA:C8	2.50	0.47
1:E:2:ASN:HB2	1:E:15:TYR:CZ	2.50	0.47
1:A:23:MET:O	1:A:27:VAL:HG23	2.15	0.46
1:I:153:SER:OG	1:I:205:ASP:OD2	2.28	0.46
1:I:158:LEU:HD23	1:I:182:PRO:HD3	1.98	0.45
1:I:103:ILE:HD13	1:I:103:ILE:H	1.82	0.44
1:I:44:PRO:HD3	1:I:69:PRO:HG2	1.98	0.44
1:A:34:ALA:O	1:A:35:SER:C	2.56	0.44
1:E:22:VAL:HG12	1:E:143:LEU:HD11	2.00	0.44
1:E:173:PRO:HA	2:F:43:TRP:CH2	2.53	0.44
1:A:20:GLU:N	1:A:21:PRO:CD	2.81	0.43
3:K:7:DT:OP2	6:K:101:HOH:O	2.21	0.43
1:I:200:VAL:HG11	2:J:10:LEU:HB3	2.01	0.43
1:A:60:LYS:HD2	1:A:117:ILE:HB	2.01	0.42
1:I:19:TRP:O	1:I:20:GLU:C	2.57	0.42
1:I:84:PHE:CD1	1:I:107:GLY:HA2	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:22:VAL:HG12	1:I:143:LEU:HD11	2.00	0.41
1:E:181:HIS:CG	1:E:182:PRO:HD2	2.55	0.41
3:C:1:DC:O4'	3:G:10:DT:H2''	2.21	0.41
1:E:44:PRO:HB2	1:E:48:LYS:HB2	2.02	0.41
1:I:51:ILE:HD12	1:I:77:VAL:HG23	2.01	0.41
1:E:70:TYR:CE1	1:E:86:LYS:HB3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/232 (94%)	205 (94%)	11 (5%)	2 (1%)	17	40
1	E	218/232 (94%)	203 (93%)	15 (7%)	0	100	100
1	I	218/232 (94%)	199 (91%)	17 (8%)	2 (1%)	17	40
2	B	48/52 (92%)	47 (98%)	1 (2%)	0	100	100
2	F	48/52 (92%)	46 (96%)	2 (4%)	0	100	100
2	J	48/52 (92%)	47 (98%)	1 (2%)	0	100	100
All	All	798/852 (94%)	747 (94%)	47 (6%)	4 (0%)	29	54

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	171	GLU
1	A	138	ASP
1	A	172	SER
1	I	0	PRO



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/213 (95%)	187 (93%)	15 (7%)	13	32
1	E	202/213 (95%)	193 (96%)	9 (4%)	27	55
1	I	202/213 (95%)	189 (94%)	13 (6%)	17	39
2	B	44/44 (100%)	38 (86%)	6 (14%)	3	8
2	F	44/44 (100%)	41 (93%)	3 (7%)	16	36
2	J	44/44 (100%)	37 (84%)	7 (16%)	2	6
All	All	738/771 (96%)	685 (93%)	53 (7%)	14	34

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	VAL
1	A	5	THR
1	A	12	THR
1	A	37	LEU
1	A	38	LEU
1	A	131	LYS
1	A	135	ILE
1	A	150	LYS
1	A	169	LYS
1	A	172	SER
1	A	177	ILE
1	A	186	ASP
1	A	188	GLN
1	A	192	ASP
1	A	196	GLU
2	B	20	LEU
2	B	25	SER
2	B	33	SER
2	B	40	SER
2	B	48	GLN
2	B	50	VAL
1	E	6	VAL

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	E	37	LEU
1	E	38	LEU
1	E	104	ASP
1	E	106	LYS
1	E	114	ASP
1	E	131	LYS
1	E	158	LEU
1	E	188	GLN
2	F	1	MET
2	F	21	ARG
2	F	40	SER
1	I	-1	ASP
1	I	3	SER
1	I	5	THR
1	I	12	THR
1	I	38	LEU
1	I	51	ILE
1	I	72	LYS
1	I	95	SER
1	I	97	SER
1	I	103	ILE
1	I	104	ASP
1	I	127	LEU
1	I	142	LYS
2	J	1	MET
2	J	3	SER
2	J	4	SER
2	J	19	SER
2	J	30	LYS
2	J	40	SER
2	J	48	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	8	HIS
1	A	16	HIS
1	A	109	ASN
1	A	165	ASN
1	A	188	GLN
1	A	206	ASN
2	B	48	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	109	ASN
1	E	151	HIS
1	E	188	GLN
1	I	8	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ORP	K	5	3	9,12,13	1.11	1 (11%)	8,16,19	0.66	0
3	ORP	C	5	3	9,12,13	1.38	1 (11%)	8,16,19	1.11	1 (12%)
3	ORP	G	5	3	9,12,13	1.29	2 (22%)	8,16,19	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ORP	K	5	3	-	0/3/17/18	0/1/1/1
3	ORP	C	5	3	-	1/3/17/18	0/1/1/1
3	ORP	G	5	3	-	2/3/17/18	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	5	ORP	O1-C1	3.11	1.47	1.39

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	5	ORP	O1-C1	2.72	1.46	1.39
3	G	5	ORP	O1-C1	2.26	1.45	1.39
3	G	5	ORP	O4-C4	-2.26	1.40	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	ORP	C1-C2-C3	2.32	107.09	103.93

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	5	ORP	C3-C4-C5-O5
3	G	5	ORP	O4-C4-C5-O5
3	C	5	ORP	C3-C4-C5-O5

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	URA	A	301	-	6,8,8	1.37	1 (16%)	4,10,10	7.58	4 (100%)
5	URA	E	301	-	6,8,8	1.21	1 (16%)	4,10,10	8.85	3 (75%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	URA	I	301	-	6,8,8	1.05	0	4,10,10	8.06	4 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	URA	A	301	-	-	-	0/1/1/1
5	URA	E	301	-	-	-	0/1/1/1
5	URA	I	301	-	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	301	URA	C4-N3	2.20	1.36	1.33
5	A	301	URA	C4-N3	2.03	1.36	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	301	URA	C6-N1-C2	13.97	121.31	114.42
5	I	301	URA	C6-N1-C2	11.91	120.29	114.42
5	A	301	URA	N1-C2-N3	-10.51	120.08	128.43
5	A	301	URA	C6-N1-C2	10.39	119.55	114.42
5	E	301	URA	N1-C2-N3	-10.17	120.34	128.43
5	I	301	URA	N1-C2-N3	-10.15	120.36	128.43
5	E	301	URA	C5-C6-N1	-3.54	119.56	123.96
5	I	301	URA	C5-C6-N1	-3.27	119.89	123.96
5	A	301	URA	C5-C4-N3	-2.61	117.57	123.31
5	A	301	URA	C5-C6-N1	-2.12	121.32	123.96
5	I	301	URA	C5-C4-N3	-2.09	118.71	123.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	220/232 (94%)	-0.34	0 100 100	47, 62, 90, 130	0
1	E	220/232 (94%)	-0.23	0 100 100	50, 74, 100, 148	0
1	I	220/232 (94%)	-0.23	0 100 100	49, 80, 111, 139	0
2	B	50/52 (96%)	-0.47	0 100 100	45, 60, 79, 97	0
2	F	50/52 (96%)	-0.41	0 100 100	63, 86, 107, 108	0
2	J	50/52 (96%)	-0.53	0 100 100	52, 61, 81, 99	0
3	C	9/10 (90%)	-0.49	0 100 100	54, 57, 75, 87	0
3	G	9/10 (90%)	-0.49	0 100 100	58, 63, 82, 89	0
3	K	9/10 (90%)	-0.41	0 100 100	67, 77, 102, 115	0
4	D	10/10 (100%)	-0.72	0 100 100	65, 72, 82, 86	0
4	H	10/10 (100%)	-0.71	0 100 100	75, 81, 96, 107	0
4	L	10/10 (100%)	-0.34	0 100 100	88, 98, 105, 119	0
All	All	867/912 (95%)	-0.32	0 100 100	45, 72, 104, 148	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ORP	K	5	12/13	0.98	0.15	56,64,78,79	0
3	ORP	G	5	12/13	0.99	0.16	50,56,65,74	0
3	ORP	C	5	12/13	0.99	0.16	45,53,60,71	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	URA	I	301	8/8	0.97	0.16	76,80,82,86	0
5	URA	E	301	8/8	0.98	0.15	61,71,82,83	0
5	URA	A	301	8/8	0.98	0.15	62,69,72,77	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.