



wwPDB EM Validation Summary Report ⓘ

Nov 8, 2022 – 06:36 AM JST

PDB ID : 5YKE
EMDB ID : EMD-6831
Title : Structure of pancreatic ATP-sensitive potassium channel bound with glibenclamide and ATPgammaS (focused refinement on TM at 4.11Å)
Authors : Chen, L.; Wu, J.X.
Deposited on : 2017-10-14
Resolution : 4.11 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	
1	C	390	
1	E	390	
1	G	390	
2	B	1582	
2	D	1582	
2	F	1582	
2	H	1582	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 30636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ATP-sensitive inward rectifier potassium channel 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	133	1028	681	174	163	10	0	0
1	C	133	1028	681	174	163	10	0	0
1	E	133	1028	681	174	163	10	0	0
1	G	133	1028	681	174	163	10	0	0

- Molecule 2 is a protein called ATP-binding cassette sub-family C member 8 isoform X2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	845	6598	4345	1102	1115	36	0	0
2	D	845	6598	4345	1102	1115	36	0	0
2	F	845	6598	4345	1102	1115	36	0	0
2	H	845	6598	4345	1102	1115	36	0	0

- Molecule 3 is 5-chloro-N-(2-{4-[(cyclohexylcarbamoyl)sulfamoyl]phenyl}ethyl)-2-methoxybenzamide (three-letter code: GBM) (formula: C₂₃H₂₈ClN₃O₅S).




Mol	Chain	Residues	Atoms					AltConf	
			Total	C	Cl	N	O		S
3	B	1	Total 33	C 23	Cl 1	N 3	O 5	S 1	0
3	D	1	Total 33	C 23	Cl 1	N 3	O 5	S 1	0
3	F	1	Total 33	C 23	Cl 1	N 3	O 5	S 1	0
3	H	1	Total 33	C 23	Cl 1	N 3	O 5	S 1	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-sensitive inward rectifier potassium channel 11






- Molecule 1: ATP-sensitive inward rectifier potassium channel 11







● Molecule 2: ATP-binding cassette sub-family C member 8 isoform X2





● Molecule 2: ATP-binding cassette sub-family C member 8 isoform X2



Q211	D212	L213	G214	V215	R216	F217	L218	Q219	V222	N223	L224	L225	M233	A240	I245	R248	A249	L253	P254	I255	R258	N262	D271	A272	Q273	A274	R275	K276	D277	THR	GLN	SER	PRO	GLN	G283	R289	R297	I300	T304	F305	R306	I307	I320	H326										
L327	G328	K329	GLU	ASN	HIS	VAL	PHE	GLN	PRO	LYS	THR	PHE	LEU	VAL	TYR	PHE	SER	SER	GLU	LEU	N354	V357	L362	R370	Q374	Y377	Y378	Y379	A380	I381	E382	T383	L387	A390	Y396	N397	K398	I399	L402	M406	LEU	SER	MET											
E411	M412	I417	C418	M419	L420	I423	D424	L428	M429	M430	F431	F432	M439	A440	M441	L452	Y453	Y454	I455	L456	G457	V458	S459	A460	L461	A464	I467	L470	A471	P472	A478	L481	H491	E494	Q498	T499	N500	E501	M502	L503	R504	G505	M506	L509										
K510	L511	A512	W514	E515	S516	I517	F518	R521	V522	E523	V524	T525	R526	R527	K528	E529	M530	R534	F545	M546	N547	F559	V563	F566	K567	E568	S569	D570	S574	V575	S579	L580	S581	L586	P589	L590	F591	L592	L593	R598	S616	ALA	GLU	ILE	ARG									
GLU	GLU	GLN	CYS	ALA	PRO	ARG	GLU	PRO	PRO	ALA	ALA	GLY	TYR	GLN	ALA	VAL	VAL	ASN	ARG	LYS	ARG	PRO	ALA	GLU	VAL	GLY	ASP	LEU	LEU	PRO	GLY	ARG	LEU	PRO	ASP	GLY	VAL	ASP	ALA	ALA	ASN	PHE	CYS	VAL	GLN									
ILE	ILE	GLY	PHE	PHE	THR	THR	THR	ASP	GLY	GLY	THR	THR	SER	ASN	ILE	THR	ILE	ILE	ILE	THR	ILE	VAL	VAL	GLN	GLY	VAL	CYS	GLY	SER	ALA	ALA	THR	GLY	THR	THR	GLY	THR	THR	GLY	GLY	THR	THR	THR	ASN										
SER	ASN	LEU	PRO	ASP	SER	GLU	GLY	GLU	ASP	SER	PRO	PRO	GLU	GLU	ALA	GLY	GLY	ASP	ARG	VAL	ALA	TYR	ALA	GLY	GLN	LYS	TRP	LEU	LEU	VAL	GLU	GLU	GLY	THR	ILE	THR	PHE	GLU	SER	GLN	TYR	LYS												
MET	VAL	ILE	GLU	ALA	CYS	SER	LEU	MET	GLN	ASP	ILE	ILE	PRO	LEU	ASP	GLN	THR	GLN	GLY	ARG	GLY	SER	GLY	TYR	GLN	ILE	TRP	ASP	ILE	VAL	TRP	GLN	GLY	VAL	PHE	VAL	VAL	GLY	THR	THR	THR	LEU												
ASP	VAL	HIS	LEU	SER	ASP	HIS	MET	TRP	VAL	GLN	GLY	LEU	GLU	GLU	ASP	VAL	VAL	VAL	VAL	VAL	VAL	GLN	TYR	LEU	PRO	ASP	TRP	ASP	TRP	ASP	ASP	GLY	GLY	GLY	GLY	THR	THR	THR	THR	THR	THR	THR	SER											
GLU	CYS	GLN	LEU	PHE	GLU	HIS	TRP	LYS	VAL	GLN	ASP	GLU	GLU	GLU	GLU	VAL	THR	THR	THR	THR	THR	GLY	SER	PRO	GLN	GLY	ALA	ALA	MET	GLY	LEU	LEU	LEU	ASP	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	SER										
GLU	GLU	ASP	ASN	ASN	LEU	SER	SER	VAL	1996	P997	W998	R999	C1001	T1002	K1003	Y1004	L1005	A1008	S1014	F1018	S1019	Q1020	L1021	L1022	K1023	H1024	V1028	A1029	I1030	D1031	L1034	M1037	T1038	D1039	S1040	ALA	LEU	VAL	LEU	SER	PRO	ALA	ALA	ARG	ASN	CYS								
SER	LEU	SER	GLN	GLU	CYS	ASP	LEU	D1060	Q1061	S1062	V1063	Y1064	A1065	M1066	V1067	F1068	T1069	L1070	L1071	C1072	S1073	L1074	G1075	I1076	V1077	L1078	V1081	V1084	T1085	K1092	V1093	A1094	K1095	R1096	L1097	H1098	R1099	S1100	L1101	R1104	I1105	I1106	L1107	A1108	P1109	M1110	R1111	F1112	F1113	E1114	T1115	T1116	G1119	S1120
L1121	L1122	M1123	R1124	F1125	S1126	S1127	D1128	C1129	M1130	T1131	I1132	I1136	L1140	R1145	S1152	A1153	L1154	T1155	V1156	L1157	S1158	Y1159	V1160	L1165	V1166	A1167	L1168	L1169	P1170	L1171	V1174	R1187	Q1190	T1195	L1198	P1199	L1200	L1201	F1204	A1205	E1206	R1215	A1216	F1217	R1218	Y1219								
Q1224	L1227	Y1230	S1238	L1239	F1240	L1241	T1242	M1245	R1246	W1247	R1251	M1252	E1253	Y1254	N1270	S1271	E1275	L1276	L1280	Y1287	A1288	L1289	S1292	L1295	M1296	R1300	M1305	F1306	I1307	Q1308	L1309	V1312	H1316	A1317	L1318	L1319	K1320	L1321	GLU	ALA	GLU	SER	TYR	GLU										
GLY	LEU	ALA	PRO	SER	LEU	ILE	PRO	VAL	ASP	LYS	ASP	GLN	ILE	GLY	ASN	SER	ALA	VAL	TYR	SER	SER	LEU	VAL	LEU	ASN	ALA	LEU	PRO	VAL	PRO	PHE	GLY	GLN	ILE	ILE	ILE	ARG	CYS	GLY	THR	GLY	GLY	LYS	LYS	SER	SER								
PHE	SER	LEU	PHE	PHE	ARG	MET	VAL	ASP	ASP	ASP	PHE	GLY	ARG	ILE	ILE	ASP	GLY	ILE	PRO	THR	THR	ARG	ARG	ILE	GLN	ASP	PRO	VAL	VAL	PHE	SER	GLY	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR						

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	74633	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.130	Depositor
Minimum map value	-0.061	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.028	Depositor
Map size (\AA)	329.15997, 329.15997, 329.15997	wwPDB
Map dimensions	312, 312, 312	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.055, 1.055, 1.055	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GBM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1055	0.58	0/1434
1	C	0.42	0/1055	0.58	0/1434
1	E	0.42	0/1055	0.58	0/1434
1	G	0.42	0/1055	0.58	0/1434
2	B	0.29	0/6738	0.45	0/9165
2	D	0.29	0/6738	0.45	0/9165
2	F	0.29	0/6738	0.45	0/9165
2	H	0.29	0/6738	0.45	0/9165
All	All	0.31	0/31172	0.47	0/42396

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1028	0	1046	42	0
1	C	1028	0	1046	43	0
1	E	1028	0	1046	42	0
1	G	1028	0	1046	43	0
2	B	6598	0	6827	142	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	6598	0	6827	145	0
2	F	6598	0	6827	145	0
2	H	6598	0	6827	142	0
3	B	33	0	28	4	0
3	D	33	0	28	3	0
3	F	33	0	28	3	0
3	H	33	0	28	3	0
All	All	30636	0	31604	698	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 698 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1125:PHE:HA	2:D:1129:CYS:SG	1.95	1.07
2:F:1125:PHE:HA	2:F:1129:CYS:SG	1.95	1.06
2:H:1125:PHE:HA	2:H:1129:CYS:SG	1.95	1.06
2:B:1125:PHE:HA	2:B:1129:CYS:SG	1.95	1.06
2:H:1124:ARG:HD2	2:H:1318:LEU:HD12	1.41	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/390 (34%)	121 (92%)	10 (8%)	0	100	100
1	C	131/390 (34%)	121 (92%)	10 (8%)	0	100	100
1	E	131/390 (34%)	121 (92%)	10 (8%)	0	100	100
1	G	131/390 (34%)	121 (92%)	10 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	827/1582 (52%)	798 (96%)	25 (3%)	4 (0%)	29	67
2	D	827/1582 (52%)	798 (96%)	25 (3%)	4 (0%)	29	67
2	F	827/1582 (52%)	798 (96%)	25 (3%)	4 (0%)	29	67
2	H	827/1582 (52%)	798 (96%)	25 (3%)	4 (0%)	29	67
All	All	3832/7888 (49%)	3676 (96%)	140 (4%)	16 (0%)	38	71

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	199	LYS
2	D	199	LYS
2	F	199	LYS
2	H	199	LYS
2	B	212	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/339 (32%)	95 (88%)	13 (12%)	5	23
1	C	108/339 (32%)	95 (88%)	13 (12%)	5	23
1	E	108/339 (32%)	95 (88%)	13 (12%)	5	23
1	G	108/339 (32%)	95 (88%)	13 (12%)	5	23
2	B	708/1371 (52%)	671 (95%)	37 (5%)	23	50
2	D	708/1371 (52%)	671 (95%)	37 (5%)	23	50
2	F	708/1371 (52%)	671 (95%)	37 (5%)	23	50
2	H	708/1371 (52%)	671 (95%)	37 (5%)	23	50
All	All	3264/6840 (48%)	3064 (94%)	200 (6%)	22	46

5 of 200 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	119	THR
2	F	1238	SER
2	H	1318	LEU
2	F	193	ARG
2	F	430	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	1135	HIS
1	E	128	GLN
2	H	485	GLN
1	G	128	GLN
1	G	152	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GBM	B	2001	-	35,35,35	2.03	4 (11%)	48,48,48	1.87	7 (14%)
3	GBM	F	2001	-	35,35,35	2.03	4 (11%)	48,48,48	1.87	7 (14%)
3	GBM	H	2001	-	35,35,35	2.03	4 (11%)	48,48,48	1.87	7 (14%)
3	GBM	D	2001	-	35,35,35	2.03	4 (11%)	48,48,48	1.87	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GBM	B	2001	-	-	8/27/35/35	0/3/3/3
3	GBM	F	2001	-	-	8/27/35/35	0/3/3/3
3	GBM	H	2001	-	-	8/27/35/35	0/3/3/3
3	GBM	D	2001	-	-	8/27/35/35	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2001	GBM	C18-S2	-10.16	1.60	1.76
3	D	2001	GBM	C18-S2	-10.16	1.60	1.76
3	F	2001	GBM	C18-S2	-10.16	1.60	1.76
3	H	2001	GBM	C18-S2	-10.16	1.60	1.76
3	B	2001	GBM	C27-C28	4.65	1.49	1.40

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2001	GBM	O5-S2-O4	-9.72	107.60	119.55
3	D	2001	GBM	O5-S2-O4	-9.72	107.60	119.55
3	F	2001	GBM	O5-S2-O4	-9.72	107.60	119.55
3	H	2001	GBM	O5-S2-O4	-9.72	107.60	119.55
3	B	2001	GBM	O7-C28-C27	3.76	122.02	116.55

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	2001	GBM	C27-C26-N10-C25
3	D	2001	GBM	C27-C26-N10-C25

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	F	2001	GBM	C27-C26-N10-C25
3	H	2001	GBM	C27-C26-N10-C25
3	B	2001	GBM	O6-C26-N10-C25


There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2001	GBM	4	0
3	F	2001	GBM	3	0
3	H	2001	GBM	3	0
3	D	2001	GBM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-6831. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 156




Y Index: 156




Z Index: 156


6.2.2 Raw map



X Index: 156



Y Index: 156




Z Index: 156


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 117




Y Index: 195




Z Index: 173


6.3.2 Raw map



X Index: 125



Y Index: 187



Z Index: 175

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.028. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 264 nm³; this corresponds to an approximate mass of 238 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.243 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.243\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.11	-	-
Author-provided FSC curve	4.11	4.81	4.19
Unmasked-calculated*	4.50	6.74	4.75

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-6831 and PDB model 5YKE. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.028 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.028).



















9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.028) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6452	 0.3180
A	 0.7229	 0.3950
B	 0.6331	 0.3060
C	 0.7229	 0.3930
D	 0.6331	 0.3070
E	 0.7229	 0.3950
F	 0.6331	 0.3070
G	 0.7229	 0.3950
H	 0.6331	 0.3060

