



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 25, 2024 – 11:48 PM EDT

PDB ID : 6YLA  
Title : Crystal structure of the SARS-CoV-2 receptor binding domain in complex with CR3022 Fab  
Authors : Huo, J.; Zhao, Y.; Ren, J.; Zhou, D.; Ginn, H.M.; Fry, E.E.; Owens, R.; Stuart, D.I.  
Deposited on : 2020-04-06  
Resolution : 2.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 3% 80% 12% 7%
1	E	213	 83% 12% 5%
2	B	229	 10% 87% 7% 5%
2	H	229	 3% 89% 5% 6%
3	C	220	 7% 93% 6%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	L	220	 <p>A horizontal bar chart representing the quality of chain. The bar is 93% green and 6% yellow. A small red square is at the beginning of the bar. The percentage '93%' is written in the center of the green portion, and '6%' is written at the end of the yellow portion.</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MLI	B	302	-	-	-	X
5	MLI	B	304	-	-	-	X
5	MLI	H	301	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 10090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	203	1604	1027	268	301	8	0	0	0
1	A	199	1582	1014	264	296	8	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	327	GLU	-	expression tag	UNP P0DTC2
E	328	THR	-	expression tag	UNP P0DTC2
E	329	GLY	-	expression tag	UNP P0DTC2
E	533	LYS	-	expression tag	UNP P0DTC2
E	534	HIS	-	expression tag	UNP P0DTC2
E	535	HIS	-	expression tag	UNP P0DTC2
E	536	HIS	-	expression tag	UNP P0DTC2
E	537	HIS	-	expression tag	UNP P0DTC2
E	538	HIS	-	expression tag	UNP P0DTC2
E	539	HIS	-	expression tag	UNP P0DTC2
A	327	GLU	-	expression tag	UNP P0DTC2
A	328	THR	-	expression tag	UNP P0DTC2
A	329	GLY	-	expression tag	UNP P0DTC2
A	533	LYS	-	expression tag	UNP P0DTC2
A	534	HIS	-	expression tag	UNP P0DTC2
A	535	HIS	-	expression tag	UNP P0DTC2
A	536	HIS	-	expression tag	UNP P0DTC2
A	537	HIS	-	expression tag	UNP P0DTC2
A	538	HIS	-	expression tag	UNP P0DTC2
A	539	HIS	-	expression tag	UNP P0DTC2


- Molecule 2 is a protein called Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	1	0
			1609	1023	261	317	8			
2	B	217	Total	C	N	O	S	0	1	0
			1615	1026	262	318	9			

- Molecule 3 is a protein called Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	219	Total	C	N	O	S	0	0	0
			1703	1070	282	347	4			
3	C	219	Total	C	N	O	S	0	0	0
			1703	1070	282	347	4			

- Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	O	S	0	0
			4	2	1	1		
4	E	1	Total	C	O	S	0	0
			4	2	1	1		
4	E	1	Total	C	O	S	0	0
			4	2	1	1		
4	E	1	Total	C	O	S	0	0
			4	2	1	1		
4	H	1	Total	C	O	S	0	0
			4	2	1	1		
4	H	1	Total	C	O	S	0	0
			4	2	1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	L	1	Total	C	O	S	0	0
			4	2	1	1		
4	L	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		
4	C	1	Total	C	O	S	0	0
			4	2	1	1		
4	C	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula:  $C_3H_2O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			7	3	4		
5	E	1	Total	C	O	0	0
			7	3	4		
5	H	1	Total	C	O	0	0
			7	3	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	C	O	0	0
			7	3	4		
5	H	1	Total	C	O	0	0
			7	3	4		
5	B	1	Total	C	O	0	0
			7	3	4		
5	B	1	Total	C	O	0	0
			7	3	4		
5	B	1	Total	C	O	0	0
			7	3	4		
5	C	1	Total	C	O	0	0
			7	3	4		
5	C	1	Total	C	O	0	0
			7	3	4		
5	C	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	E	1	Total	C	N	O	0	0
			14	8	1	5		
6	E	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	14	8	1	5	0	0


- Molecule 7 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	L	1	16	10	6	0	0
7	C	1	16	10	6	0	0

- Molecule 8 is 2-(2-METHOXYETHOXY)ETHANOL (three-letter code: PG0) (formula:  $C_5H_{12}O_3$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			8	5	3		


- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	6	Total	O	0	0
			6	6		
9	H	9	Total	O	0	0
			9	9		
9	L	9	Total	O	0	0
			9	9		
9	A	4	Total	O	0	0
			4	4		
9	B	10	Total	O	0	0
			10	10		
9	C	14	Total	O	0	0
			14	14		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike glycoprotein

Chain E: 




- Molecule 1: Spike glycoprotein

Chain A: 





- Molecule 2: Heavy Chain

Chain H: 



- Molecule 2: Heavy Chain

Chain B: 





- Molecule 3: Light chain

Chain L:  %



● Molecule 3: Light chain

Chain C:  %



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	163.12Å 163.12Å 189.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.31 – 2.42 58.79 – 2.42	Depositor EDS
% Data completeness (in resolution range)	96.8 (55.31-2.42) 92.9 (58.79-2.42)	Depositor EDS
$R_{merge}$	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.68 (at 2.42Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.213 , 0.237 0.213 , 0.236	Depositor DCC
$R_{free}$ test set	4674 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.6	Xtrriage
Anisotropy	0.285	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG0, DMS, MLI, NAG, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1625	0.44	0/2210
1	E	0.25	0/1649	0.43	0/2244
2	B	0.25	0/1660	0.47	0/2261
2	H	0.25	0/1654	0.46	0/2253
3	C	0.25	0/1741	0.45	0/2367
3	L	0.24	0/1741	0.45	0/2367
All	All	0.25	0/10070	0.45	0/13702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1582	0	1507	12	0
1	E	1604	0	1524	12	0
2	B	1615	0	1590	11	0
2	H	1609	0	1585	7	0
3	C	1703	0	1649	9	0
3	L	1703	0	1649	11	0
4	A	12	0	18	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	6	0	0
4	C	8	0	12	0	0
4	E	16	0	24	0	0
4	H	8	0	12	0	0
4	L	8	0	12	0	0
5	B	28	0	8	1	0
5	C	21	0	6	0	0
5	E	14	0	4	0	0
5	H	21	0	6	0	0
6	A	14	0	13	0	0
6	E	28	0	26	0	0
7	C	16	0	22	0	0
7	L	16	0	22	4	0
8	A	8	0	12	0	0
9	A	4	0	0	0	0
9	B	10	0	0	0	0
9	C	14	0	0	0	0
9	E	6	0	0	0	0
9	H	9	0	0	0	0
9	L	9	0	0	0	0
All	All	10090	0	9707	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:455:LEU:HD11	1:A:493:GLN:HG3	1.64	0.79
3:L:47:GLY:H	7:L:1603:1PE:H221	1.52	0.74
1:E:378:LYS:NZ	2:H:57:GLU:OE2	2.24	0.70
1:A:378:LYS:NZ	2:B:57:GLU:OE2	2.27	0.67
2:B:123:PRO:HB3	2:B:149:TYR:HB3	1.82	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/213 (92%)	179 (92%)	16 (8%)	0	100	100
1	E	201/213 (94%)	190 (94%)	11 (6%)	0	100	100
2	B	214/229 (93%)	204 (95%)	9 (4%)	1 (0%)	29	40
2	H	213/229 (93%)	208 (98%)	5 (2%)	0	100	100
3	C	217/220 (99%)	207 (95%)	10 (5%)	0	100	100
3	L	217/220 (99%)	208 (96%)	8 (4%)	1 (0%)	29	40
All	All	1257/1324 (95%)	1196 (95%)	59 (5%)	2 (0%)	47	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	130	PRO
3	L	35	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/185 (94%)	163 (94%)	11 (6%)	18	27
1	E	175/185 (95%)	167 (95%)	8 (5%)	27	42
2	B	183/194 (94%)	181 (99%)	2 (1%)	73	86
2	H	182/194 (94%)	180 (99%)	2 (1%)	73	86
3	C	194/195 (100%)	192 (99%)	2 (1%)	76	87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	L	194/195 (100%)	193 (100%)	1 (0%)	88	95
All	All	1102/1148 (96%)	1076 (98%)	26 (2%)	49	67

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	377	PHE
1	A	440	ASN
3	C	1	ASP
1	A	405	ASP
1	A	477	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

32 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DMS	L	1602	-	3,3,3	0.64	0	3,3,3	0.47	0
5	MLI	B	304	-	6,6,6	1.39	0	7,7,7	1.30	0
5	MLI	C	304	-	6,6,6	1.35	0	7,7,7	1.18	0
5	MLI	B	302	-	6,6,6	1.38	0	7,7,7	1.20	0
4	DMS	H	303	-	3,3,3	0.66	0	3,3,3	0.52	0
4	DMS	E	904	-	3,3,3	0.65	0	3,3,3	0.42	0
5	MLI	C	305	-	6,6,6	1.40	0	7,7,7	1.30	0
4	DMS	E	905	-	3,3,3	0.66	0	3,3,3	0.47	0
6	NAG	E	906	1	14,14,15	0.30	0	17,19,21	0.60	0
7	1PE	C	301	-	15,15,15	0.53	0	14,14,14	0.28	0
4	DMS	C	302	-	3,3,3	0.66	0	3,3,3	0.53	0
5	MLI	B	301	-	6,6,6	1.37	0	7,7,7	1.31	0
8	PG0	A	902	-	7,7,7	0.48	0	6,6,6	0.24	0
7	1PE	L	1603	-	15,15,15	0.53	0	14,14,14	0.25	0
4	DMS	E	902	-	3,3,3	0.66	0	3,3,3	0.52	0
4	DMS	A	903	-	3,3,3	0.66	0	3,3,3	0.51	0
5	MLI	E	903	-	6,6,6	1.40	0	7,7,7	1.37	1 (14%)
4	DMS	C	303	-	3,3,3	0.66	0	3,3,3	0.63	0
5	MLI	H	305	-	6,6,6	1.31	0	7,7,7	1.25	0
4	DMS	E	901	-	3,3,3	0.65	0	3,3,3	0.49	0
5	MLI	C	306	-	6,6,6	1.32	0	7,7,7	1.21	0
5	MLI	H	304	-	6,6,6	1.31	0	7,7,7	1.23	0
4	DMS	L	1601	-	3,3,3	0.66	0	3,3,3	0.82	0
4	DMS	B	305	-	3,3,3	0.66	0	3,3,3	0.52	0
5	MLI	H	301	-	6,6,6	1.31	0	7,7,7	1.26	0
6	NAG	A	904	1	14,14,15	0.73	1 (7%)	17,19,21	0.62	0
4	DMS	A	905	-	3,3,3	0.66	0	3,3,3	1.12	0
6	NAG	E	907	1	14,14,15	0.35	0	17,19,21	0.38	0
5	MLI	B	303	-	6,6,6	1.33	0	7,7,7	1.24	0
4	DMS	H	302	-	3,3,3	0.66	0	3,3,3	0.49	0
4	DMS	A	901	-	3,3,3	0.66	0	3,3,3	0.49	0
5	MLI	E	908	-	6,6,6	1.37	0	7,7,7	1.40	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MLI	C	305	-	-	4/4/4/4	-
5	MLI	E	903	-	-	0/4/4/4	-
5	MLI	B	304	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MLI	H	305	-	-	0/4/4/4	-
5	MLI	C	304	-	-	1/4/4/4	-
5	MLI	H	301	-	-	2/4/4/4	-
5	MLI	C	306	-	-	2/4/4/4	-
6	NAG	E	906	1	-	3/6/23/26	0/1/1/1
6	NAG	A	904	1	-	2/6/23/26	0/1/1/1
7	1PE	C	301	-	-	5/13/13/13	-
6	NAG	E	907	1	-	2/6/23/26	0/1/1/1
5	MLI	B	301	-	-	2/4/4/4	-
5	MLI	B	302	-	-	2/4/4/4	-
5	MLI	B	303	-	-	2/4/4/4	-
8	PG0	A	902	-	-	3/5/5/5	-
7	1PE	L	1603	-	-	8/13/13/13	-
5	MLI	H	304	-	-	0/4/4/4	-
5	MLI	E	908	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	904	NAG	C1-C2	2.49	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	903	MLI	O7-C2-C1	2.17	121.48	114.54
5	E	908	MLI	O9-C3-C1	2.01	120.96	114.54

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	E	906	NAG	C8-C7-N2-C2
6	E	906	NAG	O7-C7-N2-C2
6	E	907	NAG	C8-C7-N2-C2
6	E	907	NAG	O7-C7-N2-C2
6	A	904	NAG	C8-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	304	MLI	1	0
7	L	1603	1PE	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/213 (93%)	0.46	7 (3%) 44 42	70, 90, 129, 153	0
1	E	203/213 (95%)	0.22	1 (0%) 91 89	60, 81, 120, 168	0
2	B	217/229 (94%)	0.64	22 (10%) 7 6	55, 89, 141, 179	0
2	H	216/229 (94%)	0.25	6 (2%) 53 50	56, 84, 135, 170	0
3	C	219/220 (99%)	0.58	15 (6%) 17 15	57, 86, 125, 144	0
3	L	219/220 (99%)	0.05	2 (0%) 84 82	58, 83, 109, 124	0
All	All	1273/1324 (96%)	0.37	53 (4%) 36 34	55, 86, 129, 179	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	218	LYS	5.0
3	C	131	LEU	4.8
2	B	215	VAL	4.6
1	A	445	VAL	4.5
2	B	195	THR	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MLI	B	304	7/7	0.54	0.41	116,123,137,137	0
4	DMS	L	1602	4/4	0.63	0.31	160,163,164,175	0
5	MLI	C	306	7/7	0.67	0.29	115,118,138,139	0
8	PG0	A	902	8/8	0.70	0.28	93,96,104,105	0
5	MLI	H	304	7/7	0.71	0.28	126,130,132,133	0
5	MLI	B	302	7/7	0.72	0.40	120,123,134,134	0
5	MLI	E	908	7/7	0.78	0.28	97,101,109,109	0
4	DMS	H	302	4/4	0.78	0.27	144,147,149,154	0
5	MLI	H	301	7/7	0.79	0.41	125,126,127,127	0
5	MLI	C	305	7/7	0.80	0.33	131,135,136,137	0
4	DMS	C	303	4/4	0.80	0.19	149,149,150,153	0
5	MLI	H	305	7/7	0.80	0.30	134,136,139,139	0
4	DMS	A	903	4/4	0.81	0.28	142,146,146,147	0
7	1PE	C	301	16/16	0.82	0.21	84,90,95,96	0
6	NAG	E	906	14/15	0.82	0.26	82,90,102,107	0
5	MLI	E	903	7/7	0.83	0.66	132,133,134,135	0
5	MLI	B	303	7/7	0.85	0.27	135,136,137,140	0
7	1PE	L	1603	16/16	0.85	0.21	81,91,108,109	0
6	NAG	E	907	14/15	0.86	0.16	80,99,120,127	0
5	MLI	C	304	7/7	0.88	0.62	123,124,128,129	0
4	DMS	C	302	4/4	0.88	0.20	136,136,138,138	0
5	MLI	B	301	7/7	0.89	0.28	107,109,113,113	0
4	DMS	A	905	4/4	0.89	0.39	148,149,151,160	0
4	DMS	E	904	4/4	0.90	0.68	141,142,143,144	0
4	DMS	L	1601	4/4	0.90	0.32	137,138,139,147	0
4	DMS	E	905	4/4	0.91	0.26	129,130,131,132	0
6	NAG	A	904	14/15	0.92	0.11	72,97,109,117	0
4	DMS	B	305	4/4	0.93	0.17	136,137,139,139	0
4	DMS	H	303	4/4	0.94	0.18	123,124,124,125	0
4	DMS	A	901	4/4	0.95	0.20	101,102,107,113	0
4	DMS	E	902	4/4	0.95	0.22	125,129,129,130	0
4	DMS	E	901	4/4	0.96	0.21	95,95,98,98	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.