



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 16, 2023 – 06:09 AM EDT

PDB ID : 1YQ2  
Title : beta-galactosidase from *Arthrobacter* sp. C2-2 (isoenzyme C2-2-1)  
Authors : Skalova, T.; Dohnalek, J.; Spiwok, V.; Lipovova, P.; Vondrackova, E.;  
Petrokova, H.; Strnad, H.; Kralova, B.; Hasek, J.  
Deposited on : 2005-02-01  
Resolution : 1.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1024	 2% 91% 8%
1	B	1024	 2% 92% 6%
1	C	1024	 3% 93% 6%
1	D	1024	 2% 91% 8%
1	E	1024	 2% 92% 7%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	1024	<p>A horizontal bar chart representing the quality of chain. The bar is primarily green, indicating a high quality score of 91%. There is a small red segment at the beginning, labeled '2%', and a small yellow segment at the end, labeled '8%'. The total length of the bar represents 100%.</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	A	8511	-	-	X	-
5	SO4	C	8513	-	-	X	-
5	SO4	F	8516	-	-	X	-
6	PEG	B	9002	-	-	X	-
6	PEG	C	9003	-	-	X	-
6	PEG	E	9005	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 53897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1020	7882	4949	1419	1493	21	0	17	0
1	B	1020	7870	4943	1422	1484	21	0	15	0
1	C	1020	7891	4955	1426	1488	22	0	19	0
1	D	1020	7870	4942	1416	1491	21	0	14	0
1	E	1020	7888	4954	1426	1485	23	0	17	0
1	F	1020	7904	4961	1436	1486	21	0	20	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q8KRF6
A	903	MET	VAL	engineered mutation	UNP Q8KRF6
B	0	HIS	-	expression tag	UNP Q8KRF6
B	903	MET	VAL	engineered mutation	UNP Q8KRF6
C	0	HIS	-	expression tag	UNP Q8KRF6
C	903	MET	VAL	engineered mutation	UNP Q8KRF6
D	0	HIS	-	expression tag	UNP Q8KRF6
D	903	MET	VAL	engineered mutation	UNP Q8KRF6
E	0	HIS	-	expression tag	UNP Q8KRF6
E	903	MET	VAL	engineered mutation	UNP Q8KRF6
F	0	HIS	-	expression tag	UNP Q8KRF6
F	903	MET	VAL	engineered mutation	UNP Q8KRF6

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Na 2 2	0	0
3	B	2	Total Na 2 2	0	0
3	C	2	Total Na 2 2	0	0
3	D	2	Total Na 2 2	0	0
3	E	2	Total Na 2 2	0	0
3	F	2	Total Na 2 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	1	Total Cl 1 1	0	0
4	C	1	Total Cl 1 1	0	0
4	D	1	Total Cl 1 1	0	0
4	E	1	Total Cl 1 1	0	0
4	F	1	Total Cl 1 1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	E	1	Total O S 5 4 1	0	0
5	E	1	Total O S 5 4 1	0	0
5	F	1	Total O S 5 4 1	0	0
5	F	1	Total O S 5 4 1	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0
6	C	1	Total C O 7 4 3	0	0
6	D	1	Total C O 7 4 3	0	0
6	E	1	Total C O 7 4 3	0	0
6	F	1	Total C O 7 4 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1119	Total O 1119 1119	0	0
7	B	1122	Total O 1122 1122	0	0
7	C	1108	Total O 1108 1108	0	0
7	D	1089	Total O 1089 1089	0	0
7	E	969	Total O 969 969	0	0
7	F	1064	Total O 1064 1064	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: beta-galactosidase




- Molecule 1: beta-galactosidase




- Molecule 1: beta-galactosidase










- Molecule 1: beta-galactosidase




- Molecule 1: beta-galactosidase



- Molecule 1: beta-galactosidase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.09Å 205.70Å 140.46Å 90.00° 102.34° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 39.98 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.0 (40.00-1.90) 95.0 (39.98-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2	Depositor
R, $R_{free}$	0.157 , 0.195 0.158 , 0.167	Depositor DCC
$R_{free}$ test set	5792 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtrriage
Anisotropy	0.338	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.015 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	53897	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NA, SO4, CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/8190	0.72	9/11200 (0.1%)
1	B	0.67	0/8173	0.75	8/11176 (0.1%)
1	C	0.65	0/8210	0.73	11/11226 (0.1%)
1	D	0.66	0/8157	0.71	6/11157 (0.1%)
1	E	0.65	0/8198	0.72	2/11207 (0.0%)
1	F	0.66	0/8226	0.72	4/11245 (0.0%)
All	All	0.66	0/49154	0.73	40/67211 (0.1%)

There are no bond length outliers.

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	ARG	NE-CZ-NH1	-8.44	116.08	120.30
1	B	277	LEU	CA-CB-CG	8.17	134.08	115.30
1	D	79	LEU	CA-CB-CG	7.26	132.01	115.30
1	A	377	LEU	CA-CB-CG	-7.15	98.86	115.30
1	C	277	LEU	CA-CB-CG	7.06	131.54	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7882	0	7501	63	0
1	B	7870	0	7495	48	0
1	C	7891	0	7512	45	0
1	D	7870	0	7479	57	0
1	E	7888	0	7517	54	0
1	F	7904	0	7537	63	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	10	0	0	3	0
5	B	10	0	0	0	0
5	C	5	0	0	3	0
5	D	10	0	0	0	0
5	E	10	0	0	0	0
5	F	10	0	0	4	0
6	A	7	0	10	3	0
6	B	7	0	10	4	0
6	C	7	0	10	4	0
6	D	7	0	10	3	0
6	E	7	0	10	6	0
6	F	7	0	10	3	0
7	A	1119	0	0	7	1
7	B	1122	0	0	10	0
7	C	1108	0	0	11	0
7	D	1089	0	0	8	0
7	E	969	0	0	4	0
7	F	1064	0	0	13	1
All	All	53897	0	45101	340	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 340 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:8516:SO4:O1	7:F:9184:HOH:O	1.52	1.26
1:C:130[B]:GLN:HG3	7:C:9125:HOH:O	1.27	1.24
1:F:773:ARG:NH1	7:F:9942:HOH:O	1.73	1.21
5:A:8511:SO4:S	7:A:9895:HOH:O	2.03	1.14
1:C:325[A]:ARG:NH1	7:C:9898:HOH:O	1.79	1.14

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:9663:HOH:O	7:F:9129:HOH:O[2_545]	1.95	0.25

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1035/1024 (101%)	1005 (97%)	30 (3%)	0	100	100
1	B	1033/1024 (101%)	1004 (97%)	29 (3%)	0	100	100
1	C	1037/1024 (101%)	1006 (97%)	30 (3%)	1 (0%)	51	42
1	D	1032/1024 (101%)	1002 (97%)	30 (3%)	0	100	100
1	E	1035/1024 (101%)	1007 (97%)	28 (3%)	0	100	100
1	F	1038/1024 (101%)	1003 (97%)	35 (3%)	0	100	100
All	All	6210/6144 (101%)	6027 (97%)	182 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	523	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	815/803 (102%)	792 (97%)	23 (3%)	43	36
1	B	813/803 (101%)	789 (97%)	24 (3%)	41	33
1	C	816/803 (102%)	802 (98%)	14 (2%)	60	57
1	D	811/803 (101%)	789 (97%)	22 (3%)	44	38
1	E	816/803 (102%)	792 (97%)	24 (3%)	42	35
1	F	817/803 (102%)	800 (98%)	17 (2%)	53	48
All	All	4888/4818 (102%)	4764 (98%)	124 (2%)	49	41

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	622	LYS
1	F	234	ARG
1	D	311	ARG
1	F	166	ARG
1	F	512	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	130	GLN
1	F	512	GLN
1	D	755	GLN
1	F	330	HIS
1	F	584	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 24 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	SO4	B	8502	-	4,4,4	0.14	0	6,6,6	0.43	0
6	PEG	F	9006	-	6,6,6	0.43	0	5,5,5	0.58	0
5	SO4	F	8506	-	4,4,4	0.17	0	6,6,6	0.36	0
5	SO4	D	8504	-	4,4,4	0.21	0	6,6,6	0.29	0
6	PEG	A	9001	-	6,6,6	0.55	0	5,5,5	0.33	0
5	SO4	A	8511	-	4,4,4	0.19	0	6,6,6	0.31	0
6	PEG	E	9005	-	6,6,6	0.72	0	5,5,5	0.63	0
5	SO4	E	8505	-	4,4,4	0.17	0	6,6,6	0.33	0
5	SO4	D	8514	-	4,4,4	0.15	0	6,6,6	0.31	0
5	SO4	E	8515	-	4,4,4	0.14	0	6,6,6	0.46	0
5	SO4	F	8516	-	4,4,4	0.12	0	6,6,6	0.27	0
6	PEG	C	9003	-	6,6,6	0.60	0	5,5,5	0.61	0
6	PEG	D	9004	-	6,6,6	0.55	0	5,5,5	0.40	0
5	SO4	A	8501	-	4,4,4	0.06	0	6,6,6	0.23	0
5	SO4	C	8513	-	4,4,4	0.14	0	6,6,6	0.25	0
5	SO4	B	8512	-	4,4,4	0.16	0	6,6,6	0.25	0
6	PEG	B	9002	-	6,6,6	0.60	0	5,5,5	0.43	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	F	9006	-	-	2/4/4/4	-
6	PEG	A	9001	-	-	2/4/4/4	-
6	PEG	E	9005	-	-	2/4/4/4	-
6	PEG	C	9003	-	-	1/4/4/4	-
6	PEG	D	9004	-	-	2/4/4/4	-
6	PEG	B	9002	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	9001	PEG	O2-C3-C4-O4
6	D	9004	PEG	O2-C3-C4-O4
6	E	9005	PEG	O1-C1-C2-O2
6	E	9005	PEG	O2-C3-C4-O4
6	F	9006	PEG	O1-C1-C2-O2

There are no ring outliers.

10 monomers are involved in 33 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	F	9006	PEG	3	0
5	F	8506	SO4	1	0
6	A	9001	PEG	3	0
5	A	8511	SO4	3	0
6	E	9005	PEG	6	0
5	F	8516	SO4	3	0
6	C	9003	PEG	4	0
6	D	9004	PEG	3	0
5	C	8513	SO4	3	0
6	B	9002	PEG	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1020/1024 (99%)	-0.42	25 (2%) 57 60	8, 16, 31, 48	6 (0%)
1	B	1020/1024 (99%)	-0.46	23 (2%) 60 63	7, 15, 30, 49	6 (0%)
1	C	1020/1024 (99%)	-0.47	27 (2%) 56 58	7, 14, 30, 46	11 (1%)
1	D	1020/1024 (99%)	-0.41	25 (2%) 57 60	6, 15, 30, 47	8 (0%)
1	E	1020/1024 (99%)	-0.46	19 (1%) 66 69	7, 15, 29, 47	20 (1%)
1	F	1020/1024 (99%)	-0.48	19 (1%) 66 69	7, 15, 30, 45	15 (1%)
All	All	6120/6144 (99%)	-0.45	138 (2%) 60 63	6, 15, 30, 49	66 (1%)

The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	236	GLY	6.9
1	E	235	SER	6.8
1	C	237	ALA	6.1
1	F	235	SER	6.1
1	A	236	GLY	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	PEG	A	9001	7/7	0.73	0.23	38,39,41,41	0
6	PEG	C	9003	7/7	0.79	0.20	28,30,33,36	0
6	PEG	E	9005	7/7	0.82	0.17	28,29,33,33	0
5	SO4	F	8516	5/5	0.84	0.25	33,33,36,36	5
6	PEG	D	9004	7/7	0.85	0.18	31,33,35,35	0
6	PEG	F	9006	7/7	0.85	0.15	31,35,36,37	0
6	PEG	B	9002	7/7	0.86	0.16	28,32,33,34	0
5	SO4	E	8515	5/5	0.88	0.21	26,26,29,29	5
3	NA	B	7502	1/1	0.90	0.12	44,44,44,44	0
5	SO4	C	8513	5/5	0.90	0.23	37,38,40,40	5
3	NA	D	7504	1/1	0.91	0.18	39,39,39,39	0
3	NA	F	7506	1/1	0.91	0.21	47,47,47,47	0
5	SO4	B	8512	5/5	0.91	0.18	30,30,32,32	5
3	NA	C	7503	1/1	0.91	0.08	41,41,41,41	0
5	SO4	D	8514	5/5	0.92	0.16	31,31,32,32	5
5	SO4	A	8511	5/5	0.92	0.15	23,25,27,28	5
5	SO4	D	8504	5/5	0.92	0.13	54,55,55,56	0
5	SO4	F	8506	5/5	0.94	0.10	34,34,35,35	5
5	SO4	E	8505	5/5	0.96	0.10	40,41,42,43	0
3	NA	A	7501	1/1	0.97	0.14	38,38,38,38	0
4	CL	A	8011	1/1	0.97	0.04	30,30,30,30	0
5	SO4	A	8501	5/5	0.97	0.11	29,30,31,31	5
3	NA	C	7513	1/1	0.97	0.06	29,29,29,29	0
3	NA	B	7512	1/1	0.97	0.06	32,32,32,32	0
3	NA	E	7505	1/1	0.97	0.22	38,38,38,38	0
4	CL	F	8016	1/1	0.98	0.03	32,32,32,32	0
3	NA	A	7511	1/1	0.98	0.07	30,30,30,30	0
2	MG	C	7003	1/1	0.98	0.04	19,19,19,19	0
5	SO4	B	8502	5/5	0.98	0.14	30,30,32,33	0
3	NA	F	7516	1/1	0.98	0.06	33,33,33,33	0
3	NA	D	7514	1/1	0.98	0.05	30,30,30,30	0
4	CL	B	8012	1/1	0.99	0.03	27,27,27,27	0
4	CL	C	8013	1/1	0.99	0.03	30,30,30,30	0
4	CL	D	8014	1/1	0.99	0.05	28,28,28,28	0
3	NA	E	7515	1/1	0.99	0.14	31,31,31,31	0
2	MG	B	7002	1/1	0.99	0.04	18,18,18,18	0
2	MG	A	7001	1/1	0.99	0.03	17,17,17,17	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	D	7004	1/1	0.99	0.04	20,20,20,20	0
2	MG	E	7005	1/1	1.00	0.03	15,15,15,15	0
2	MG	F	7006	1/1	1.00	0.02	17,17,17,17	0
4	CL	E	8015	1/1	1.00	0.02	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.