



## Full wwPDB X-ray Structure Validation Report ⓘ

Sep 4, 2024 – 04:10 PM JST

PDB ID : 8YQ8  
Title : Quadruple mutant (N51I+C59R+S108N+I164L) Plasmodium falciparum dihydrofolate reductase-thymidylate synthase (PfDHFR-TS V1/S) complexed with FB8, NADPH and dUMP  
Authors : Vanichtanankul, J.; Vitsupakorn, D.; Saeyang, T.; Arwon, U.; Hoarau, M.; Decharuangsilp, S.; Kamchonwongpaisan, S.; Yuthavong, Y.  
Deposited on : 2024-03-19  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	608	 5% 78% 8% 14%
1	B	608	 10% 75% 9% 15%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional dihydrofolate reductase-thymidylate synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	524	Total 4364	C 2828	N 718	O 793	S 25	0	0	0
1	B	518	Total 4312	C 2799	N 712	O 777	S 24	0	0	0

- Molecule 2 is 4-[4-[4-[2,4-bis(azanyl)-6-ethyl-pyrimidin-5-yl]oxybutoxy]phenyl]benzoic acid (three-letter code: A1LZS) (formula: C<sub>23</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 31	C 23	N 4	O 4	0	0
2	B	1	Total 31	C 23	N 4	O 4	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	48	21	7	17	3	0	0
3	B	1	48	21	7	17	3	0	0

- Molecule 4 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $C_9H_{13}N_2O_8P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
4	A	1	20	9	2	8	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	B	1	20	9	2	8	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	314	Total	O	0	0
			314	314		
5	B	274	Total	O	0	0
			274	274		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.76Å 156.41Å 165.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.10 – 2.10 26.10 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (26.10-2.10) 99.4 (26.10-2.10)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.195 , 0.240 0.204 , 0.245	Depositor DCC
$R_{free}$ test set	4451 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 61.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9462	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, A1LZS, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/4467	0.96	10/6034 (0.2%)
1	B	0.49	0/4413	0.92	3/5961 (0.1%)
All	All	0.51	0/8880	0.94	13/11995 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	566	ARG	NE-CZ-NH1	6.68	123.64	120.30
1	B	421	ARG	NE-CZ-NH2	-6.25	117.18	120.30
1	A	471	ARG	NE-CZ-NH2	-6.10	117.25	120.30
1	A	59	ARG	NE-CZ-NH2	5.88	123.24	120.30
1	A	59	ARG	CD-NE-CZ	5.59	131.43	123.60
1	A	471	ARG	CD-NE-CZ	5.58	131.41	123.60
1	A	566	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	A	82	ASN	CB-CA-C	5.42	121.23	110.40
1	A	510	ARG	NE-CZ-NH2	-5.07	117.77	120.30
1	B	377	ARG	NE-CZ-NH1	5.07	122.83	120.30
1	A	122	ARG	NE-CZ-NH1	5.04	122.82	120.30
1	B	473	LEU	CB-CG-CD1	-5.04	102.43	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	494	CYS	CB-CA-C	-5.02	100.35	110.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	402	ARG	Sidechain
1	B	345	ARG	Sidechain
1	B	566	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4364	0	4317	26	0
1	B	4312	0	4287	32	0
2	A	31	0	0	0	0
2	B	31	0	0	0	0
3	A	48	0	26	0	0
3	B	48	0	26	2	0
4	A	20	0	11	1	0
4	B	20	0	11	2	0
5	A	314	0	0	4	0
5	B	274	0	0	4	0
All	All	9462	0	8678	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:SER:HB3	3:B:702:NDP:O2N	1.93	0.66
1:B:346:THR:HB	1:B:348:VAL:HG23	1.78	0.64
1:A:70:TYR:CE2	1:A:74:LYS:HE3	2.37	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:491:HIS:CE1	4:B:703:UMP:O4	2.55	0.59
1:B:112:ILE:HB	1:B:117:LYS:HD3	1.84	0.58
1:A:605:ASP:O	1:A:606:MET:C	2.43	0.56
1:A:332:ILE:HD13	1:A:560:LEU:HD22	1.87	0.56
1:A:307:ILE:HB	1:A:337:MET:CE	2.36	0.55
1:A:328:TYR:CZ	1:A:332:ILE:HD11	2.42	0.54
1:B:312:PHE:HA	1:B:565:ASN:HD21	1.73	0.53
1:B:487:LEU:N	1:B:487:LEU:HD23	2.23	0.53
1:A:437:PHE:CE2	1:A:478:ASN:HB2	2.43	0.53
1:B:582:ASN:HB3	5:B:814:HOH:O	2.11	0.51
1:B:309:PRO:HA	1:B:312:PHE:CD2	2.46	0.50
1:B:328:TYR:CZ	1:B:332:ILE:HD11	2.47	0.50
1:A:375:PHE:HD1	5:A:1035:HOH:O	1.94	0.50
1:B:309:PRO:HA	1:B:312:PHE:HD2	1.78	0.49
1:B:59:ARG:O	1:B:63:THR:HG23	2.13	0.48
1:A:354:PHE:CE2	1:B:506:ILE:HG13	2.49	0.48
1:B:114:LYS:HA	1:B:117:LYS:HE2	1.96	0.47
1:A:308:HIS:O	1:A:311:ASP:HB2	2.15	0.47
1:B:373:LYS:HG3	1:B:598:HIS:CE1	2.50	0.47
1:A:506:ILE:HG13	1:B:354:PHE:CE2	2.50	0.46
1:B:332:ILE:HD13	1:B:560:LEU:HD22	1.97	0.46
1:A:8:VAL:O	1:A:76:LYS:HE3	2.17	0.45
1:A:109:TRP:CZ2	1:A:117:LYS:HD2	2.52	0.45
1:A:109:TRP:CE2	1:A:117:LYS:HD2	2.52	0.45
1:A:81:LEU:O	1:A:82:ASN:HB2	2.17	0.45
1:B:490:CYS:SG	4:B:703:UMP:C6	3.09	0.45
1:B:333:TYR:CE1	1:B:337:MET:HE3	2.52	0.44
1:A:470:ARG:HD3	5:B:948:HOH:O	2.17	0.44
1:B:376:LEU:HD12	1:B:593:ILE:HG13	1.99	0.44
1:B:145:LYS:HA	1:B:145:LYS:HE2	1.99	0.44
1:A:127:LEU:HD23	1:A:143:ILE:HG13	2.00	0.44
1:A:491:HIS:CE1	4:A:703:UMP:O4	2.71	0.43
1:B:473:LEU:HD12	1:B:473:LEU:N	2.34	0.43
1:B:312:PHE:HB3	1:B:315:TYR:HB3	2.01	0.43
1:B:106:ARG:HE	3:B:702:NDP:P2B	2.42	0.42
1:B:152:LEU:HD11	1:B:156:LEU:HD11	2.02	0.42
1:A:532:ILE:HD13	1:A:583:ILE:HD13	2.01	0.42
1:A:329:LEU:HD22	1:A:564:LEU:HD12	2.02	0.42
1:B:122:ARG:O	1:B:124:ASN:ND2	2.52	0.42
1:B:452:GLY:HA2	5:B:966:HOH:O	2.19	0.42
1:B:569:TYR:HB3	1:B:570:PRO:HD2	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:584:GLU:HG3	5:B:814:HOH:O	2.20	0.41
1:A:78:CYS:O	1:A:82:ASN:N	2.46	0.41
1:A:4:GLN:O	1:A:8:VAL:HG23	2.20	0.41
5:A:1029:HOH:O	1:B:470:ARG:HD3	2.21	0.41
1:A:133:LYS:HD3	5:A:1092:HOH:O	2.20	0.41
1:A:346:THR:HG22	1:A:348:VAL:HG23	2.03	0.41
1:B:17:CYS:HA	1:B:39:GLY:O	2.20	0.41
1:A:569:TYR:HB3	1:A:570:PRO:HD2	2.02	0.41
1:A:594:GLN:HB2	5:A:914:HOH:O	2.21	0.40
1:A:19:LYS:HG2	1:A:36:THR:HG22	2.03	0.40
1:B:307:ILE:HG22	1:B:337:MET:HE1	2.03	0.40
1:B:336:MET:CE	1:B:560:LEU:HB2	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	514/608 (84%)	495 (96%)	19 (4%)	0	100	100
1	B	506/608 (83%)	490 (97%)	15 (3%)	1 (0%)	44	45
All	All	1020/1216 (84%)	985 (97%)	34 (3%)	1 (0%)	48	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	429	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/570 (86%)	485 (99%)	5 (1%)	73	79
1	B	484/570 (85%)	468 (97%)	16 (3%)	33	36
All	All	974/1140 (85%)	953 (98%)	21 (2%)	47	53

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	LYS
1	A	218	ASN
1	A	307	ILE
1	A	375	PHE
1	A	487	LEU
1	B	97	LYS
1	B	114	LYS
1	B	115	LYS
1	B	131	LEU
1	B	132	LYS
1	B	133	LYS
1	B	145	LYS
1	B	189	SER
1	B	295	PHE
1	B	310	ASN
1	B	345	ARG
1	B	573	THR
1	B	582	ASN
1	B	600	GLU
1	B	601	LYS
1	B	602	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	171	GLN
1	A	394	ASN
1	A	424	ASN
1	A	554	ASN
1	B	316	ASN
1	B	394	ASN
1	B	407	ASN
1	B	424	ASN
1	B	554	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NDP	A	702	-	45,52,52	0.69	1 (2%)	53,80,80	0.87	1 (1%)
3	NDP	B	702	-	45,52,52	0.70	1 (2%)	53,80,80	0.85	1 (1%)
4	UMP	A	703	-	21,21,21	0.76	0	31,31,31	1.11	3 (9%)
2	A1LZS	B	701	-	32,33,33	0.50	0	41,44,44	1.22	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	UMP	B	703	-	21,21,21	0.81	1 (4%)	31,31,31	1.22	3 (9%)
2	A1LZS	A	701	-	32,33,33	0.93	1 (3%)	41,44,44	1.01	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	A	702	-	-	2/30/77/77	0/5/5/5
3	NDP	B	702	-	-	5/30/77/77	0/5/5/5
4	UMP	A	703	-	-	1/10/22/22	0/2/2/2
2	A1LZS	B	701	-	-	4/19/19/19	0/3/3/3
4	UMP	B	703	-	-	1/10/22/22	0/2/2/2
2	A1LZS	A	701	-	-	5/19/19/19	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	A1LZS	C6-C3	-3.66	1.34	1.40
3	B	702	NDP	P2B-O2B	2.54	1.64	1.59
4	B	703	UMP	P-OP3	-2.48	1.45	1.54
3	A	702	NDP	P2B-O2X	-2.07	1.46	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	A1LZS	C7-O1-C6	6.68	134.41	114.23
2	A	701	A1LZS	C7-O1-C6	3.49	124.77	114.23
4	B	703	UMP	O2-C2-N1	3.27	127.14	122.79
4	A	703	UMP	O2-C2-N1	2.94	126.69	122.79
4	B	703	UMP	C1'-N1-C6	-2.83	115.97	121.55
3	A	702	NDP	C5A-C6A-N6A	2.65	124.38	120.35
2	A	701	A1LZS	C4-N1-C3	2.36	118.21	116.24
3	B	702	NDP	O2A-PA-O1A	2.23	123.26	112.24
4	B	703	UMP	C1'-N1-C2	2.15	121.88	117.64
4	A	703	UMP	C1'-N1-C2	2.03	121.63	117.64
4	A	703	UMP	OP3-P-OP2	2.02	115.34	107.64

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	A1LZS	C5-C6-O1-C7
2	A	701	A1LZS	C3-C6-O1-C7
2	B	701	A1LZS	C5-C6-O1-C7
2	B	701	A1LZS	C3-C6-O1-C7
2	B	701	A1LZS	O1-C7-C8-C9
2	A	701	A1LZS	O1-C7-C8-C9
2	B	701	A1LZS	O2-C10-C9-C8
2	A	701	A1LZS	C8-C7-O1-C6
3	B	702	NDP	C2D-C1D-N1N-C2N
2	A	701	A1LZS	C7-C8-C9-C10
3	B	702	NDP	O4D-C1D-N1N-C2N
3	A	702	NDP	O4D-C1D-N1N-C2N
3	A	702	NDP	C2D-C1D-N1N-C2N
3	B	702	NDP	C2D-C1D-N1N-C6N
4	A	703	UMP	O4'-C4'-C5'-O5'
3	B	702	NDP	O4D-C1D-N1N-C6N
3	B	702	NDP	C2B-O2B-P2B-O1X
4	B	703	UMP	O4'-C4'-C5'-O5'


There are no ring outliers.



3 monomers are involved in 5 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	702	NDP	2	0
4	A	703	UMP	1	0
4	B	703	UMP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	524/608 (86%)	0.12	29 (5%) 32 34	14, 27, 64, 105	0
1	B	518/608 (85%)	0.50	63 (12%) 10 10	14, 33, 74, 99	0
All	All	1042/1216 (85%)	0.31	92 (8%) 17 18	14, 29, 71, 105	0

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	307	ILE	5.3
1	B	346	THR	4.5
1	B	107	THR	4.5
1	B	131	LEU	4.4
1	B	75	TYR	4.0
1	B	348	VAL	3.8
1	A	343	SER	3.7
1	B	106	ARG	3.6
1	B	168	VAL	3.6
1	A	606	MET	3.5
1	B	143	ILE	3.5
1	B	127	LEU	3.5
1	A	348	VAL	3.5
1	B	114	LYS	3.4
1	A	346	THR	3.4
1	B	132	LYS	3.3
1	A	22	SER	3.3
1	B	312	PHE	3.3
1	B	285	GLU	3.2
1	B	307	ILE	3.2
1	B	347	GLY	3.2
1	B	140	VAL	3.2
1	A	306	SER	3.1
1	A	75	TYR	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	310	ASN	3.0
1	B	602	ILE	2.9
1	B	97	LYS	2.9
1	B	118	PRO	2.9
1	B	45	VAL	2.9
1	B	605	ASP	2.9
1	B	309	PRO	2.8
1	A	347	GLY	2.8
1	B	130	THR	2.8
1	A	308	HIS	2.8
1	B	345	ARG	2.8
1	A	304	LYS	2.8
1	B	116	PHE	2.7
1	B	105	GLY	2.7
1	B	126	ILE	2.7
1	B	108	ASN	2.7
1	B	109	TRP	2.7
1	A	309	PRO	2.7
1	B	110	GLU	2.7
1	A	8	VAL	2.7
1	A	82	ASN	2.6
1	B	144	ASN	2.6
1	B	98	LEU	2.6
1	B	555	ASN	2.6
1	A	4	GLN	2.6
1	A	284	ASP	2.6
1	A	312	PHE	2.6
1	B	193	CYS	2.6
1	B	308	HIS	2.6
1	B	112	ILE	2.5
1	A	72	LYS	2.5
1	B	9	PHE	2.5
1	A	285	GLU	2.5
1	B	111	SER	2.5
1	B	117	LYS	2.5
1	A	349	GLY	2.5
1	A	400	ASN	2.4
1	B	74	LYS	2.4
1	B	87	ASP	2.4
1	B	133	LYS	2.4
1	B	113	PRO	2.4
1	A	293	PHE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	344	ASP	2.4
1	B	205	TYR	2.4
1	A	6	CYS	2.3
1	B	129	ARG	2.3
1	A	311	ASP	2.3
1	B	104	MET	2.3
1	B	128	SER	2.3
1	A	97	LYS	2.2
1	B	20	VAL	2.2
1	B	80	TYR	2.2
1	B	306	SER	2.2
1	A	605	ASP	2.2
1	B	229	THR	2.1
1	B	167	SER	2.1
1	A	283	ASP	2.1
1	A	44	GLY	2.1
1	B	121	ASN	2.1
1	B	3	GLU	2.1
1	B	600	GLU	2.1
1	B	47	PRO	2.1
1	B	82	ASN	2.1
1	A	344	ASP	2.1
1	B	43	LYS	2.0
1	B	169	VAL	2.0
1	B	81	LEU	2.0
1	B	151	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	A1LZS	B	701	31/31	0.83	0.15	36,66,100,101	0
3	NDP	B	702	48/48	0.85	0.13	37,58,82,85	0
2	A1LZS	A	701	31/31	0.93	0.10	17,36,60,65	0
4	UMP	B	703	20/20	0.94	0.11	27,39,50,50	0
4	UMP	A	703	20/20	0.96	0.09	25,35,42,45	0
3	NDP	A	702	48/48	0.97	0.07	21,27,33,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around NDP B 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around A1LZS A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around UMP B 703:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UMP A 703:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.