



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2023 – 08:21 PM EDT

PDB ID : 1YTV  
Title : Maltose-binding protein fusion to a C-terminal fragment of the V1a vasopressin receptor  
Authors : Adikesavan, N.V.; Mahmood, S.S.; Stanley, S.; Xu, Z.; Wu, N.; Thibonnier, M.; Shoham, M.  
Deposited on : 2005-02-11  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	 2% 86% 14%
1	B	366	 3% 80% 19%
2	M	84	 2% 7% 93%
2	N	84	 4% 6% 94%
3	C	2	 100%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	D	2	 100%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	366	2842	1831	461	544	6	0	0	0
1	B	366	2842	1831	461	544	6	0	0	0

- Molecule 2 is a protein called Vasopressin V1a receptor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	M	6	42	21	9	12	0	0	0
2	N	5	34	17	7	10	0	0	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	381	ASN	-	linker	UNP P37288
M	382	SER	-	linker	UNP P37288
M	383	SER	-	linker	UNP P37288
M	384	SER	-	linker	UNP P37288
M	385	ASN	-	linker	UNP P37288
M	386	ASN	-	linker	UNP P37288
M	387	ASN	-	linker	UNP P37288
M	388	ASN	-	linker	UNP P37288
M	389	ASN	-	linker	UNP P37288
M	390	ASN	-	linker	UNP P37288
M	391	ASN	-	linker	UNP P37288
M	392	ASN	-	linker	UNP P37288
M	393	ASN	-	linker	UNP P37288
M	394	ASN	-	linker	UNP P37288
M	395	LEU	-	linker	UNP P37288
M	396	GLY	-	linker	UNP P37288

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	397	ILE	-	linker	UNP P37288
M	398	GLU	-	linker	UNP P37288
M	399	GLU	-	linker	UNP P37288
M	400	ASN	-	linker	UNP P37288
M	401	LEU	-	linker	UNP P37288
M	402	TYR	-	linker	UNP P37288
M	403	PHE	-	linker	UNP P37288
M	404	GLN	-	linker	UNP P37288
M	405	GLY	-	linker	UNP P37288
N	381	ASN	-	linker	UNP P37288
N	382	SER	-	linker	UNP P37288
N	383	SER	-	linker	UNP P37288
N	384	SER	-	linker	UNP P37288
N	385	ASN	-	linker	UNP P37288
N	386	ASN	-	linker	UNP P37288
N	387	ASN	-	linker	UNP P37288
N	388	ASN	-	linker	UNP P37288
N	389	ASN	-	linker	UNP P37288
N	390	ASN	-	linker	UNP P37288
N	391	ASN	-	linker	UNP P37288
N	392	ASN	-	linker	UNP P37288
N	393	ASN	-	linker	UNP P37288
N	394	ASN	-	linker	UNP P37288
N	395	LEU	-	linker	UNP P37288
N	396	GLY	-	linker	UNP P37288
N	397	ILE	-	linker	UNP P37288
N	398	GLU	-	linker	UNP P37288
N	399	GLU	-	linker	UNP P37288
N	400	ASN	-	linker	UNP P37288
N	401	LEU	-	linker	UNP P37288
N	402	TYR	-	linker	UNP P37288
N	403	PHE	-	linker	UNP P37288
N	404	GLN	-	linker	UNP P37288
N	405	GLY	-	linker	UNP P37288

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			22	12	10			
3	D	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	337	Total	O	0	0
			337	337		
4	M	6	Total	O	0	0
			6	6		
4	B	296	Total	O	0	0
			296	296		
4	N	2	Total	O	0	0
			2	2		







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.10Å 66.56Å 115.72Å 90.00° 95.99° 90.00°	Depositor
Resolution (Å)	34.43 – 1.80 34.43 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (34.43-1.80) 99.6 (34.43-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 1.81Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.199 , 0.225 0.208 , 0.232	Depositor DCC
$R_{free}$ test set	3609 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.7	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 56.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6445	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/2911	0.54	0/3952
1	B	0.32	0/2911	0.53	0/3952
2	M	0.30	0/41	0.45	0/54
2	N	0.44	0/33	0.74	0/43
All	All	0.32	0/5896	0.54	0/8001

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2842	0	2812	39	0
1	B	2842	0	2812	59	0
2	M	42	0	32	0	0
2	N	34	0	26	0	0
3	C	22	0	19	0	0
3	D	22	0	19	0	0
4	A	337	0	0	8	0
4	B	296	0	0	10	0
4	M	6	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	N	2	0	0	0	0
All	All	6445	0	5720	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ILE:HG22	1:A:291:LYS:HE3	1.40	1.02
1:B:16:ILE:H	1:B:16:ILE:HD12	1.35	0.89
1:A:314:GLY:HA3	4:A:892:HOH:O	1.79	0.81
1:A:136:LEU:HD21	1:A:140:PRO:HD3	1.68	0.76
1:A:345:PRO:HG2	1:A:350:MET:SD	2.26	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/366 (100%)	360 (99%)	4 (1%)	0	100	100
1	B	364/366 (100%)	357 (98%)	7 (2%)	0	100	100
2	M	4/84 (5%)	2 (50%)	2 (50%)	0	100	100
2	N	3/84 (4%)	3 (100%)	0	0	100	100
All	All	735/900 (82%)	722 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/293 (100%)	290 (99%)	3 (1%)	76	71
1	B	293/293 (100%)	287 (98%)	6 (2%)	55	44
2	M	6/80 (8%)	6 (100%)	0	100	100
2	N	5/80 (6%)	5 (100%)	0	100	100
All	All	597/746 (80%)	588 (98%)	9 (2%)	65	56

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	323	GLU
1	B	349	GLN
1	B	174	LEU
1	B	192	ILE
1	B	216	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	296	ASN
2	N	381	ASN
1	B	349	GLN
1	A	339	GLN
1	B	286	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GLC	C	1	3	11,11,12	1.25	1 (9%)	15,15,17	2.03	4 (26%)
3	GLC	C	2	3	11,11,12	0.92	0	15,15,17	1.48	2 (13%)
3	GLC	D	1	3	11,11,12	1.26	1 (9%)	15,15,17	2.03	4 (26%)
3	GLC	D	2	3	11,11,12	1.04	0	15,15,17	1.25	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	C	1	3	-	0/2/19/22	0/1/1/1
3	GLC	C	2	3	-	0/2/19/22	0/1/1/1
3	GLC	D	1	3	-	0/2/19/22	0/1/1/1
3	GLC	D	2	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	GLC	O5-C5	2.39	1.48	1.43
3	D	1	GLC	O5-C5	2.35	1.48	1.43

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	GLC	C1-O5-C5	3.96	117.56	112.19
3	C	1	GLC	C2-C3-C4	-3.88	104.19	110.89
3	D	1	GLC	O2-C2-C3	-3.79	102.54	110.14
3	D	1	GLC	C2-C3-C4	-3.73	104.44	110.89
3	C	1	GLC	O2-C2-C3	-3.71	102.71	110.14


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	366/366 (100%)	0.05	9 (2%) 57 52	11, 19, 35, 50	0
1	B	366/366 (100%)	0.10	11 (3%) 50 44	11, 19, 36, 56	0
2	M	6/84 (7%)	2.53	2 (33%) 0 0	22, 36, 63, 68	0
2	N	5/84 (5%)	2.79	3 (60%) 0 0	26, 27, 50, 54	0
All	All	743/900 (82%)	0.12	25 (3%) 45 39	11, 19, 36, 68	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	LYS	10.7
1	B	16	ILE	9.4
2	N	385	ASN	7.6
2	M	386	ASN	6.4
1	B	68	GLY	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GLC	D	1	11/12	0.91	0.11	12,14,17,17	0
3	GLC	C	1	11/12	0.92	0.12	12,15,18,22	0


*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GLC	C	2	11/12	0.97	0.07	10,11,13,14	0
3	GLC	D	2	11/12	0.98	0.08	8,10,13,14	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.