

Full wwPDB X-ray Structure Validation Report (i)

Jan 15, 2024 – 12:53 pm GMT

PDB ID	:	6YWJ
Title	:	Arabidopsis aspartate transcarbamoylase mutant F161A complex with UMP
Authors	:	Ramon Maiques, S.; Del Cano Ochoa, F.; Bellin, L.; Mohlmann, T.
Deposited on		
Resolution	:	2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4 02b-467
·		
Mogul	:	1.8.4, CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	332	78%	15%	• 6%
1	В	332	76%	17%	• 6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4946 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	312	Total	С	Ν	0	\mathbf{S}	0	0	0
		312	2448	1549	416	471	12	0		
1	В	211	Total	С	Ν	0	S	0	0	0
I B	311	2436	1541	413	470	12	0	0	U	

• Molecule 1 is a protein called PYRB.

There are 48 discrepancies between th	e modelled and reference sequences:
---------------------------------------	-------------------------------------


Chain	Residue	Modelled	Actual	Comment	Reference
А	59	MET	-	initiating methionine	UNP A0A178VJE3
А	60	GLY	_	expression tag	UNP A0A178VJE3
А	61	SER	_	expression tag	UNP A0A178VJE3
А	62	SER	-	expression tag	UNP A0A178VJE3
А	63	HIS	-	expression tag	UNP A0A178VJE3
А	64	HIS	-	expression tag	UNP A0A178VJE3
А	65	HIS	-	expression tag	UNP A0A178VJE3
А	66	HIS	-	expression tag	UNP A0A178VJE3
А	67	HIS	-	expression tag	UNP A0A178VJE3
А	68	HIS	-	expression tag	UNP A0A178VJE3
А	69	SER	-	expression tag	UNP A0A178VJE3
А	70	SER	-	expression tag	UNP A0A178VJE3
А	71	GLY	-	expression tag	UNP A0A178VJE3
А	72	LEU	-	expression tag	UNP A0A178VJE3
А	73	GLU	-	expression tag	UNP A0A178VJE3
А	74	VAL	-	expression tag	UNP A0A178VJE3
А	75	LEU	-	expression tag	UNP A0A178VJE3
А	76	PHE	-	expression tag	UNP A0A178VJE3
А	77	GLN	-	expression tag	UNP A0A178VJE3
А	78	GLY	-	expression tag	UNP A0A178VJE3
А	79	PRO	-	expression tag	UNP A0A178VJE3
А	80	HIS	-	expression tag	UNP A0A178VJE3
А	81	MET	-	expression tag	UNP A0A178VJE3
А	161	ALA	PHE	conflict	UNP A0A178VJE3
В	59	MET	-	initiating methionine	UNP A0A178VJE3



01110

Chain	Residue	Modelled	Actual	Comment	Reference
В	60	GLY	-	expression tag	UNP A0A178VJE3
В	61	SER	-	expression tag	UNP A0A178VJE3
В	62	SER	-	expression tag	UNP A0A178VJE3
В	63	HIS	-	expression tag	UNP A0A178VJE3
В	64	HIS	-	expression tag	UNP A0A178VJE3
В	65	HIS	-	expression tag	UNP A0A178VJE3
В	66	HIS	-	expression tag	UNP A0A178VJE3
В	67	HIS	-	expression tag	UNP A0A178VJE3
В	68	HIS	-	expression tag	UNP A0A178VJE3
В	69	SER	-	expression tag	UNP A0A178VJE3
В	70	SER	-	expression tag	UNP A0A178VJE3
В	71	GLY	-	expression tag	UNP A0A178VJE3
В	72	LEU	-	expression tag	UNP A0A178VJE3
В	73	GLU	-	expression tag	UNP A0A178VJE3
В	74	VAL	-	expression tag	UNP A0A178VJE3
В	75	LEU	-	expression tag	UNP A0A178VJE3
В	76	PHE	-	expression tag	UNP A0A178VJE3
В	77	GLN	-	expression tag	UNP A0A178VJE3
В	78	GLY	-	expression tag	UNP A0A178VJE3
В	79	PRO	-	expression tag	UNP A0A178VJE3
В	80	HIS	-	expression tag	UNP A0A178VJE3
В	81	MET	-	expression tag	UNP A0A178VJE3
В	161	ALA	PHE	conflict	UNP A0A178VJE3


• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 3 is URIDINE-5'-MONOPHOSPHATE (three-letter code: U5P) (formula: $C_9H_{13}N_2O_9P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Λ	1	Total	С	Ν	0	Р	0	0
J A	1	21	9	2	9	1	0	0	
9	D	1	Total	С	Ν	Ο	Р	0	0
J	3 B	1	21	9	2	9	1	0	U

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	3	Total O 3 3	0	0
4	В	5	Total O 5 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:	78%	15% • 6%
MET GLY SER SER SER HIS HIS HIS HIS HIS CLU CLU CLU CLU CLV CLU CLV CLU CLV CLU CLV CLU CLV CLV CLV CLV CLV CLV CLV CLV CLV CLV	R95 R96 N97 1101 1101 1128 8134 8134 8134 8134 8136 7135 7136 7136 7136 7154 7154	0210 H215 H215 1221 1233 1233 1233 1236 7235 7235 7236
L244 R248 V250 V256 V256 V256 V256 V256 V228 V228 V228 V228 V2305 V2305 V2305 V306	L324 A325 K329 K329 K329 F330 1331 V331 V332 V339 M344 1346 1346 1346 1346 1346	1.00 0371 1.378 1.379 1.381 1.381 1.381 1.381 1.390
• Molecule 1: PYRB		
Chain B:	76%	17% • 6%
MET MET OLY SER SER HIS HIS HIS HIS SER CLU CLU CLU CLU CLU CLU CLU CLU CLU CLU	Less 1864 1887 1887 1888 1889 1893 193 193 193 193 1128 1128 1128 1128 1128 1128 1128 112	8163 1184 1184 1203 1203 1203 1203 1203 1203 1221
0222 1225 1225 1223 0234 0234 0235 1236 1236 1236 1236 1236 7266 7266 7266 7266 7266 7266 7266 7	E271 1272 1272 1226 2281 7282 7282 7307 5301 5301 7307 5314 5315 7316 7316 7316 7316 7316	F330 F331 F331 F334 F337 F337 F337 F337 F337 F341 F341 F341 F341 F341 F341 F341 F341
V368 1386 1386		

• Molecule 1: PYRB



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants	104.34Å 104.34Å 127.84Å	Denesiten
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.30 - 2.40	Depositor
Resolution (A)	48.30 - 2.40	EDS
% Data completeness	100.0 (48.30-2.40)	Depositor
(in resolution range)	99.8 (48.30-2.40)	EDS
R _{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.07 (at 2.39 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D	0.138 , 0.190	Depositor
R, R_{free}	0.141 , 0.178	DCC
R_{free} test set	1539 reflections (4.98%)	wwPDB-VP
Wilson B-factor $(Å^2)$	43.9	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 25.9	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.070 for h,-h-k,-l	Xtriage
Depented twinning fraction	0.483 for H, K, L	Depositor
Reported twinning fraction	0.517 for -K, -H, -L	Depositor
Outliers	0 of 30890 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4946	wwPDB-VP
Average B, all atoms $(Å^2)$	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, $\rm U5P$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.78	0/2486	0.97	1/3346~(0.0%)
1	В	0.81	3/2472~(0.1%)	0.93	0/3327
All	All	0.80	3/4958~(0.1%)	0.95	1/6673~(0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
1	В	163	SER	CA-CB	-5.75	1.44	1.52
1	В	156	GLU	CD-OE2	5.43	1.31	1.25
1	В	143	SER	CA-CB	-5.32	1.45	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	379	ARG	NE-CZ-NH2	-5.15	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2448	0	2476	29	0
1	В	2436	0	2463	28	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	А	6	0	8	0	0
2	В	6	0	8	0	0
3	А	21	0	11	1	0
3	В	21	0	11	0	0
4	А	3	0	0	0	0
4	В	5	0	0	0	0
All	All	4946	0	4977	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + 1	A + 0	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:318:GLU:OE1	1:A:318:GLU:N	2.21	0.72
1:B:301:SER:O	1:B:341:GLN:NE2	2.22	0.72
1:A:221:LEU:C	1:A:221:LEU:HD12	2.12	0.70
1:B:264:LYS:HE2	1:B:266:TYR:OH	1.95	0.66
1:B:221:LEU:C	1:B:221:LEU:HD12	2.18	0.64
1:A:236:ILE:HD12	1:A:238:VAL:HG23	1.80	0.63
1:B:280:LYS:NZ	1:B:292:GLU:OE2	2.31	0.63
1:B:89:GLU:OE1	1:B:91:LYS:HE3	1.98	0.63
1:B:203:ILE:HB	1:B:204:PRO:HD2	1.81	0.62
1:B:337:LEU:HD11	1:B:360:VAL:HG13	1.80	0.62
1:A:360:VAL:HG12	1:A:360:VAL:O	2.02	0.59
1:A:236:ILE:HD12	1:A:238:VAL:CG2	2.33	0.59
1:B:235:GLY:HA2	1:B:262:ASP:O	2.04	0.58
1:B:83:GLU:N	1:B:83:GLU:OE1	2.38	0.57
1:B:82:PHE:CE1	1:B:385:LEU:HD13	2.40	0.57
1:A:97:MET:O	1:A:101:ILE:HG13	2.05	0.56
1:A:235:GLY:HA2	1:A:262:ASP:O	2.05	0.56
1:A:85:SER:OG	1:A:86:ASP:OD2	2.24	0.54
1:B:233:LEU:HA	1:B:236:ILE:HD13	1.90	0.54
1:A:378:ILE:O	1:A:381:ALA:HB3	2.07	0.53
1:B:270:PRO:HG3	1:B:330:PHE:CE2	2.43	0.53
1:A:328:GLY:O	1:A:331:ILE:HG22	2.12	0.50
1:A:128:THR:HG21	1:A:138:ARG:HG3	1.92	0.50
1:B:93:PHE:O	1:B:259:LYS:HE3	2.12	0.50
1:B:265:ILE:HD12	1:B:288:VAL:HG11	1.94	0.50
1:A:221:LEU:HD12	1:A:221:LEU:O	2.13	0.49
1:A:360:VAL:O	1:A:360:VAL:CG1	2.61	0.48



A + am 1	A + a	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:332:VAL:HB	1:A:356:ILE:HG12	1.97	0.47
1:A:336:LEU:O	1:A:340:MET:HG3	2.16	0.46
1:A:136:ARG:HD2	1:A:136:ARG:C	2.35	0.46
1:A:304:ASP:O	1:A:344:ALA:HA	2.14	0.46
1:A:266:TYR:CD2	1:A:299:VAL:HG13	2.51	0.46
1:B:87:VAL:HG23	1:B:386:LEU:HD11	1.97	0.46
1:A:339:VAL:HG23	1:A:339:VAL:O	2.14	0.45
1:A:298:GLU:OE2	1:A:298:GLU:HA	2.17	0.45
1:B:329:LYS:HB3	1:B:329:LYS:HE2	1.81	0.45
1:B:264:LYS:HG2	1:B:266:TYR:CE1	2.52	0.44
1:A:325:ALA:O	1:A:329:LYS:NZ	2.38	0.44
1:B:82:PHE:CE2	1:B:84:LEU:HB2	2.53	0.43
1:B:225:THR:HB	1:B:347:MET:HE3	2.00	0.43
1:A:306:VAL:HB	1:A:346:ILE:HD13	2.00	0.43
1:A:233:LEU:O	1:A:236:ILE:HG12	2.18	0.42
1:A:324:GLU:HA	1:A:324:GLU:OE1	2.19	0.42
1:B:98:LEU:HD23	1:B:98:LEU:HA	1.86	0.42
1:B:280:LYS:CE	1:B:292:GLU:OE2	2.68	0.42
1:B:184:ILE:O	1:B:205:VAL:HA	2.19	0.42
1:B:241:VAL:HA	1:B:268:VAL:O	2.20	0.42
1:B:256:LEU:HD23	1:B:256:LEU:HA	1.95	0.42
1:B:104:VAL:O	1:B:108:MET:HG2	2.20	0.42
1:B:334:LYS:HA	1:B:334:LYS:HD2	1.86	0.41
1:A:248:ARG:NH1	3:A:402:U5P:O3'	2.51	0.41
1:A:368:TYR:O	1:A:371:GLN:HB3	2.21	0.41
1:B:272:ILE:HD11	1:B:316:PHE:HA	2.02	0.41
1:A:207:ASN:HD21	1:A:210:ASP:HB2	1.85	0.41
1:A:258:ALA:CB	1:A:288:VAL:HG21	2.52	0.40
1:B:222:ASP:OD2	1:B:307:TYR:OH	2.30	0.40
1:A:244:LEU:HD23	1:A:250:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	А	310/332~(93%)	297~(96%)	13 (4%)	0	100	100
1	В	309/332~(93%)	296 (96%)	13 (4%)	0	100	100
All	All	619/664~(93%)	593~(96%)	26 (4%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	262/279~(94%)	251~(96%)	11 (4%)	30 47
1	В	260/279~(93%)	247~(95%)	13~(5%)	24 40
All	All	522/558~(94%)	498 (95%)	24~(5%)	27 43

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	95	ARG
1	А	107	GLU
1	А	134	SER
1	А	154	THR
1	А	176	THR
1	А	215	HIS
1	А	221	LEU
1	А	237	SER
1	А	262	ASP
1	А	297	MET
1	А	368	TYR
1	В	81	MET
1	В	91	LYS
1	В	128	THR
1	В	195	ARG



Contr	naca from	i preui	bus puye
Mol	Chain	\mathbf{Res}	Type
1	В	215	HIS
1	В	221	LEU
1	В	261	LYS
1	В	282	TYR
1	В	314	GLU
1	В	327	ARG
1	В	329	LYS
1	В	331	ILE
1	В	368	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Res	Link	Bond lengths			Bond angles			
	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	GOL	В	401	-	$5,\!5,\!5$	0.23	0	$5,\!5,\!5$	0.40	0



Mal	Mol Type Chain	Chain	Dec	Link	Bo	ond leng	\mathbf{ths}	Bond angles		
INIOI		nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
3	U5P	А	402	-	22,22,22	0.38	0	33,33,33	0.95	2 (6%)
3	U5P	В	402	-	22,22,22	0.81	1 (4%)	33,33,33	0.81	0
2	GOL	А	401	-	$5,\!5,\!5$	0.31	0	$5,\!5,\!5$	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	В	401	-	-	2/4/4/4	-
3	U5P	А	402	-	-	3/10/26/26	0/2/2/2
3	U5P	В	402	-	-	3/10/26/26	0/2/2/2
2	GOL	А	401	-	-	4/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	В	402	U5P	P-O5'	2.01	1.66	1.60

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	402	U5P	O2'-C2'-C1'	-3.30	98.98	110.02
3	А	402	U5P	O5'-P-O1P	-2.30	100.03	106.47

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	401	GOL	C1-C2-C3-O3
3	В	402	U5P	C3'-C4'-C5'-O5'
3	В	402	U5P	O4'-C4'-C5'-O5'
2	А	401	GOL	O1-C1-C2-C3
3	А	402	U5P	O4'-C4'-C5'-O5'
2	А	401	GOL	O1-C1-C2-O2
2	А	401	GOL	O2-C2-C3-O3
2	В	401	GOL	O2-C2-C3-O3
3	А	402	U5P	C3'-C4'-C5'-O5'
3	А	402	U5P	C4'-C5'-O5'-P



Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	В	402	U5P	C4'-C5'-O5'-P
2	В	401	GOL	C1-C2-C3-O3

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mo	bl	Chain	Res	Type	Clashes	Symm-Clashes
3		А	402	U5P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	> #RSRZ>2		$OWAB(Å^2)$	Q < 0.9	
1	А	312/332~(93%)	-0.55	0	100	100	26, 44, 68, 90	0
1	В	311/332~(93%)	-0.57	0	100	100	30, 43, 64, 73	0
All	All	623/664~(93%)	-0.56	0	100	100	26, 44, 65, 90	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	GOL	А	401	6/6	0.96	0.12	$26,\!30,\!35,\!38$	0
2	GOL	В	401	6/6	0.98	0.11	28,33,34,34	0
3	U5P	В	402	21/21	0.98	0.10	24,33,38,41	0
3	U5P	А	402	21/21	0.99	0.11	22,35,39,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers



as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

