

wwPDB X-ray Structure Validation Summary Report (i)

Aug 16, 2023 – 05:41 AM EDT

PDB ID	:	1YXW
Title	:	A common binding site for disiallylactose and a tri-peptide in the C-fragment
		of tetanus neurotoxin
Authors	:	Jayaraman, S.; Eswaramoorthy, S.; Kumaran, D.; Swaminathan, S.
Deposited on		
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	441	71%	25%	•

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TYR	А	5292	-	-	Х	-
3	GLU	А	5293	-	-	Х	-



1YXW

2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 3802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Tetanus toxin (Tentoxylysin).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	441	$\begin{array}{c} \text{Total} \\ 3564 \end{array}$	C 2282	N 597	O 676	${ m S} 9$	0	0	0


• Molecule 2 is TYROSINE (three-letter code: TYR) (formula: $C_9H_{11}NO_3$).



Mol	Chain	Residues	At	ton	ıs		ZeroOcc	AltConf
2	А	1	Total 12	С 9	N 1	O 2	0	0


• Molecule 3 is GLUTAMIC ACID (three-letter code: GLU) (formula: $C_5H_9NO_4$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total 9	_	N 1	0 3	0	0
0	A	1	9	5	1	3	0	

• Molecule 4 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	Λ	1	Total	С	Ν	0	0	0
4	A	1	15	11	2	2	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	А	202	Total 202	O 202	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.



• Molecule 1: Tetanus toxin (Tentoxylysin)



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	66.76Å 79.18Å 90.27Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.59 - 2.20	Depositor
% Data completeness	91.7 (39.59-2.20)	Depositor
(in resolution range)	51.1 (55.55-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R _{sym}	0.12	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.224 , 0.274	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3802	wwPDB-VP
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	l Chain Bond lengths		Bond angles		
	Unam	RMSZ	RMSZ $\# Z > 5$		# Z > 5
1	А	0.38	1/3645~(0.0%)	0.66	3/4945~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

\mathbf{Mol}	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	1183	ASN	C-N	-7.80	1.16	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	1183	ASN	O-C-N	-10.50	105.90	122.70
1	А	1183	ASN	C-N-CA	8.16	142.11	121.70
1	А	1183	ASN	CA-C-N	5.63	129.58	117.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	1183	ASN	Peptide,Mainchain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3564	0	3515	127	0
2	А	12	0	8	6	0
3	А	9	0	6	14	0
4	А	15	0	10	4	0
5	А	202	0	0	3	0
All	All	3802	0	3539	135	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1216:ASN:H	3:A:5293:GLU:HG3	1.20	1.02
1:A:980:MET:HA	1:A:989:SER:HB2	1.45	0.98
1:A:989:SER:HB3	1:A:1066:GLU:HA	1.50	0.93
1:A:1216:ASN:N	3:A:5293:GLU:HG3	1.84	0.92
1:A:1143:LYS:HZ3	4:A:5294:TRP:HH2	1.22	0.86

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	439/441 (100%)	403 (92%)	32 (7%)	4 (1%)	17 16



All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	945	ASP
1	А	989	SER
1	А	1184	ASN
1	А	990	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric (Outliers	Percentiles	
1	А	397/397~(100%)	382~(96%)	15 (4%)	33 42	

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	1072	LEU
1	А	1246	ARG
1	А	1097	ASN
1	А	1299	LEU
1	А	1208	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	А	1120	ASN
1	А	1141	GLN
1	А	1274	GLN
1	А	1203	ASN
1	А	1207	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
	Type	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLU	А	5293	4,2	7,8,9	0.85	0	4,9,11	1.19	0
2	TYR	А	5292	3	11,12,13	0.61	0	$12,\!15,\!17$	0.64	0
4	TRP	А	5294	3	14,16,16	0.88	0	16,22,22	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLU	А	5293	4,2	-	2/6/7/9	-
2	TYR	А	5292	3	-	3/5/6/8	0/1/1/1
4	TRP	А	5294	3	-	6/7/8/8	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	А	5292	TYR	O-C-CA-CB
3	А	5293	GLU	N-CA-CB-CG
3	А	5293	GLU	C-CA-CB-CG
4	А	5294	TRP	C-CA-CB-CG
2	А	5292	TYR	CA-CB-CG-CD2

There are no ring outliers.

3 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	5293	GLU	14	0
2	А	5292	TYR	6	0
4	А	5294	TRP	4	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	А	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1183:ASN	С	1184:ASN	N	1.16



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

