



Full wwPDB EM Validation Report ⓘ

Jul 7, 2024 – 08:13 am BST

PDB ID : 7YYN
EMDB ID : EMD-14384
Title : Mammalian Dicer in the dicing state with pre-miR-15a substrate
Authors : Zanova, M.; Zapletal, D.; Kubicek, K.; Stefl, R.; Pinkas, M.; Novacek, J.
Deposited on : 2022-02-18
Resolution : 6.21 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 6.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	59	
2	A	1765	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 13111 atoms, of which 6287 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 59-nt precursor of miR-15a.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			P
1	B	55	1763	523	590	205	390	55	0	0

- Molecule 2 is a protein called Isoform 2 of Endoribonuclease Dicer.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	A	705	11348	3643	5697	954	1026	28	0	0

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-51	MET	-	initiating methionine	UNP Q8R418
A	-50	VAL	-	expression tag	UNP Q8R418
A	-49	TRP	-	expression tag	UNP Q8R418
A	-48	SER	-	expression tag	UNP Q8R418
A	-47	HIS	-	expression tag	UNP Q8R418
A	-46	PRO	-	expression tag	UNP Q8R418
A	-45	GLN	-	expression tag	UNP Q8R418
A	-44	PHE	-	expression tag	UNP Q8R418
A	-43	GLU	-	expression tag	UNP Q8R418
A	-42	LYS	-	expression tag	UNP Q8R418
A	-41	GLY	-	expression tag	UNP Q8R418
A	-40	GLY	-	expression tag	UNP Q8R418
A	-39	GLY	-	expression tag	UNP Q8R418
A	-38	SER	-	expression tag	UNP Q8R418
A	-37	GLY	-	expression tag	UNP Q8R418
A	-36	GLY	-	expression tag	UNP Q8R418
A	-35	GLY	-	expression tag	UNP Q8R418
A	-34	SER	-	expression tag	UNP Q8R418
A	-33	GLY	-	expression tag	UNP Q8R418
A	-32	GLY	-	expression tag	UNP Q8R418
A	-31	SER	-	expression tag	UNP Q8R418

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-30	ALA	-	expression tag	UNP Q8R418
A	-29	TRP	-	expression tag	UNP Q8R418
A	-28	SER	-	expression tag	UNP Q8R418
A	-27	HIS	-	expression tag	UNP Q8R418
A	-26	PRO	-	expression tag	UNP Q8R418
A	-25	GLN	-	expression tag	UNP Q8R418
A	-24	PHE	-	expression tag	UNP Q8R418
A	-23	GLU	-	expression tag	UNP Q8R418
A	-22	LYS	-	expression tag	UNP Q8R418
A	-21	GLY	-	expression tag	UNP Q8R418
A	-20	ASP	-	expression tag	UNP Q8R418
A	-19	TYR	-	expression tag	UNP Q8R418
A	-18	PRO	-	expression tag	UNP Q8R418
A	-17	TYR	-	expression tag	UNP Q8R418
A	-16	ASP	-	expression tag	UNP Q8R418
A	-15	VAL	-	expression tag	UNP Q8R418
A	-14	PRO	-	expression tag	UNP Q8R418
A	-13	ASP	-	expression tag	UNP Q8R418
A	-12	TYR	-	expression tag	UNP Q8R418
A	-11	ALA	-	expression tag	UNP Q8R418
A	-10	GLY	-	expression tag	UNP Q8R418
A	-9	THR	-	expression tag	UNP Q8R418
A	-8	GLU	-	expression tag	UNP Q8R418
A	-7	ASN	-	expression tag	UNP Q8R418
A	-6	LEU	-	expression tag	UNP Q8R418
A	-5	TYR	-	expression tag	UNP Q8R418
A	-4	PHE	-	expression tag	UNP Q8R418
A	-3	GLN	-	expression tag	UNP Q8R418
A	-2	GLY	-	expression tag	UNP Q8R418
A	-1	LEU	-	expression tag	UNP Q8R418
A	0	VAL	-	expression tag	UNP Q8R418
A	1	ASP	-	expression tag	UNP Q8R418
A	872	SER	THR	conflict	UNP Q8R418
A	1322	ALA	GLU	engineered mutation	UNP Q8R418
A	1381	SER	ALA	conflict	UNP Q8R418
A	1569	ALA	GLU	engineered mutation	UNP Q8R418
A	1679	GLY	-	expression tag	UNP Q8R418
A	1680	ARG	-	expression tag	UNP Q8R418
A	1681	GLY	-	expression tag	UNP Q8R418
A	1682	GLU	-	expression tag	UNP Q8R418
A	1683	ASN	-	expression tag	UNP Q8R418
A	1684	LEU	-	expression tag	UNP Q8R418

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1685	TYR	-	expression tag	UNP Q8R418
A	1686	PHE	-	expression tag	UNP Q8R418
A	1687	GLN	-	expression tag	UNP Q8R418
A	1688	GLY	-	expression tag	UNP Q8R418
A	1689	ALA	-	expression tag	UNP Q8R418
A	1690	SER	-	expression tag	UNP Q8R418
A	1691	ASP	-	expression tag	UNP Q8R418
A	1692	TYR	-	expression tag	UNP Q8R418
A	1693	LYS	-	expression tag	UNP Q8R418
A	1694	ASP	-	expression tag	UNP Q8R418
A	1695	HIS	-	expression tag	UNP Q8R418
A	1696	ASP	-	expression tag	UNP Q8R418
A	1697	GLY	-	expression tag	UNP Q8R418
A	1698	ASP	-	expression tag	UNP Q8R418
A	1699	TYR	-	expression tag	UNP Q8R418
A	1700	LYS	-	expression tag	UNP Q8R418
A	1701	ASP	-	expression tag	UNP Q8R418
A	1702	HIS	-	expression tag	UNP Q8R418
A	1703	ASP	-	expression tag	UNP Q8R418
A	1704	GLY	-	expression tag	UNP Q8R418
A	1705	SER	-	expression tag	UNP Q8R418
A	1706	HIS	-	expression tag	UNP Q8R418
A	1707	HIS	-	expression tag	UNP Q8R418
A	1708	HIS	-	expression tag	UNP Q8R418
A	1709	HIS	-	expression tag	UNP Q8R418
A	1710	HIS	-	expression tag	UNP Q8R418
A	1711	HIS	-	expression tag	UNP Q8R418
A	1712	HIS	-	expression tag	UNP Q8R418
A	1713	HIS	-	expression tag	UNP Q8R418


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 59-nt precursor of miR-15a



- Molecule 2: Isoform 2 of Endoribonuclease Dicer



	E1615		
SER	M1616		
GLY	E1617		
ARG	P1618		
GLY	E1619		
GLU	T1620		
ASN	A1621		
LEU	K1622		
TYR	F1623		
PHE	S1624		
GLN	P1625		
GLY	A1626		
ALA	E1627		
SER	H1628		
ASP	L1629		
TYR	T1629		
ASP	Y1630		
ASP	D1631		
TYR	G1632		
LYS	ASP		
ASP	K1633		
HIS	V1634		
ASP	R1635		
GLY	V1636		
SER	T1637		
HIS	H1638		
HIS	E1639		
HIS	V1640		
HIS	G1641		
HIS	V1642		
HIS	K1643		
HIS	G1644		
HIS	K1645		
HIS	F1646		
HIS	K1647		
HIS	G1648		
HIS	V1649		
HIS	G1650		
HIS	R1651		
HIS	S1652		
HIS	Y1653		
HIS	R1654		
HIS	I1655		
HIS	A1659		
HIS	R1662		
HIS	R1663		
HIS	A1664		
HIS	L1665		
HIS	R1666		
HIS	S1667		
HIS	L1668		
HIS	K1669		
HIS	A1670		
HIS	M1671		
HIS	Q1672		
PRO	GLN		
VAL	VAL		
PRO	PRO		
ASN	ASN		

SER
GLY
ARG
GLU
ASN
LEU
TYR
PHE
GLY
GLY
ALA
SER
ASP
TYR
LYS
ASP
HIS
ASP
GLY
ASP
TYR
LYS
ASP
HIS
ASP
GLY
SER
HIS
HIS
HIS
HIS
HIS
HIS
HIS
HIS
HIS
HIS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	84697	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	55	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.298	Depositor
Minimum map value	-0.141	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.112	Depositor
Map size (Å)	317.952, 317.952, 317.952	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.828, 0.828, 0.828	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	1.49	6/1309 (0.5%)	2.22	82/2036 (4.0%)
2	A	0.69	0/5790	1.03	23/7851 (0.3%)
All	All	0.89	6/7099 (0.1%)	1.37	105/9887 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	17
2	A	0	10
All	All	0	27

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	52	C	C4-N4	-7.37	1.27	1.33
1	B	29	G	C2-N2	-7.25	1.27	1.34
1	B	8	A	C5-C4	-5.30	1.35	1.38
1	B	26	G	C2-N2	-5.08	1.29	1.34
1	B	29	G	N1-C2	-5.05	1.33	1.37
1	B	9	C	C4-N4	-5.00	1.29	1.33

All (105) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1	U	O4'-C1'-N1	14.24	119.60	108.20
1	B	40	G	P-O3'-C3'	9.83	131.50	119.70
1	B	52	C	N3-C4-C5	8.71	125.39	121.90
1	B	40	G	C4'-C3'-O3'	8.42	129.84	113.00
1	B	58	C	N3-C2-O2	-8.18	116.18	121.90
2	A	550	ARG	NE-CZ-NH1	8.15	124.38	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	738	TYR	CB-CG-CD2	-8.10	116.14	121.00
2	A	1076	ARG	NE-CZ-NH1	8.03	124.32	120.30
1	B	8	A	C5-C6-N1	7.83	121.62	117.70
1	B	5	A	C5-C6-N1	7.75	121.58	117.70
2	A	1607	ARG	NE-CZ-NH1	7.72	124.16	120.30
2	A	1328	TYR	CB-CG-CD1	-7.65	116.41	121.00
2	A	614	TYR	CB-CG-CD1	-7.64	116.41	121.00
1	B	39	A	C4-C5-C6	-7.46	113.27	117.00
1	B	1	U	N1-C2-N3	7.29	119.27	114.90
1	B	2	A	C5-C6-N1	7.22	121.31	117.70
1	B	39	A	C5-C6-N1	7.21	121.30	117.70
2	A	734	TYR	CB-CG-CD2	-7.16	116.70	121.00
1	B	41	G	C5'-C4'-O4'	7.04	117.55	109.10
1	B	24	A	N1-C6-N6	-7.02	114.39	118.60
1	B	5	A	C4-C5-C6	-6.98	113.51	117.00
1	B	10	A	C4-C5-C6	-6.88	113.56	117.00
1	B	24	A	C5-C6-N1	6.79	121.09	117.70
1	B	1	U	N3-C2-O2	-6.72	117.50	122.20
1	B	39	A	N1-C6-N6	-6.71	114.57	118.60
1	B	44	A	C5-C6-N1	6.69	121.04	117.70
1	B	59	A	C4'-C3'-C2'	-6.66	95.94	102.60
1	B	59	A	C3'-C2'-C1'	-6.64	96.18	101.50
1	B	8	A	C4-C5-C6	-6.58	113.71	117.00
2	A	1628	ARG	NE-CZ-NH1	6.57	123.58	120.30
1	B	59	A	C4-C5-C6	-6.53	113.74	117.00
1	B	12	A	C4-C5-C6	-6.48	113.76	117.00
1	B	10	A	C5-C6-N1	6.46	120.93	117.70
1	B	44	A	N1-C6-N6	-6.46	114.73	118.60
1	B	2	A	N1-C6-N6	-6.45	114.73	118.60
2	A	821	TYR	CB-CG-CD1	-6.45	117.13	121.00
1	B	9	C	N3-C4-C5	6.42	124.47	121.90
1	B	51	G	C4'-C3'-C2'	-6.30	96.30	102.60
2	A	562	ARG	NE-CZ-NH1	6.17	123.38	120.30
1	B	3	G	C5-C6-N1	6.16	114.58	111.50
1	B	36	U	O4'-C1'-N1	6.15	113.12	108.20
1	B	40	G	OP2-P-O3'	6.15	118.72	105.20
1	B	22	G	N1-C6-O6	-6.11	116.23	119.90
1	B	57	U	N3-C2-O2	-6.10	117.93	122.20
1	B	28	U	O4'-C1'-N1	6.09	113.07	108.20
2	A	1510	TYR	CB-CG-CD2	-6.09	117.35	121.00
1	B	15	G	C5-C6-N1	6.04	114.52	111.50
1	B	42	C	N3-C2-O2	-6.04	117.67	121.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	58	C	N1-C2-O2	5.94	122.46	118.90
2	A	1130	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	B	34	G	C5-C6-N1	5.87	114.43	111.50
1	B	10	A	N1-C6-N6	-5.86	115.09	118.60
1	B	25	U	N3-C2-O2	-5.81	118.13	122.20
1	B	19	U	O4'-C1'-N1	5.80	112.84	108.20
1	B	55	C	N3-C2-O2	-5.74	117.88	121.90
1	B	52	C	N1-C2-O2	5.69	122.31	118.90
2	A	676	TYR	CB-CG-CD1	-5.67	117.60	121.00
1	B	52	C	N3-C2-O2	-5.66	117.94	121.90
1	B	59	A	C5-C6-N1	5.65	120.53	117.70
1	B	2	A	C4-C5-C6	-5.65	114.17	117.00
1	B	46	A	C5-C6-N1	5.63	120.52	117.70
2	A	562	ARG	NE-CZ-NH2	-5.62	117.49	120.30
2	A	1076	ARG	CD-NE-CZ	5.60	131.45	123.60
1	B	58	C	N3-C4-C5	5.60	124.14	121.90
1	B	40	G	C5-C6-N1	5.59	114.29	111.50
1	B	21	U	N1-C2-N3	5.56	118.24	114.90
1	B	44	A	C4-C5-C6	-5.54	114.23	117.00
1	B	51	G	C5-C6-N1	5.54	114.27	111.50
1	B	23	G	N1-C6-O6	-5.53	116.58	119.90
1	B	20	G	N1-C6-O6	-5.52	116.59	119.90
2	A	733	TYR	CB-CG-CD2	-5.51	117.69	121.00
1	B	24	A	C4-C5-C6	-5.50	114.25	117.00
2	A	1431	TYR	CB-CG-CD1	-5.48	117.71	121.00
2	A	1147	TYR	CB-CG-CD2	-5.46	117.72	121.00
1	B	55	C	O4'-C1'-N1	5.44	112.56	108.20
2	A	1653	TYR	C-N-CA	5.42	135.25	121.70
2	A	508	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	13	A	C5-C6-N1	5.39	120.40	117.70
1	B	1	U	C5'-C4'-O4'	5.34	115.52	109.10
1	B	56	C	C5'-C4'-O4'	5.34	115.51	109.10
1	B	1	U	C5-C6-N1	-5.28	120.06	122.70
1	B	1	U	N1-C1'-C2'	-5.28	106.19	112.00
1	B	43	C	N3-C2-O2	-5.27	118.21	121.90
1	B	35	G	C5-C6-N1	5.26	114.13	111.50
1	B	57	U	N1-C2-N3	5.26	118.06	114.90
1	B	47	C	O4'-C1'-N1	5.23	112.39	108.20
1	B	25	U	O4'-C1'-N1	5.23	112.38	108.20
2	A	1459	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	B	20	G	C5-C6-N1	5.21	114.10	111.50
1	B	43	C	N3-C4-C5	5.16	123.96	121.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	U	O4'-C1'-N1	5.15	112.32	108.20
1	B	40	G	C2-N3-C4	5.14	114.47	111.90
1	B	15	G	N1-C6-O6	-5.13	116.82	119.90
1	B	50	U	O4'-C1'-N1	5.13	112.31	108.20
2	A	1119	TYR	CB-CG-CD2	-5.13	117.92	121.00
1	B	47	C	N3-C2-O2	-5.12	118.31	121.90
1	B	59	A	N1-C6-N6	-5.12	115.53	118.60
1	B	5	A	C5-N7-C8	-5.12	101.34	103.90
1	B	50	U	C5-C6-N1	-5.10	120.15	122.70
1	B	52	C	C5'-C4'-O4'	5.09	115.21	109.10
1	B	1	U	C1'-O4'-C4'	-5.04	105.87	109.90
2	A	1591	TYR	CB-CG-CD1	-5.02	117.99	121.00
1	B	22	G	O4'-C1'-N9	5.02	112.22	108.20
1	B	7	C	O4'-C1'-N1	5.00	112.20	108.20
1	B	16	G	C5'-C4'-C3'	-5.00	108.00	116.00

There are no chirality outliers.

All (27) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	1431	TYR	Sidechain
2	A	1510	TYR	Sidechain
2	A	1662	ARG	Sidechain
2	A	507	ARG	Sidechain
2	A	508	ARG	Sidechain
2	A	668	ARG	Sidechain
2	A	699	ARG	Sidechain
2	A	722	PHE	Sidechain
2	A	727	TYR	Sidechain
2	A	833	ARG	Sidechain
1	B	25	U	Sidechain
1	B	26	G	Sidechain
1	B	28	U	Sidechain
1	B	37	G	Sidechain
1	B	38	C	Sidechain
1	B	39	A	Sidechain
1	B	40	G	Sidechain
1	B	43	C	Sidechain
1	B	46	A	Sidechain
1	B	48	U	Sidechain
1	B	49	G	Sidechain
1	B	5	A	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	50	U	Sidechain
1	B	51	G	Sidechain
1	B	53	U	Sidechain
1	B	59	A	Sidechain
1	B	6	G	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1173	590	583	3	0
2	A	5651	5697	5696	10	0
All	All	6824	6287	6279	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:698:TYR:CD1	2:A:738:TYR:CE2	2.97	0.53
1:B:1:U:H5''	2:A:621:ARG:HH11	1.79	0.47
2:A:1090:THR:HG23	2:A:1462:PHE:CE2	2.51	0.45
1:B:18:U:H2'	1:B:19:U:C6	2.53	0.43
2:A:619:ILE:HD11	2:A:805:ILE:CD1	2.50	0.42
2:A:762:LEU:HD13	2:A:822:ARG:HH21	1.83	0.42
1:B:9:C:H2'	1:B:10:A:C8	2.54	0.42
2:A:1626:ALA:HB2	2:A:1653:TYR:CD1	2.55	0.42
2:A:734:TYR:CG	2:A:799:LEU:CD1	3.04	0.41
2:A:713:TYR:CE2	2:A:751:ASP:HB2	2.56	0.41
2:A:696:PRO:HB3	2:A:698:TYR:CE2	2.55	0.41
2:A:615:ILE:HA	2:A:619:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	693/1765 (39%)	684 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	625/1561 (40%)	621 (99%)	4 (1%)	86	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	706	ARG
2	A	1067	ASN
2	A	1082	ASP
2	A	1315	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	B	53/59 (89%)	9 (16%)	1 (1%)

All (9) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	B	16	G
1	B	25	U
1	B	26	G
1	B	27	U
1	B	28	U
1	B	35	G
1	B	40	G
1	B	41	G
1	B	53	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	B	40	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14384. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 196




Y Index: 182




Z Index: 217


6.3.2 Raw map



X Index: 186



Y Index: 177




Z Index: 226


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.112. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 51 nm^3 ; this corresponds to an approximate mass of 46 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.161 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.161 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.21	-	-
Author-provided FSC curve	8.27	9.74	8.42
Unmasked-calculated*	8.17	9.56	8.29


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 8.27 differs from the reported value 6.21 by more than 10 %

The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.17 differs from the reported value 6.21 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14384 and PDB model 7YYN. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.112 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.112).

9.4 Atom inclusion [i](#)



At the recommended contour level, 48% of all backbone atoms, 46% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.112) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	■ 0.4630	■ 0.1330
A	■ 0.4490	■ 0.1320
B	■ 0.5220	■ 0.1410

