



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 15, 2022 – 12:55 pm GMT

PDB ID : 7Z04
Title : 10 mM Rb⁺ soak of beryllium fluoride inhibited Na⁺,K⁺-ATPase, E2-BeFx (rigid body model)
Authors : Fruergaard, M.U.; Dach, I.; Andersen, J.L.; Ozol, M.; Shasavar, A.; Quistgaard, E.M.; Poulsen, H.; Fedosova, N.U.; Nissen, P.
Deposited on : 2022-02-22
Resolution : 7.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 7.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1004 (10.00-3.90)
Clashscore	141614	1069 (10.00-3.90)
Ramachandran outliers	138981	1002 (10.00-3.90)
Sidechain outliers	138945	1002 (10.00-3.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	1020	
1	C	1020	
2	B	302	
2	D	302	
3	E	64	
3	G	64	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium/potassium-transporting ATPase subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	996	7726	4922	1301	1456	47	0	0	0
1	C	996	7726	4922	1301	1456	47	0	0	0


- Molecule 2 is a protein called Sodium/potassium-transporting ATPase subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	289	2368	1534	386	435	13	0	0	0
2	D	289	2368	1534	386	435	13	0	0	0

- Molecule 3 is a protein called FXYP domain-containing ion transport regulator.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	G	32	255	174	37	44	0	0	0
3	E	32	255	174	37	44	0	0	0

- Molecule 4 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Be	F		
4	A	1	4	1	3	0	0
4	C	1	4	1	3	0	0


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0
5	C	1	1	1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Sodium/potassium-transporting ATPase subunit alpha-1







- Molecule 2: Sodium/potassium-transporting ATPase subunit beta-1

Chain B: 44% 44% 8%



- Molecule 2: Sodium/potassium-transporting ATPase subunit beta-1


Chain D: 41% 49% 5%



V301
K302
S303

- Molecule 3: FXYP domain-containing ion transport regulator

Chain G: 34% 14% 50%



- Molecule 3: FXYP domain-containing ion transport regulator

Chain E: 27% 23% 50%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.00Å 119.15Å 498.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 7.50 48.98 – 7.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (25.00-7.50) 99.9 (48.98-7.50)	Depositor EDS
R_{merge}	0.30	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 7.37Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.302 , 0.338 0.318 , 0.337	Depositor DCC
R_{free} test set	943 reflections (9.81%)	wwPDB-VP
Wilson B-factor (Å ²)	373.0	Xtrriage
Anisotropy	0.729	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.058 for k,h,-l	Xtrriage
F_o, F_c correlation	0.50	EDS
Total number of atoms	20708	wwPDB-VP
Average B, all atoms (Å ²)	480.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	6/7876 (0.1%)	0.80	13/10688 (0.1%)
1	C	0.65	6/7876 (0.1%)	0.78	12/10688 (0.1%)
2	B	0.67	2/2431 (0.1%)	0.79	1/3279 (0.0%)
2	D	0.64	0/2431	0.76	2/3279 (0.1%)
3	E	0.62	0/261	0.83	0/354
3	G	0.62	0/261	0.78	1/354 (0.3%)
All	All	0.66	14/21136 (0.1%)	0.79	29/28642 (0.1%)

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	325	VAL	C-N	8.50	1.50	1.34
1	A	325	VAL	C-N	8.42	1.50	1.34
2	B	159	CYS	CB-SG	7.75	1.95	1.82
2	B	282	ASN	CA-C	-6.52	1.35	1.52
1	C	63	GLU	CD-OE1	-6.30	1.18	1.25

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	125	LEU	CA-CB-CG	8.91	135.78	115.30
1	A	171	MET	CA-CB-CG	8.34	127.47	113.30
2	B	98	TYR	CA-CB-CG	7.77	128.17	113.40
1	A	73	LEU	CA-CB-CG	6.97	131.33	115.30
1	C	123	LEU	CA-CB-CG	-6.79	99.69	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7726	0	7778	647	0
1	C	7726	0	7778	734	0
2	B	2368	0	2338	237	0
2	D	2368	0	2338	175	0
3	E	255	0	259	11	0
3	G	255	0	259	13	0
4	A	4	0	0	1	0
4	C	4	0	0	1	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
All	All	20708	0	20750	1769	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 1769 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:300:PHE:CD1	1:C:317:LEU:HB2	1.14	1.61
1:A:96:LEU:CD1	1:A:285:ILE:HD13	1.34	1.58
2:D:102:VAL:CG1	2:D:169:TYR:CE2	1.87	1.54
1:C:302:LEU:HA	1:C:305:ILE:CG2	1.39	1.52
1:A:783:PHE:CE2	1:A:800:ILE:HD11	1.42	1.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	994/1020 (98%)	825 (83%)	144 (14%)	25 (2%)	5	32
1	C	994/1020 (98%)	822 (83%)	137 (14%)	35 (4%)	3	25
2	B	287/302 (95%)	217 (76%)	56 (20%)	14 (5%)	2	20
2	D	287/302 (95%)	231 (80%)	43 (15%)	13 (4%)	2	22
3	E	30/64 (47%)	26 (87%)	3 (10%)	1 (3%)	4	26
3	G	30/64 (47%)	26 (87%)	4 (13%)	0	100	100
All	All	2622/2772 (95%)	2147 (82%)	387 (15%)	88 (3%)	3	26

5 of 88 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	115	GLU
1	A	1010	VAL
2	B	160	SER
2	B	169	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	847/864 (98%)	831 (98%)	16 (2%)	57	75
1	C	847/864 (98%)	823 (97%)	24 (3%)	43	65
2	B	259/268 (97%)	250 (96%)	9 (4%)	36	59
2	D	259/268 (97%)	248 (96%)	11 (4%)	30	54
3	E	26/51 (51%)	26 (100%)	0	100	100
3	G	26/51 (51%)	26 (100%)	0	100	100
All	All	2264/2366 (96%)	2204 (97%)	60 (3%)	44	65

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	204	CYS
2	D	163	ASN
1	C	698	CYS
2	D	159	CYS
2	D	273	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	713	ASN
2	D	256	GLN
1	C	912	HIS
1	C	122	ASN
1	C	613	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BEF	C	1101	1	0,3,3	-	-	-		
4	BEF	A	1101	1	0,3,3	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1101	BEF	1	0
4	A	1101	BEF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates


Unable to reproduce the depositors R factor - this section is therefore empty.


6.4 Ligands


Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.