



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 1, 2021 – 06:11 pm BST

PDB ID : 6Z3N  
Title : Apo Structure of a Hydrolase from Pseudomonas aeruginosa PAO1  
Authors : Jin, Y.; Rizkallah, P.; Bell, H.  
Deposited on : 2020-05-21  
Resolution : 1.58 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.19  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.19


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	205	 4% 78% 8% 14%
1	BBB	205	 4% 79% 7% 14%
1	CCC	205	 3% 80% 5% 14%
1	DDD	205	 4% 77% 7% 14%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	AAA	177	1408	881	261	263	2	1	0	0	0
1	BBB	177	1417	886	263	265	2	1	0	1	0
1	CCC	177	1408	881	261	263	2	1	0	0	0
1	DDD	177	1408	881	261	263	2	1	0	0	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-20	MSE	-	initiating methionine	UNP Q9I686
AAA	-19	GLY	-	expression tag	UNP Q9I686
AAA	-18	SER	-	expression tag	UNP Q9I686
AAA	-17	SER	-	expression tag	UNP Q9I686
AAA	-16	HIS	-	expression tag	UNP Q9I686
AAA	-15	HIS	-	expression tag	UNP Q9I686
AAA	-14	HIS	-	expression tag	UNP Q9I686
AAA	-13	HIS	-	expression tag	UNP Q9I686
AAA	-12	HIS	-	expression tag	UNP Q9I686
AAA	-11	HIS	-	expression tag	UNP Q9I686
AAA	-10	SER	-	expression tag	UNP Q9I686
AAA	-9	SER	-	expression tag	UNP Q9I686
AAA	-8	GLY	-	expression tag	UNP Q9I686
AAA	-7	GLU	-	expression tag	UNP Q9I686
AAA	-6	ASN	-	expression tag	UNP Q9I686
AAA	-5	LEU	-	expression tag	UNP Q9I686
AAA	-4	TYR	-	expression tag	UNP Q9I686
AAA	-3	PHE	-	expression tag	UNP Q9I686
AAA	-2	GLN	-	expression tag	UNP Q9I686
AAA	-1	SER	-	expression tag	UNP Q9I686
AAA	0	HIS	-	expression tag	UNP Q9I686

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	1	MSE	-	expression tag	UNP Q9I686
BBB	-20	MSE	-	initiating methionine	UNP Q9I686
BBB	-19	GLY	-	expression tag	UNP Q9I686
BBB	-18	SER	-	expression tag	UNP Q9I686
BBB	-17	SER	-	expression tag	UNP Q9I686
BBB	-16	HIS	-	expression tag	UNP Q9I686
BBB	-15	HIS	-	expression tag	UNP Q9I686
BBB	-14	HIS	-	expression tag	UNP Q9I686
BBB	-13	HIS	-	expression tag	UNP Q9I686
BBB	-12	HIS	-	expression tag	UNP Q9I686
BBB	-11	HIS	-	expression tag	UNP Q9I686
BBB	-10	SER	-	expression tag	UNP Q9I686
BBB	-9	SER	-	expression tag	UNP Q9I686
BBB	-8	GLY	-	expression tag	UNP Q9I686
BBB	-7	GLU	-	expression tag	UNP Q9I686
BBB	-6	ASN	-	expression tag	UNP Q9I686
BBB	-5	LEU	-	expression tag	UNP Q9I686
BBB	-4	TYR	-	expression tag	UNP Q9I686
BBB	-3	PHE	-	expression tag	UNP Q9I686
BBB	-2	GLN	-	expression tag	UNP Q9I686
BBB	-1	SER	-	expression tag	UNP Q9I686
BBB	0	HIS	-	expression tag	UNP Q9I686
BBB	1	MSE	-	expression tag	UNP Q9I686
CCC	-20	MSE	-	initiating methionine	UNP Q9I686
CCC	-19	GLY	-	expression tag	UNP Q9I686
CCC	-18	SER	-	expression tag	UNP Q9I686
CCC	-17	SER	-	expression tag	UNP Q9I686
CCC	-16	HIS	-	expression tag	UNP Q9I686
CCC	-15	HIS	-	expression tag	UNP Q9I686
CCC	-14	HIS	-	expression tag	UNP Q9I686
CCC	-13	HIS	-	expression tag	UNP Q9I686
CCC	-12	HIS	-	expression tag	UNP Q9I686
CCC	-11	HIS	-	expression tag	UNP Q9I686
CCC	-10	SER	-	expression tag	UNP Q9I686
CCC	-9	SER	-	expression tag	UNP Q9I686
CCC	-8	GLY	-	expression tag	UNP Q9I686
CCC	-7	GLU	-	expression tag	UNP Q9I686
CCC	-6	ASN	-	expression tag	UNP Q9I686
CCC	-5	LEU	-	expression tag	UNP Q9I686
CCC	-4	TYR	-	expression tag	UNP Q9I686
CCC	-3	PHE	-	expression tag	UNP Q9I686
CCC	-2	GLN	-	expression tag	UNP Q9I686

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
CCC	-1	SER	-	expression tag	UNP Q9I686
CCC	0	HIS	-	expression tag	UNP Q9I686
CCC	1	MSE	-	expression tag	UNP Q9I686
DDD	-20	MSE	-	initiating methionine	UNP Q9I686
DDD	-19	GLY	-	expression tag	UNP Q9I686
DDD	-18	SER	-	expression tag	UNP Q9I686
DDD	-17	SER	-	expression tag	UNP Q9I686
DDD	-16	HIS	-	expression tag	UNP Q9I686
DDD	-15	HIS	-	expression tag	UNP Q9I686
DDD	-14	HIS	-	expression tag	UNP Q9I686
DDD	-13	HIS	-	expression tag	UNP Q9I686
DDD	-12	HIS	-	expression tag	UNP Q9I686
DDD	-11	HIS	-	expression tag	UNP Q9I686
DDD	-10	SER	-	expression tag	UNP Q9I686
DDD	-9	SER	-	expression tag	UNP Q9I686
DDD	-8	GLY	-	expression tag	UNP Q9I686
DDD	-7	GLU	-	expression tag	UNP Q9I686
DDD	-6	ASN	-	expression tag	UNP Q9I686
DDD	-5	LEU	-	expression tag	UNP Q9I686
DDD	-4	TYR	-	expression tag	UNP Q9I686
DDD	-3	PHE	-	expression tag	UNP Q9I686
DDD	-2	GLN	-	expression tag	UNP Q9I686
DDD	-1	SER	-	expression tag	UNP Q9I686
DDD	0	HIS	-	expression tag	UNP Q9I686
DDD	1	MSE	-	expression tag	UNP Q9I686

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total C O 4 2 2	0	0
2	CCC	1	Total C O 4 2 2	0	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Ni 1 1	0	0
3	BBB	1	Total Ni 1 1	0	0
3	CCC	1	Total Ni 1 1	0	0
3	DDD	1	Total Ni 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	CCC	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	AAA	153	Total 153	O 153	0	0
5	BBB	124	Total 124	O 124	0	0
5	CCC	145	Total 147	O 147	0	2
5	DDD	144	Total 145	O 145	0	1


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydrolase




- Molecule 1: Hydrolase



- Molecule 1: Hydrolase



- Molecule 1: Hydrolase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.31Å 85.90Å 122.64Å 90.00° 102.34° 90.00°	Depositor
Resolution (Å)	59.90 – 1.58 59.90 – 1.58	Depositor EDS
% Data completeness (in resolution range)	81.2 (59.90-1.58) 81.2 (59.90-1.58)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 1.58Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.137 , 0.200 0.140 , 0.201	Depositor DCC
$R_{free}$ test set	4868 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.8	Xtrriage
Anisotropy	0.103	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6228	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.77	1/1438 (0.1%)	0.86	1/1951 (0.1%)
1	BBB	0.80	0/1447	0.88	2/1963 (0.1%)
1	CCC	0.80	0/1438	0.88	2/1951 (0.1%)
1	DDD	0.92	5/1438 (0.3%)	0.86	1/1951 (0.1%)
All	All	0.83	6/5761 (0.1%)	0.87	6/7816 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	DDD	81	GLU	CD-OE1	11.63	1.38	1.25
1	DDD	81	GLU	CD-OE2	9.83	1.36	1.25
1	DDD	47	GLU	CD-OE1	8.92	1.35	1.25
1	DDD	47	GLU	CD-OE2	5.64	1.31	1.25
1	AAA	47	GLU	CD-OE1	5.08	1.31	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	43	MSE	CG-SE-CE	7.38	115.14	98.90
1	CCC	43	MSE	CG-SE-CE	5.93	111.95	98.90
1	AAA	27	ASN	CB-CA-C	5.80	122.00	110.40
1	DDD	48	LEU	CA-CB-CG	-5.68	102.24	115.30
1	CCC	132	ARG	NE-CZ-NH2	5.29	122.94	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1408	0	1375	6	0
1	BBB	1417	0	1382	6	0
1	CCC	1408	0	1375	8	0
1	DDD	1408	0	1375	10	0
2	AAA	4	0	3	0	0
2	CCC	4	0	3	0	0
3	AAA	1	0	0	0	0
3	BBB	1	0	0	0	0
3	CCC	1	0	0	0	0
3	DDD	1	0	0	0	0
4	CCC	6	0	8	3	0
5	AAA	153	0	0	1	0
5	BBB	124	0	0	0	0
5	CCC	147	0	0	1	0
5	DDD	145	0	0	0	0
All	All	6228	0	5521	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:DDD:101:ARG:HD3	1:DDD:106:PRO:C	2.21	0.61
1:DDD:102:VAL:HG13	1:DDD:103:GLY:N	2.15	0.61
1:BBB:39:VAL:HG12	1:BBB:43:MSE:HE2	1.86	0.56
1:BBB:103:GLY:C	1:BBB:105:GLY:H	2.11	0.54
1:CCC:104:ASP:O	1:CCC:105:GLY:O	2.27	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	175/205 (85%)	170 (97%)	4 (2%)	1 (1%)	25	8
1	BBB	176/205 (86%)	170 (97%)	4 (2%)	2 (1%)	14	3
1	CCC	175/205 (85%)	169 (97%)	4 (2%)	2 (1%)	14	3
1	DDD	175/205 (85%)	167 (95%)	7 (4%)	1 (1%)	25	8
All	All	701/820 (86%)	676 (96%)	19 (3%)	6 (1%)	17	4

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	103	GLY
1	CCC	105	GLY
1	BBB	104	ASP
1	BBB	102	VAL
1	CCC	104	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	144/165 (87%)	138 (96%)	6 (4%)	30	7
1	BBB	145/165 (88%)	143 (99%)	2 (1%)	67	45
1	CCC	144/165 (87%)	140 (97%)	4 (3%)	43	17
1	DDD	144/165 (87%)	136 (94%)	8 (6%)	21	4
All	All	577/660 (87%)	557 (96%)	20 (4%)	36	11

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	DDD	31	SER
1	DDD	101	ARG
1	DDD	137	GLN
1	DDD	107	LYS
1	BBB	30	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	CCC	501	-	5,5,5	0.55	0	5,5,5	0.38	0
2	ACT	AAA	501	-	1,3,3	5.43	1 (100%)	0,3,3	0.00	-
2	ACT	CCC	502	-	1,3,3	4.47	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	CCC	501	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	501	ACT	CH3-C	5.43	1.55	1.48
2	CCC	502	ACT	CH3-C	4.47	1.54	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	CCC	501	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	176/205 (85%)	0.38	8 (4%) 33 33	20, 30, 73, 113	0
1	BBB	176/205 (85%)	0.52	9 (5%) 28 28	20, 29, 67, 127	0
1	CCC	176/205 (85%)	0.38	6 (3%) 45 46	20, 30, 64, 144	3 (1%)
1	DDD	176/205 (85%)	0.31	8 (4%) 33 33	20, 30, 73, 108	2 (1%)
All	All	704/820 (85%)	0.40	31 (4%) 34 34	20, 30, 70, 144	5 (0%)

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	CCC	103	GLY	13.8
1	BBB	102	VAL	13.6
1	BBB	106	PRO	13.1
1	BBB	103	GLY	10.6
1	BBB	104	ASP	10.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	CCC	501	6/6	0.91	0.27	69,80,86,110	0
3	NI	DDD	201	1/1	0.93	0.10	61,61,61,61	1
2	ACT	CCC	502	4/4	0.95	0.07	66,66,69,71	0
2	ACT	AAA	501	4/4	0.96	0.16	56,61,66,68	0
3	NI	CCC	503	1/1	0.97	0.14	63,63,63,63	1
3	NI	AAA	502	1/1	0.98	0.07	56,56,56,56	1
3	NI	BBB	201	1/1	0.99	0.09	56,56,56,56	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.