



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2023 – 01:08 pm GMT

PDB ID : 7Z3R
Title : Crystal structure of the mouse leptin:LepR-IgCRH2 complex to 2.95 Å resolution.
Authors : Verstraete, K.; Verschueren, K.; Savvides, S.N.; Tsigotaki, A.
Deposited on : 2022-03-02
Resolution : 2.95 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	
2	B	330	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3563 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leptin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1089	685	184	217	3	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP P41160

- Molecule 2 is a protein called Leptin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	306	2453	1570	418	451	14	0	0	0

There are 25 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	304	ALA	-	expression tag	UNP P48356
B	305	HIS	-	expression tag	UNP P48356
B	306	HIS	-	expression tag	UNP P48356
B	307	HIS	-	expression tag	UNP P48356
B	308	HIS	-	expression tag	UNP P48356
B	309	HIS	-	expression tag	UNP P48356
B	310	HIS	-	expression tag	UNP P48356
B	311	PRO	-	expression tag	UNP P48356
B	312	GLY	-	expression tag	UNP P48356
B	313	GLY	-	expression tag	UNP P48356
B	314	PRO	-	expression tag	UNP P48356
B	315	GLY	-	expression tag	UNP P48356
B	316	SER	-	expression tag	UNP P48356
B	317	GLU	-	expression tag	UNP P48356
B	318	ASN	-	expression tag	UNP P48356

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	319	LEU	-	expression tag	UNP P48356
B	320	TYR	-	expression tag	UNP P48356
B	321	PHE	-	expression tag	UNP P48356
B	322	GLN	-	expression tag	UNP P48356
B	323	GLY	-	expression tag	UNP P48356
B	324	GLY	-	expression tag	UNP P48356
B	325	SER	-	expression tag	UNP P48356
B	326	SER	-	expression tag	UNP P48356
B	327	GLY	-	expression tag	UNP P48356
B	602	SER	CYS	engineered mutation	UNP P48356

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	B	1	14	8	1	5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	B	7	7	7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leptin



- Molecule 2: Leptin receptor



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	132.68Å 132.68Å 249.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.34 – 2.95 66.34 – 2.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (66.34-2.95) 100.0 (66.34-2.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.96Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (3-FEB-2022)	Depositor
R, R_{free}	0.223 , 0.251 0.210 , 0.242	Depositor DCC
R_{free} test set	1194 reflections (6.61%)	wwPDB-VP
Wilson B-factor (Å ²)	101.5	Xtrriage
Anisotropy	0.064	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 82.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3563	wwPDB-VP
Average B, all atoms (Å ²)	111.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/1101	0.51	0/1495
2	B	0.28	0/2520	0.53	0/3434
All	All	0.28	0/3621	0.52	0/4929

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1089	0	1126	8	0
2	B	2453	0	2421	9	0
3	B	14	0	13	0	0
4	B	7	0	0	0	0
All	All	3563	0	3560	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:349:HIS:CD2	2:B:382:VAL:HG21	2.31	0.66
1:A:84:GLN:HE22	1:A:127:LYS:NZ	1.94	0.65
1:A:114:SER:O	1:A:115:LYS:HG2	1.99	0.63
2:B:458:SER:HB2	2:B:515:HIS:HA	1.85	0.59
2:B:334:PHE:HB2	2:B:349:HIS:HB2	1.89	0.54
1:A:111:LEU:O	1:A:114:SER:O	2.25	0.53
1:A:84:GLN:HE22	1:A:127:LYS:HZ2	1.56	0.52
2:B:352:TYR:HB3	2:B:359:ILE:HG12	1.93	0.51
2:B:446:ARG:HG3	2:B:495:GLU:HG2	1.93	0.50
2:B:337:LYS:HB3	2:B:517:LEU:HD21	1.95	0.48
2:B:562:PRO:HB2	2:B:564:ASN:HD22	1.78	0.48
1:A:91:SER:HB3	1:A:94:VAL:HG22	1.97	0.47
1:A:64:PRO:O	1:A:82:TYR:OH	2.31	0.42
1:A:84:GLN:HE22	1:A:127:LYS:HZ3	1.66	0.41
2:B:351:ILE:HD11	2:B:386:VAL:HG22	2.02	0.41
2:B:564:ASN:HD22	2:B:564:ASN:H	1.69	0.40
1:A:127:LYS:HE3	1:A:127:LYS:HB3	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/148 (93%)	127 (93%)	7 (5%)	3 (2%)	6	28
2	B	304/330 (92%)	296 (97%)	8 (3%)	0	100	100
All	All	441/478 (92%)	423 (96%)	15 (3%)	3 (1%)	22	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	LYS
1	A	65	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	69	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	131/136 (96%)	126 (96%)	5 (4%)	33 66
2	B	282/300 (94%)	276 (98%)	6 (2%)	53 80
All	All	413/436 (95%)	402 (97%)	11 (3%)	44 74

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LEU
1	A	116	SER
1	A	127	LYS
1	A	162	ASP
1	A	166	GLU
2	B	357	GLN
2	B	414	GLN
2	B	446	ARG
2	B	455	LEU
2	B	458	SER
2	B	574	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	49	GLN
1	A	84	GLN
1	A	109	HIS
1	A	151	GLN
2	B	329	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	349	HIS
2	B	564	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	701	2	14,14,15	0.26	0	17,19,21	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	701	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/148 (95%)	1.23	27 (19%) 1 0	82, 114, 170, 187	0
2	B	306/330 (92%)	1.14	48 (15%) 2 1	78, 108, 140, 159	0
All	All	447/478 (93%)	1.17	75 (16%) 1 1	78, 110, 157, 187	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	452	ILE	4.9
1	A	72	LEU	4.5
2	B	601	LEU	4.3
2	B	631	VAL	4.1
2	B	632	MET	3.9
2	B	511	ILE	3.9
1	A	54	LYS	3.6
2	B	462	LEU	3.5
2	B	552	LEU	3.5
2	B	351	ILE	3.4
1	A	134	VAL	3.3
1	A	92	GLN	3.2
1	A	110	LEU	3.1
1	A	131	LEU	3.0
1	A	97	ILE	3.0
2	B	426	ILE	3.0
2	B	596	LEU	2.9
2	B	449	PRO	2.7
1	A	112	ALA	2.7
1	A	24	ILE	2.7
2	B	536	LEU	2.7
1	A	139	LEU	2.6
2	B	630	LEU	2.6
1	A	119	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	GLN	2.5
2	B	398	ARG	2.5
2	B	430	ILE	2.5
2	B	370	LEU	2.5
2	B	541	VAL	2.5
2	B	503	LEU	2.5
1	A	100	ASP	2.5
1	A	70	LEU	2.5
1	A	135	LEU	2.5
2	B	352	TYR	2.5
1	A	79	LEU	2.4
1	A	143	GLU	2.4
2	B	562	PRO	2.4
2	B	598	VAL	2.4
2	B	423	LEU	2.4
2	B	402	LYS	2.4
2	B	527	VAL	2.3
1	A	82	TYR	2.3
2	B	498	PHE	2.3
1	A	60	LEU	2.3
2	B	617	LEU	2.3
2	B	609	VAL	2.3
2	B	432	ILE	2.2
2	B	587	PHE	2.2
1	A	94	VAL	2.2
1	A	147	LEU	2.2
2	B	517	LEU	2.2
2	B	579	ILE	2.2
1	A	28	GLN	2.2
2	B	374	ILE	2.2
2	B	501	ILE	2.2
2	B	607	VAL	2.2
1	A	145	VAL	2.1
1	A	42	ILE	2.1
2	B	529	PRO	2.1
2	B	408	VAL	2.1
2	B	358	ILE	2.1
2	B	477	ILE	2.1
2	B	597	LEU	2.1
2	B	359	ILE	2.1
1	A	101	LEU	2.1
2	B	542	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	364	ILE	2.0
2	B	570	ILE	2.0
1	A	27	VAL	2.0
2	B	532	VAL	2.0
2	B	545	ILE	2.0
2	B	494	TYR	2.0
1	A	165	PRO	2.0
2	B	394	LEU	2.0
2	B	331	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	701	14/15	0.70	0.21	155,155,155,155	0

6.5 Other polymers [i](#)

There are no such residues in this entry.