



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2020 – 09:05 AM GMT

PDB ID : 6Z4X  
Title : Structure of the CAK complex form Chaetomium thermophilum bound to ATP-gamma-S  
Authors : Peissert, S.; Kuper, J.; Kisker, C.  
Deposited on : 2020-05-26  
Resolution : 2.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	 3% 55% 18% 24%
1	D	425	 2% 55% 19% 24%
2	B	437	 2% 55% 18% 25%
2	E	437	 0% 58% 15% 27%
3	C	69	 3% 75% 17% 5%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	69	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into segments: a small red segment at the start, followed by a large green segment labeled '74%', then a yellow segment labeled '22%', and finally a small grey segment at the end. A '%' symbol is positioned above the red segment, and two dots are positioned below the grey segment.</p>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11470 atoms, of which 28 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLIN domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	312	2522	1599	449	463	11	0	0	0
1	D	314	2533	1605	451	466	11	0	0	0


- Molecule 2 is a protein called Protein kinase domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	322	2613	1678	462	460	13	0	0	0
2	E	323	2620	1682	463	462	13	0	0	0

- Molecule 3 is a protein called RING-type domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	66	538	344	90	102	2	0	0	0
3	F	67	550	353	91	104	2	0	0	0

- Molecule 4 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter code: AGS) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	H	N	O	P			S
4	B	1	45	10	14	5	12	3	1	0	0
4	E	1	45	10	14	5	12	3	1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	B	1	1	1	0	0
5	E	1	1	1	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
6	B	1	1	1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	B	1	1	1	0	0



VAL  
PHE  
GLY  
VAL  
PRO  
LEU  
LYS  
LYS  
LYS  
GLN  
LEU  
LEU  
GLU  
ASP  
PRO  
PHE  
GLY  
PRO  
PRO  
PRO  
LEU  
ALA  
GLY

• Molecule 2: Protein kinase domain-containing protein



MET  
ALA  
ALA  
SER  
PRO  
LEU  
ILE  
ALA  
PRO  
THR  
THR  
ASP  
ALA  
LEU  
GLN  
GLN  
SER  
ARG  
PRO  
GLY  
SER  
THR  
SER  
SER  
PRO  
PHE  
LYS  
LYS  
ARG  
GLN  
PRO  
GLN  
GLN  
PRO  
SER  
GLN  
PRO  
PRO  
SER  
SER  
THR  
ALA  
PRO  
SER  
SER  
SER  
ASN  
ALA  
ASN  
ASP  
THR  
PRO  
GLN  
LEU  
LEU  
GLY  
SER  
ARG  
PRO  
ASN  
PRO  
ILE

THR  
ILE  
ALA  
ASN  
VAL  
THR  
SER  
ALA  
PRO  
PHE  
SER  
SER  
ARG  
PRO  
ALA  
ILE  
THR  
P78  
E82  
Q83  
R90  
K97  
L98  
G99  
E100  
G101  
T102  
V119  
I124  
Q127  
A128  
Q129  
D132  
D137  
A138  
V139  
L142  
H151  
L157  
I158  
S159  
V160  
D165  
M167  
L168  
L172

L175  
P176  
L177  
L180  
E181  
L184  
Y192  
K198  
A199  
W200  
L204  
W210  
E213  
H218  
R219  
D220  
L221  
K222  
P223  
N224  
M225  
L226  
L227  
L228  
A229  
D238  
L241  
A246  
R251  
M252  
T253  
A254  
M255  
V256  
I257  
T258  
R259  
W260  
R271  
G274  
V277  
D278  
W280

F286  
L289  
R292  
L296  
F297  
G298  
L307  
L312  
L326  
D331  
P332  
F349  
V352  
L359  
L360  
Q365  
K366  
D367  
P368  
K369  
K370  
L372  
R375  
E376  
R377  
W382  
T385  
L395  
P396  
S399  
GLY  
GLY  
ASP  
LEU  
MET  
GLY  
ALA  
ASP  
LEU  
LYS  
ASN  
ARG

ARG  
PRO  
GLY  
MET  
VAL  
GLU  
ASP  
GLU  
ASN  
PRO  
THR  
PRO  
ARG  
GLY  
SER  
VAL  
LYS  
ARG  
ALA  
LYS  
LEU  
ASP  
PHE  
GLY  
GLN  
LEU  
LYS

• Molecule 2: Protein kinase domain-containing protein



MET  
ALA  
ALA  
SER  
PRO  
LEU  
ILE  
ALA  
PRO  
THR  
THR  
ASP  
ALA  
LEU  
GLN  
GLN  
SER  
ARG  
PRO  
GLY  
SER  
THR  
SER  
SER  
PRO  
PHE  
LYS  
ARG  
GLN  
PRO  
GLN  
PRO  
SER  
SER  
THR  
ALA  
PRO  
SER  
SER  
SER  
ASN  
ALA  
ASN  
ASP  
THR  
PRO  
GLN  
GLY  
GLY  
SER  
ARG  
PRO  
ASN  
ILE


THR  
ILE  
ALA  
ASN  
VAL  
THR  
SER  
ALA  
PRO  
PHE  
SER  
SER  
ARG  
PRO  
ALA  
ILE  
T77  
F78  
D79  
P80  
R81  
E82  
Q83  
R90  
R96  
K97  
L98  
M99  
E100  
G101  
T102  
A103  
A104  
R112  
V119  
K122  
K123  
I124  
Q129  
D132  
D137  
A138  
V139  
L142  
M153  
L157  
V160

D165  
Q166  
M167  
L168  
V171  
L172  
L175  
E181  
I184  
Y192  
G193  
I197  
L204  
A207  
R219  
K222  
F223  
M224  
M225  
L226  
L227  
A246  
R251  
M252  
T253  
A254  
M255  
V256  
P263  
E264  
E265  
G269  
A270  
R271  
G274  
V277  
F286  
T290  
I291  
R292

F295  
L296  
L326  
D331  
P332  
F349  
G350  
A351  
L359  
L360  
Q365  
L366  
D367  
P368  
K369  
K370  
M377  
S399  
GLY  
GLY  
GLU  
GLU  
ASP  
LYS  
MET  
GLY  
ALA  
ALA  
LEU  
LEU  
LYS  
ARG  
ARG  
ARG  
PRO  
GLY  
MET  
VAL  
GLU  
ASP  
GLU  
ASN  
GLY  
PRO  
ARG  
GLY  
SER  
LYS  
VAL  
ALA  
ALA  
LYS  
LEU  
ASP  
PHE  
GLY


GLN  
LEU  
LYS

• Molecule 3: RING-type domain-containing protein



GLY  
PRO  
TYR  
D273  
R286  
V287  
R288  
L296  
D297  
K298  
Y299  
R300  
L306  
L313  
A323  
F324  
A325  
G326  
L327  
A328  
V329  
F330  
L331  
A337  
G338

• Molecule 3: RING-type domain-containing protein







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.97Å 84.90Å 154.48Å 90.00° 96.20° 90.00°	Depositor
Resolution (Å)	46.65 – 2.98 46.65 – 2.98	Depositor EDS
% Data completeness (in resolution range)	61.1 (46.65-2.98) 61.1 (46.65-2.98)	Depositor EDS
$R_{merge}$	0.28	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.47 (at 2.96Å)	Xtrriage
Refinement program	BUSTER, PHENIX 1.18rc4_3812	Depositor
R, $R_{free}$	0.203 , 0.246 0.208 , 0.248	Depositor DCC
$R_{free}$ test set	1469 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.9	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 39.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	11470	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AGS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2574	0.39	0/3471
1	D	0.25	0/2585	0.39	0/3486
2	B	0.24	0/2684	0.40	0/3635
2	E	0.24	0/2691	0.41	0/3646
3	C	0.25	0/552	0.38	0/742
3	F	0.26	0/565	0.39	0/760
All	All	0.25	0/11651	0.40	0/15740

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2522	0	2517	55	0
1	D	2533	0	2527	57	0
2	B	2613	0	2624	50	0
2	E	2620	0	2630	44	0
3	C	538	0	502	11	0
3	F	550	0	511	13	0
4	B	31	14	12	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	31	14	12	1	0
5	B	1	0	0	0	0
5	E	1	0	0	0	0
6	B	1	0	0	0	0
7	B	1	0	0	0	0
All	All	11442	28	11335	207	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:253:THR:HG22	2:E:255:ASN:H	1.28	0.96
2:B:253:THR:HG22	2:B:255:ASN:H	1.34	0.92
1:A:381:ARG:HD3	1:A:385:ASP:HA	1.56	0.86
3:C:313:LEU:H	3:C:313:LEU:HD22	1.48	0.77
1:A:375:ARG:NH2	2:B:82:GLU:OE2	2.19	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/425 (72%)	297 (97%)	9 (3%)	0	100	100
1	D	308/425 (72%)	298 (97%)	10 (3%)	0	100	100
2	B	320/437 (73%)	311 (97%)	8 (2%)	1 (0%)	41	74
2	E	321/437 (74%)	312 (97%)	8 (2%)	1 (0%)	41	74
3	C	64/69 (93%)	64 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	F	65/69 (94%)	65 (100%)	0	0	100	100
All	All	1384/1862 (74%)	1347 (97%)	35 (2%)	2 (0%)	51	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	257	ILE
2	E	257	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/354 (75%)	264 (99%)	3 (1%)	73	90
1	D	268/354 (76%)	265 (99%)	3 (1%)	73	90
2	B	275/370 (74%)	270 (98%)	5 (2%)	59	83
2	E	276/370 (75%)	270 (98%)	6 (2%)	52	80
3	C	55/57 (96%)	51 (93%)	4 (7%)	14	42
3	F	56/57 (98%)	53 (95%)	3 (5%)	22	55
All	All	1197/1562 (77%)	1173 (98%)	24 (2%)	55	81

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	313	LEU
1	D	185	VAL
3	F	288	ARG
3	C	327	LEU
1	D	23	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	372	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	AGS	B	501	5	26,33,33	1.94	4 (15%)	26,52,52	1.47	4 (15%)
4	AGS	E	501	-	26,33,33	1.94	4 (15%)	26,52,52	1.60	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AGS	B	501	5	-	4/17/38/38	0/3/3/3
4	AGS	E	501	-	-	2/17/38/38	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	501	AGS	PG-S1G	8.11	2.08	1.90
4	E	501	AGS	PG-S1G	8.00	2.08	1.90
4	E	501	AGS	O4'-C1'	2.84	1.45	1.41
4	B	501	AGS	C5-C4	2.51	1.47	1.40
4	E	501	AGS	C5-C4	2.47	1.47	1.40

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	501	AGS	N3-C2-N1	-3.96	122.49	128.68
4	B	501	AGS	N3-C2-N1	-3.82	122.71	128.68
4	E	501	AGS	C2'-C3'-C4'	3.38	109.20	102.64
4	E	501	AGS	PA-O3A-PB	-2.95	122.71	132.83
4	B	501	AGS	PA-O3A-PB	-2.92	122.80	132.83

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	501	AGS	PB-O3B-PG-O2G
4	E	501	AGS	PB-O3B-PG-O2G
4	E	501	AGS	PB-O3B-PG-O3G
4	B	501	AGS	PB-O3A-PA-O1A
4	B	501	AGS	PG-O3B-PB-O2B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	501	AGS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	312/425 (73%)	0.06	12 (3%) 40 24	31, 65, 100, 127	0
1	D	314/425 (73%)	-0.24	7 (2%) 62 42	14, 40, 81, 124	0
2	B	322/437 (73%)	-0.26	8 (2%) 57 38	18, 40, 86, 110	0
2	E	323/437 (73%)	-0.35	6 (1%) 66 46	13, 31, 79, 124	0
3	C	66/69 (95%)	-0.14	2 (3%) 50 31	38, 61, 84, 94	0
3	F	67/69 (97%)	-0.37	1 (1%) 73 54	17, 37, 67, 96	0
All	All	1404/1862 (75%)	-0.21	36 (2%) 56 36	13, 44, 88, 127	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	TYR	5.4
3	C	338	GLY	5.2
1	D	75	PRO	4.5
3	F	338	GLY	4.3
1	D	262	TYR	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MG	E	502	1/1	0.90	0.12	36,36,36,36	0
4	AGS	E	501	31/31	0.91	0.23	35,77,136,146	0
4	AGS	B	501	31/31	0.91	0.20	29,76,118,144	0
5	MG	B	502	1/1	0.95	0.11	19,19,19,19	0
6	CL	B	503	1/1	0.98	0.12	43,43,43,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.